

Labelling people as 'High Risk': A tyranny of eminence?

Teppo L N Järvinen

Whenever a doctor cannot do good, he must be kept from doing harm.

Hippocrates

Nowadays, being at 'high risk' of having a disease has become a disease in and of itself. Sweeping educational programmes at all levels of healthcare now turn an otherwise healthy person's 'high' blood pressure, elevated serum lipids or low bone density into chronic conditions having increased risk of a potentially bad event.¹ But what represents 'high risk'? This question lies at the heart of modern medicine, particularly with respect to pharmacological primary prevention.

Advocates of this evolution argue simply that primary prevention saves lives. However, permissive labelling of conditions as diseases may not be entirely harmless. On an individual patient level, possible disadvantages include making relatively healthy individuals perceive themselves as 'sick', and almost every treatment has inherent risks.² On a societal level, we probably all still remember the discouraging effect of the new European guidelines on cardiovascular disease classified most adult Norwegians—among the healthiest populations in the world—to be at 'high risk' of cardiovascular disease.³ If these guidelines were applied to the Norwegian healthcare system, the focus on hypertension would drain the entire primary healthcare budget.

'HIGH RISK': HOW LOW CAN WE GO?

Current debate on 'High Risk as a Disease' twirls around the 'appropriate' threshold for defining something as a disease. The recently introduced osteoporosis guideline put out by medical experts of the National Osteoporosis Foundation (NOF, USA) recommends osteoporosis medications if a person's 10-year probability of sustaining a hip fracture is 3% or over. Applying these NOF recommendations to a large prospective cohort study, at least 72% of Caucasian women >65 years of age and 93% of those >75 years of age, in the USA, would be recommended for drug

therapy.⁴ The new cholesterol guideline similarly colonises virtually the entire elderly population into the realm of 'sick'.

UNDERSTANDING RISK: ARE THE BLIND LEADING THE BLIND?

But who are the right people to determine the threshold for 'high risk'? Advocates of the hegemony of medical experts argue that doctors—as content experts—should define diseases.⁵ If we assume doctors are truly more competent in making value judgements about the lives of their patients than the patients sitting in front of them, should we not have proof that doctors can do the job? Sadly, despite medical education and clinical experience, doctors do not seem to possess the required skill.⁶

Even more discouragingly, patients might not fully agree with our perceptions. For example, the above noted NOF threshold is more than 15 times *lower* than what patients would consider a 10-year fracture risk justifying initiation of bone-targeted pharmacotherapy (50%).⁷

Similarly, there is a huge gap on what the two stakeholders consider 'effective treatment'. Patients generally expect >20% *absolute* risk reduction (in heart attacks) for preventive pharmacotherapy to opt for treatment.⁸ In contrast, doctors world-wide began prescribing enthusiastically when an

osteoporosis drug was shown to increase the probability of avoiding a hip fracture from 97.9% to 98.9%. Indeed, this 1% *absolute* risk advantage convinced our peers; mind you, after being framed as a 50% *relative* risk reduction.

DOES THE HIPPOCRATIC OATH OBLIGE US TO INTERVENE?

One may wonder why we still intervene when not only our patients disagree with our views on what constitutes a plausible risk to be treated or an effective treatment but also when the preventive efforts are 10 times more resource consuming than the treatment for the event to be prevented? Most doctors argue that, in matters of life and death, we have no other option. However, in other disciplines affecting the health and well-being of humans, we easily make decisions based on cost. For example, teachers in primary school are well aware of the fact that there are numerous children in each class with learning disabilities who might come from troubled families and who would be naturally at 'high risk' of becoming illiterate. But does the increased risk of consequent social deprivation drive us to fund and execute large-scale special needs teaching for all 'at-risk kids'?

TREATING 'HIGH RISK'—A MOCKERY OF SHARED DECISION-MAKING?

Let us get back to the pivotal question: Is treating 'High Risk' a viable concept? A strategy of obtaining a single estimate of a patient's risk and reducing this risk effectively is appealing. However, evidence from behavioural sciences suggests that we are



Figure 1 Risk is the probability that something bad or unpleasant will happen. Intuitively, it seems like a very simple concept. However, there is strong evidence that both doctors' and patients' comprehension of RISK is poor.

Correspondence to Professor Teppo LN Järvinen, Clinical Medicine/Department of Orthopaedics and Traumatology, University of Helsinki and Helsinki University Central Hospital, Töölö Hospital/Topeliuksenkatu 5, Helsinki 00014, Finland; teppo.jarvinen@helsinki.fi

generally poor in making probability decisions.⁹ And, despite laudable efforts to improve the communication and comprehension of both the concept of risk and the anticipated treatment benefit, risk-illiteracy of the gravest magnitude still affects both doctors and patients (figure 1).⁶ But without accurate and *common* comprehension of these key aspects, there is no basis for shared decisions.¹⁰ And without shared decision-making, pharmacological primary prevention becomes a tyranny of eminence.

Acknowledgements Alan Cassels helped by adding comments and additional arguments and with the editing.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.



OPEN ACCESS

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

To cite Järvinen TLN. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2015-095286

Accepted 1 September 2015

Br J Sports Med 2015;0:1–2.
doi:10.1136/bjsports-2015-095286

REFERENCES

- 1 Moynihan R. Surrogates under scrutiny: fallible correlations, fatal consequences. *BMJ* 2011;343:d5160.
- 2 Tikkinen KA, Leinonen JS, Guyatt GH, *et al.* What is a disease? Perspectives of the public, health professionals and legislators. *BMJ Open* 2012;2:pii: e001632.
- 3 Getz L, Sigurdsson JA, Hetlevik I, *et al.* Estimating the high risk group for cardiovascular disease in the Norwegian HUNT 2 population according to the

2003 European guidelines: modelling study. *BMJ* 2005;331:551.

- 4 Donaldson MG, Cawthon PM, Lui LY, *et al.* Estimates of the proportion of older white women who would be recommended for pharmacologic treatment by the new U.S. National Osteoporosis Foundation Guidelines. *J Bone Miner Res* 2009;24:675–80.
- 5 Moynihan R. Medicalization. A new deal on disease definition. *BMJ* 2011;342:d2548.
- 6 Martyn C. Risky business: doctors' understanding of statistics. *BMJ* 2014;349:g5619.
- 7 Douglas F, Petrie KJ, Cundy T, *et al.* Differing perceptions of intervention thresholds for fracture risk: a survey of patients and doctors. *Osteoporosis Int* 2012;23:2135–40.
- 8 Trewby PN, Reddy AV, Trewby CS, *et al.* Are preventive drugs preventive enough? A study of patients' expectation of benefit from preventive drugs. *Clin Med* 2002;2:527–33.
- 9 Kahneman D, Tversky A. Prospect theory—analysis of decision under risk. *Econometrica* 1979;47:263–91.
- 10 Hoffmann TC, Montori VM, Del Mar C. The connection between evidence-based medicine and shared decision making. *JAMA* 2014;312:1295–6.



Labelling people as 'High Risk': A tyranny of eminence?

Teppo L N Järvinen

Br J Sports Med published online November 16, 2015

Updated information and services can be found at:

<http://bjsm.bmj.com/content/early/2015/10/01/bjsports-2015-095286>

These include:

Supplementary Material

Supplementary material can be found at:

<http://bjsm.bmj.com/content/suppl/2015/11/16/bjsports-2015-095286.DC1>

References

This article cites 10 articles, 5 of which you can access for free at:

<http://bjsm.bmj.com/content/early/2015/10/01/bjsports-2015-095286#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:

<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (287)

[Press releases](#) (50)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>