Ankle ligament injuries

Per A F H Renström, Lars Konradson

Injuries to the ligaments of the ankle are often called "low ankle sprains". If the tibiofibular ligament or the syndesmosis is injured it is called a "high ankle sprain".

Inversion sprains with injury to the lateral ligaments of the ankle/foot complex are by far the most common. They occur with an estimated frequency of one injury per 10 000 people per day, amounting to about 27 000 injuries each day in the United States.^{1,2} Although many injuries are treated outside medical establishments, 7-10% of those who are visiting the emergency departments of the hospitals in Scandinavia have sprained ankles.³

Ankle injuries are the most common injuries in sports and recreational activity.⁴⁻¹² For this reason, probably, these injuries tend to occur primarily to young people.¹³ The sprained ankle also remains the most common injury regardless of whether the sport is primarily an upper or a lower extremity sport. Garrick⁷ noted that injuries to the ankle accounted for 53% of injuries occurring during basketball and for 31% of those occurring during soccer. Reviewing 41 soccer teams, Ekstrand and Tropp¹⁴ found ankle sprains to account for 17 to 21% of the injuries. The "high ankle sprain" usually occurs as the result of an eversion injury in combination with fractures or lesions to the deltoid ligament. Isolated syndesmosis injuries occur in 3% of the cases. This article will deal mainly with the lateral ligament complex.

Ankle biomechanics

The passive stability of the ankle is the responsibility of the ligaments and the bony constraints of the ankle joint, while the active stability depends on muscular support. The talus has no muscular insertion. Active motion depends on the long foot muscles inserting into other tarsal or metatarsal bones. Dorsiflexion and inversion are effected by the extensor hallucis longus and the anterior tibial muscles. Dorsiflexion and eversion are guided by the peroneus tertius muscles and extensor digitorum longus and brevis muscles. Plantar flexion and eversion are effected by the peroneus longus and brevis muscles. Plantar flexion and inversion are regulated by the flexor hallucis longus, the flexor digitorum, and the posterior tibial muscles.¹⁵

The ligaments of the ankle can be divided into the lateral group, the medial group, and the ligaments of the syndesmosis.

The lateral ankle ligament complex is traditionally considered to consist of the anterior talofibular (ATFL), the calcaneofibular (CFL), and the posterior talofibular (PTFL) ligaments. However, in inversion the subtalar ligaments, especially the cervical

ligament, the interosseous ligament, and the ligaments spanning the calcaneocuboid and the talonavicular joints, also have to be considered.

Many studies have been done on talotibial ligaments to gain insight into how they function together to stabilise the joint. Of the talotibial ligaments, the ATFL is a thin 6-10 mm wide, 20 mm long, and 2 mm thick¹⁶ weak ligament, being essentially a thickening of the anterior ankle joint capsule. It passes from the distal anterior origin of the lateral malleolus to the talus in front of the proximal part of the lateral articular surface. In neutral position its direction is parallel to a long axis of the foot and in full plantar flexion it is more parallel with the tibia (fig 1). The CFL is a 20-25 mm long rounded ligament with a diameter of 6-8 mm.¹⁶ It is an extra-articular ligament closely associated with the peroneal tendon sheath. It runs obliquely downwards and backwards to be attached to the lateral surface of the calcaneus. There is a great variety in its direction and in its attachment sites.¹⁵ A rupture to this ligament will also cause a rupture of the tendon sheath, and occasionally also damage the peroneal tendons.

The ATFL acts as a primary restraint against plantar flexion, as well as internal rotation of the foot.¹⁷ In strain studies Renström *et al*¹⁸ found that the strain of the ATFL significantly increases with increasing plantar flexion. In the neutral position the ligament is relaxed.¹⁹

The CFL does not have an independent role in talotibial joint stability, but acts instead as a guide for the axis of subtalar motion.¹⁷ In dorsiflexion the ligament has increased strain.¹⁸ In a normal standing position the ligaments remain relaxed.

The lateral talocalcaneal (LTCL) extends from the talus to the calcaneus and blends its fibres with CFL and ATFL fibres. The exact incidence of injury to this ligament is not known. Transecting the subtalar ligaments results in very limited increase in motion when measured in degrees, but as they have very limited motion in the first place the increase after rupture is about 40%.²⁰ The incidence of rupture here is also unknown.

The PTFL connects the posterolateral tubercle of the talus to the medial aspect of the lateral malleolus. The PTFL has an average diameter of 6 mm. In plantar flexion and in the neutral position the ligament is relaxed, whereas in dorsiflexion the ligament is tensed.¹⁵ The clinical significance of PTFL injuries is somewhat unclear, but it is not commonly damaged. The ATFL, CFL, and PTFL ligaments function as a unit for the talotibial complex, though one may resist a specific motion depending upon foot position.²¹

Department of
Orthopaedics and
Rehabilitation,
McClure
Musculoskeletal
Research Center, The
University of Vermont,
Stafford Hall 428B,
Burlington, Vermont
05405-0084 USA
P A F H Renström

Department of
Orthopaedics,
Gentofte Hospital,
University of
Copenhagen,
Copenhagen, Denmark
L Konradson

Correspondence to:
Dr Renström.

Accepted for publication
17 December 1996

Figure 1 (A) Anterior talofibular (atf) ligament runs parallel to the axis of the foot when the foot is in neutral position. Cf = calcaneofibular ligament. (B) When the foot is in plantar flexion, the anterior talofibular ligament assumes a course parallel to the axis of the tibia and fibula.

Through the full range of motion the ATFL and CFL act in synergy²²⁻²⁵ (fig 1). As the foot plantar flexes, the strain in the ATFL increases while the strain in the CFL decreases.¹⁸ Shybut *et al*²⁵ measured ankle ligament loads directly by using implanted buckle transducers. The results indicated that ligament loads remain low within the functional range of motion (10 degrees of dorsiflexion to 20 degrees of plantar flexion). This supports the concept that ankle ligaments act as kinematic guides rather than primary restraints during normal activity.

Stormont *et al*²⁶ studied the stabilising capacity of the ligaments and articular surface in the ankle with and without physiological loading. With loading, the results indicated that the articular surface becomes an important stabiliser, accounting for 30% of stability in rotation and 100% of stability in inversion. Without loading, the results indicated that the primary and secondary ligamentous constraints vary with testing modes and ankle position.

Mechanism of injury

The extent of tissue damage that will occur with the trauma depends not only on the mechanism and magnitude of the forces that act on the ankle but also on the position of the foot and ankle during the trauma.¹⁵

The most common mechanism causing lateral ligament injuries is a situation where the ankle goes into a combination of plantar flexion and inversion. The ATFL tears first followed by rupture of the anterolateral capsule. With further inversion, the CFL will be ruptured followed by variable injury to the PTFL and the anterior part of the deltoid ligaments.²⁷

With weight bearing, the articular surface can provide 30% of stability in rotation, and 100% stability in inversion.²⁶ This ability is a function not only of the axial load but also of the close packed position.²⁸ Ankle destabilisation thus occurs during loading and unloading, but the joint is stable once it is fully loaded.

Frequency of lesions caused by inversion sprains

Owing to the ATFL's vulnerable position in plantar flexion, it is the most commonly ruptured ligament in a lateral ankle sprain.^{13 23 29-34} In 1964 Broström¹⁶ surgically explored 105 sprained ankles and found an isolated ATFL tear present in about two thirds of the cases. The second most common injury was a combined rupture of the ATFL and CFL, occurring in about 20 to 25% of the cases.^{13 16 34} Other isolated ligamentous injuries are relatively uncommon.^{13 16} The PTFL, for instance, is a very strong ligament and is rarely injured except in severe ankle trauma.^{29 35}

Ligamentous lesions after acute inversion sprains cannot, however, be seen in such a limited scope. Thus Broström¹³ noted that in a group of 321 patients with acute ankle sprains, 19% had signs of injury to the bifurcate or the dorsal calcaneocuboid ligaments, or both. Gerner-Smidt³⁶ in a combined study of children and adults found that 22% of the patients sustained lesions to the talonavicular or the calcaneocuboid ligaments, or both, and Hølmer *et al*³⁷ found clinical evidence of isolated calcaneocuboid/calcaneocervical or talonavicular ligament lesions in 15-25% of inversion injuries. Meyer *et al*³⁸ evaluated 40 patients with acute ankle sprains (it is not mentioned, but they must have been considered clinically grade II to III) with subtalar arthrograms.³⁹ Apart from lesions to the lateral tibiotalar joint ligaments, 17 patients (43%) showed contrast leaks into the sinus tarsi, suggesting interosseous ligament rupture. Some of these patients were operated on and the cervical ligament was often found to be severed too.

Chondral or osteochondral lesions of the talar dome have been noted by Taga *et al*⁴⁰ in 89% and by van Dijk¹⁵ in 66% of acute inversion injuries. In 1991 Grana⁴¹ found chondral lesions in 80% and osteochondral lesions in 6.5% of acute ankle injuries. Lesions to peroneal tendons ranging from total tendon rupture or insertion site fracture to longitudinal slits occur, and the tendon injuries are often overlooked.⁴² Injury to the superficial and deep peroneal nerves ranging from the rare condition of complete nerve palsy^{43 44} to, what seems to be much more common, discrete conduction velocity changes have been noted.^{45 46} In rarer cases fractures to the cuboid, the anterior process of the calcaneus, and the lateral process of the talus have been diagnosed after an inversion injury.⁴⁷⁻⁴⁹

Grading lateral ligament injury

Clinically, sprains of the lateral ankle have been classified in three groups based on severity.⁴ A grade I injury involves stretch of the ligament without macroscopic tearing, little swelling or tenderness, slight or no functional loss, and no mechanical instability of the joint. A grade II injury is a partial macroscopic tear of the ligament with moderate pain, swelling, and tenderness over the affected structures. Some loss of motion and mild or moderate instability

of the joint occurs. In a grade III injury, there is complete rupture of the ligament with severe swelling, haemorrhage, and tenderness. There is loss of ability to bear weight on the foot, limited function, and considerable abnormal motion and instability of the joint. An improved and validated classification of ankle sprains is needed. The terms mentioned above were validated by Lindenfeld⁵⁰ and found to be quite subjective. Others prefer to classify lateral ligament injuries as single or double ligament tears. A single injury would imply lesions to only the ATFL, a double injury would affect both the ATFL and the CFL. The distinction in clinical terms is difficult here also.

Diagnosis

Diagnosing the extent of an acute lateral ligament injury is not considered very accurate because of pain, swelling, and muscle tenderness.¹⁵ In the first few days after injury local palpational pain is often diffuse, with no maximal point of tenderness.^{151 52} The extent of swelling does not depend only on the magnitude of the ligament injury but also on the initial treatment. Extensive swelling is a predictor of ligament rupture, but the positive predictive value has been found to be only 60–70%.^{53–55} The characteristic haematoma suggesting ligament rupture usually does not develop during the first few days, and joint range of motion is mainly determined by the severity of pain and does not differ between simple sprains and ligament rupture.^{54 55} Furthermore, the interobserver variation of the acute examination concerning tenderness, swelling, discoloration, and the anterior drawer sign is considerable.⁵⁶ The anterior drawer sign seems to be a good predictor of lateral ligament disruption. Broström³⁰ found that the anterior drawer sign was positive in all of 239 patients with ligament rupture. His examination was, however, done with general or local anaesthesia. Without anaesthesia he could only elicit a positive drawer sign in two patients. Based on these results van Dijk³⁷ suggests that **the physical examination should be delayed four to five days after the initial injury.** The specificity and sensitivity of delayed physical examination for the presence or absence of a rupture to the ATFL were found to be 84% and 96% respectively, and the delayed examination, done by observers of varying degrees of experience, gave information of ligament quality that equalled that of arthrography. After four to five days a combination of tenderness at the level of the anterior talofibular ligament, lateral haematoma discoloration, and a positive drawer sign indicated a ligament lesion in 95% of the cases. A negative drawer test and the absence of discoloration always indicated an intact ligament.

Radiographs to exclude fractures are suggested according to the Ottawa strategy for ankle injury.⁵⁸ Plain radiographs are taken if there is bone tenderness at the tip or posterior aspect of the lateral malleolus, at the tip or posterior aspect of the medial malleolus, at the

navicular tuberosity or at the base of the fifth metatarsal, or if the patient is unable to bear weight immediately after injury and at the initial examination. With these rules the authors could reduce the use of radiographs from 80% to 63% of the injuries.

Treatment of lateral ligament lesions

TREATMENT OF GRADE I AND II SPRAINS

In the presence of a grade I or grade II injury it is universally agreed that recovery is fast with non-operative management and the prognosis is good.^{4 55 59} Jackson *et al*⁵⁹ found that early mobilisation resulted in a disability of eight days for a grade I and 15 days for a grade II injury. **Functional treatment including early motion and use of ankle support and early weight bearing is today the accepted treatment for grade I and II ankle sprains.**

TREATMENT OF GRADE III SPRAINS

In the case of grade III lateral ligament lesions the treatment of choice—that is, whether to operate, to immobilise in a cast, or to allow early controlled mobilisation, is more controversial. The key consideration being the question of whether ligament healing with adequate tension can be achieved equally well with early controlled mobilisation as with direct visualisation and suturing.

If the ankle is kept from excess inversion or dorsal or plantar flexion the strain in the lateral ligaments remains low,¹⁸ allowing for adequate healing conditions without elongation within this range of guided motion. With the development of lower extremity magnetic resonance imaging scanning, studies have visualised that total ligament (grade III) ruptures do heal with ligament continuity if there is early mobilisation. There is thus experimental evidence supporting ankle ligament healing in the presence of early mobilising treatment.

Clinically many uncontrolled, non-randomised studies have shown mechanical stability and satisfactory subjective results after both operative and conservative treatment. Of the many papers available, only 12 could be considered prospective and randomised^{30 55 60–69} when a comparison between operative and conservative treatments was performed by Kannus and Renström.³⁵ Six of these considered the three treatment groups in question: operation, immobilising and early mobilising modalities,^{30 62 63 65 66 69} and the results and conclusions based on these reports will be mentioned in the following. Duration of follow up ranged from six months to 3.8 years, which is considered adequate time to identify persistent disability. Results were evaluated using selected outcome parameters. Return to work or physical activity was reported in four of the studies that included three treatment modalities.^{30 61 66 69} They concluded that return to work was two to four times faster after functional treatment than after operation or immobilisation in a cast. Return to pre-injury level of activity was found to be faster after conservative treatment than after operative treatment in four cases.^{61 62 66 69} The opposite was found in

three studies^{30 55 63} and no difference was found in the remaining five.

Pain, swelling, or stiffness with activity could be evaluated in four of the studies involving three treatment groups.^{62 63 66 69} The studies failed to show any differences. In the three studies that included three therapy groups and included mobility of the ankle,^{30 66 69} mobility was found to be superior after functional treatment compared with the other two methods.

Better mechanical stability has been the primary argument for operation. Assessment of mechanical stability calls for an objective measurement with the evaluation of talar tilt and anterior drawer on stress radiographs. Based on five studies evaluating the three different treatments,^{62 65 66 68 69} equal stability results were found.

Functional instability is not a well defined entity but designates repeated inversion injuries that are either unprovoked or the result of very little provocation. Functional instability seems to be a late disability in 15 to 60% of lateral ligament injuries depending on the definition applied. In the studies that reported on functional instability^{30 62 66 69 71} opinions diverged, but no one treatment seemed superior to the others in minimising the chances of late instability. The scientific basis for treating chronic functional instability is discussed later in this review.

The review of Kannus and Renström³⁵ concludes that **functional treatment is the treatment of choice when treating grade III acute lesions of the lateral ligaments**. Recent comparative studies do not change this view. Konradsen *et al*² and Eiff *et al*³ concluded that in first time lateral ankle sprains, although both immobilisation and early mobilisation prevent late residual problems and ankle instability, early mobilisation allows early return to work and, possibly, is more comfortable for patients. In a recent prospective study comparing surgery with functional treatment in ankle ligament tears Kaikkonen and coworkers⁷⁴ found that early mobilisation gave better results than surgery plus mobilisation in treatment of complete tears of lateral ligaments of the ankle. Nine months after injury excellent to good scores were achieved in 87% of the functionally treated patients and 60% of the surgically treated patients respectively. In a meta-analytical article Schrier⁷⁵ found functional treatment was superior to immobilisation in casts, reducing both ankle instability and the cost of treatment.

Sommer and Schreiber⁷⁶ considered the cost-benefit aspects by comparing immobilisation in a plaster cast followed by immobilisation with a brace, with early mobilisation using a brace only. Early functional therapy proved to be the treatment with the least direct costs. Leanderson and Wredmark⁷⁷ in a study of 73 patients with grade II and III sprains treated with either early mobilisation or immobilisation found that the socioeconomic savings were potentially significant with ankle brace treatment. Nationally, the potential yearly savings in Sweden using this treatment were estimated to

be eight million American dollars. The treatment of choice thus remains functional treatment.

If acute surgery is considered necessary the indications could according to Leach and Schepesis⁷⁸ be (a) a history of momentary talocrural dislocation with complete ligamentous disruption, (b) a major clinical anterior drawer sign, (c) 10 degrees more tilt on the affected side with stress inversion testing, (d) clinical or radiographic suspicion of tears in both the ATFL and CFL, and (e) osteochondral fracture. Most techniques described for repair of acute ligament injuries are similar to that of Broström.³⁰ The results after acute surgery are in general very good with return to sport in 10 to 12 weeks. This must still be compared with the three to six weeks after functional treatment.

Acute treatment protocol

BIOLOGICAL BACKGROUND

In the post-injury phases of an acute severe ankle injury, an ideal treatment and rehabilitation programme should fulfil four requirements³⁵:

- 1 The RICE principle: rest, ice (cold), compression, and elevation aims to minimise haemorrhage, swelling, inflammation, cellular metabolism, and pain in order to offer the best possible conditions for the healing process.^{79 80}
- 2 Protection of the injured ligaments during the first one to three weeks is required. In this phase of healing (the proliferation phase), protective ankle support is followed by undisturbed fibroblast invasion of the injured area, which leads to undisturbed proliferation and production of collagen fibres. Mobilisation too early in inversion leads to more prolonged type III collagen formation with weaker healing tissue than during optimal immobilisation.⁸⁰⁻⁸² Protection is also needed to prevent secondary injuries and early distension and lengthening of the injured ligaments.
- 3 About three weeks after the injury the maturation phase of the collagen and the formation of final scar tissue begins.⁸¹⁻⁸³ In this phase the injured ligaments need controlled mobilisation, and perhaps even more importantly, the ankle itself must avoid the deleterious effects of immobilisation on joint cartilage, bone, muscles, tendons, and ligaments.^{8 83-90} Controlled stretching of muscles and movement of the joint enhance the orientation of collagen fibres parallel with the stress lines (that is, the normal collagen fibres of the ligaments), and they can prevent the atrophy caused by immobilisation.^{80 82} Repeated exercises will also increase the mechanical and structural properties of the ligaments.⁹¹
- 4 About four to eight weeks after the injury the new collagen fibres begin to withstand almost normal stress, and the goal for rehabilitation is rapid recovery and full return to work and sports.

If treated according to the guidelines mentioned above, each component of the ankle is

ready for a gradually increasing mobilisation and rehabilitation programme, keeping in mind that final maturation and remodelling of the injured ligaments takes a long time—from six to 12 months.

Residual disability after inversion injury

Inadequate rehabilitation is the primary cause of residual disability after ankle sprains. Many athletes return to sports before they are fully rehabilitated and therefore they are often subjected to reinjury or additional injury. Examination may show loss of range of motion, especially limited dorsiflexion, and atrophy of lower leg muscles.

Residual disability after ankle inversion injury can be divided into primarily instability problems and pain-giving entities.

CHRONIC ANKLE INSTABILITY

Chronic ankle instability can be subdivided into mechanical, functional, and subtalar instability and the sinus tarsi syndrome.

Mechanical instability

Mechanical instability is characterised by ankle mobility beyond the physiological range of motion, which is identified on the basis of a positive anterior drawer or talar tilt test, or both.⁹² The radiographic criteria for mechanical instability vary. Most authors agree, however, that mechanical instability is present when there is more than 10 mm of anterior translation on one side or the side-to-side difference is over 3 mm or the talar tilt is more than 9 degrees on one side, or the side-to-side difference is more than 3 degrees.⁹³ Pure mechanical instability of the ankle is seldom the sole independent reason for the development of late symptoms. Mann *et al*, showed that 81% of patients with radiographically documented instability experienced recurrent sprains.⁹⁴

Functional instability

Functional instability is the chronic disability after an ankle inversion injury that can be attributed to lateral ankle ligament deficiency and is characterised by “giving way” problems. Forty eight per cent of patients with an acute first time sprain have recurrent sprains and 26% report frequent sprains.⁹⁴ The definition of the symptom is ambiguous and the pathogenesis is unclear. It does not seem to be dependent on the grade of the initial injury⁶⁹ nor does it correlate with the degree of initial mechanical instability.^{61 64 69-95} No histological ligamentous changes except scarring have been found in chronic ankle disability.³⁰

As anatomical studies showed the presence of mechanoreceptors in joint ligaments and capsule, and clinical studies showed decreased postural control in patients with functional instability, Freeman *et al* suggested that functional instability was due to motor incoordination secondary to a proprioceptive disorder.⁹⁶ Postural control was later measured objectively using stabilometry by Tropp⁹⁵ who, like Freeman, found increased postural sway in subjects with functionally unstable ankles. It is, how-

ever, clear that functional instability is a complex syndrome where mechanical, proprioceptive, and muscular disabilities either alone or in combination are at fault. Known factors include mechanically insufficient ligaments, peroneal muscle weakness,⁹⁵ subtalar instability,⁹⁷ and proprioceptive deficit.^{95 96}

Although increased postural swaying has been used as a measure for a proprioceptive deficit, there is no direct causal connection between swaying with an increased amplitude when balancing on one foot and sustaining recurrent inversion injuries. Rather it seems reasonable to assume that a common and superior deficit is responsible for both disabilities. The nature of such a disability remains unclear. The latency of the peroneal muscles has been found to be significantly slower in functionally unstable ankles by some⁹⁸⁻¹⁰⁰ but not by others.¹⁰¹⁻¹⁰³ An increased error in detecting ankle inversion movement or in matching ankle inversion angles has again been found by some¹⁰⁴⁻¹⁰⁶ but not by others.¹⁰⁷ There is, however, evidence that (a) a proprioceptive deficit may result after an acute ankle inversion injury,^{46 70 108} (b) proprioceptive defects measured in different ways are seen in subjects with functional instability, and (c) models exist that may explain how measurable kinaesthetic deficits may cause an increased frequency of ankle inversion injuries (Konradson *et al*, unpublished data).

Subtalar instability

The incidence of subtalar sprains is unknown, but it is widely accepted that most subtalar ligamentous injuries occur in combination with injuries of the lateral ligaments of the ankle (fig 2). Subtalar instability is estimated to be present in about 10% of patients with lateral ankle instability. The symptoms of chronic subtalar instability include giving way episodes during activity and a feeling of instability when walking on uneven ground. These symptoms may coincide with chronic talocrural instability and therefore careful clinical examination is necessary. Localised tenderness on palpation over the subtalar joint is suggestive of a subtalar sprain. The diagnosis can be verified with subtalar arthrography³⁸ or a subtalar stress view or stress tomography.

Functional treatment similar to the regimen used for lateral sprains is the treatment of choice. Surgery with non-anatomical procedures has been described.^{109 110} Anatomical repairs, however, show equally good results.^{97 111}

Sinus tarsi syndrome

The sinus tarsi syndrome is often accompanied by a feeling of instability and giving way of the ankle. Seventy per cent of the patients will have experienced a severe inversion sprain.⁹² If the calcaneofibular ligament is torn the interosseous talocalcaneal ligament occupying the sinus will often also be sprained. This sprain will cause synovitis in the sinus tarsi with rather localised pain. The diagnosis can be made based on a complaint of pain and tenderness of the sinus tarsi combined with the giving way

Figure 2 Forceful inversion of the hindfoot in neutral flexion. Tearing of the calcaneofibular, the cervical, and the interosseous ligaments. (From Meyer JM, Garcia J, Hoffmeyer B, Fritchy D. The subtalar sprain. A roentgenographic study. *Clin Orthop* 1988;226:169–73.)

feeling. The pain and the giving way feeling can usually be relieved by local injections of anaesthetic and corticosteroids. Excision of tissue filling the sinus tarsi can give good results if conservative treatment fails.

TREATMENT OF CHRONIC INSTABILITY

Rehabilitation

With the uncertain pathogenesis of functional ankle instability the elements of treatment protocols will rest primarily on personal experiences and not so much on scientific data. Before deciding upon surgical treatment, a well planned rehabilitation programme based on peroneal muscle strengthening and coordination training should be carried out. Regaining muscle strength is primarily achieved by increasing neural activation through functionally oriented training and dynamometer training.¹¹² In functional training the patient is required to activate his/her musculature in normally occurring activities. Dynamic stability, including proprioceptive information, motor control, and appropriate muscle force development, will be enforced by the training.

Treatment of the proprioceptive ankle deficit consists of a progressive series of coordination exercises to re-educate the ankle. Tropp⁹⁵ found a significant improvement in balance during single limb stance in subjects with a functionally unstable ankle after six weeks of balance board exercises (fig 3). The same group reported a decrease in the frequency of inversion injuries and ankle giving way feelings when compared with an untrained control group with unstable ankles. Tropp⁹⁵ also found that the subjects in question showed a change in their strategy of single limb stance, going from a broken chain strategy before training to the normal inverted pendulum strategy after completed training. So it may well be, as suggested by Glencross and Thornton,¹⁰⁴ that

Figure 3 Tilt board training is an excellent way to return neuromuscular control to the injured ankle.

rehabilitation was as much a relearning programme as it was a physical recovery after injury. Often the injured subjects themselves have a clear feeling of a perception/proprioception deficit in the ankle joint area—they cannot “feel” the ankle as they used to. When this feeling returns it seems to coincide with complete rehabilitation. Leanderson *et al*¹⁰⁸ found that return of single limb balance to normal values coincided with a subjective feeling of full rehabilitation in dancers. It can, however, be argued that in this group balance during single limb stance was as much a specific functional test as an unspecific test for ankle proprioception.

In summary, training proprioceptive function through coordination exercises using unilateral balance boards, uniaxial and multiaxial teeter boards, and jumping ropes relies primarily on empirical data but produces very favourable results. Restitution of normal function relies on the individual's subjective feeling of return to normal sensation. Full effect of training is achieved within 10–12 weeks as evaluated by postural sway.⁹⁵

About 50% of patients with functional instability will regain satisfactory functional stability after such a programme.¹¹³

EXTERNAL SUPPORT

When considering the mechanical effect of the different external support modalities it is important to remember that even though they do increase the resistance to inversion, the effect cannot be compared with that of active evertor muscles. Ashton-Miller *et al*¹¹⁴ found that active evertor muscles, acting isometrically on a 15 degrees inverted ankle, can provide more than three times the protection against

further ankle inversion than tape or an orthosis worn inside a three-quarter-top shoe.

Tape

The primary purpose of taping is to provide increased stiffness to the ankle and a semirigid and sometimes rigid splinting around it. Studies have shown that taping the ankle prevents ankle injuries. Garrick *et al* found that the combination of taping prophylactically and the use of high top shoes in basketball reduced the frequency of ankle sprains.^{114a} Lindenberger *et al*¹¹⁵ found that tape had a significant prophylactic effect in handball team players who were studied prospectively for two full seasons. Six top level teams participated in the first year. In three teams the players wore tape and in the other three teams the players did not. During the first season 13 ankle sprains occurred—all in the non-taped group. In the second season nine teams participated. Twenty one ankle sprains occurred and 20 of these were in the non-taped group.

Contrary to this, several studies have shown that tape loses as much as 50% of its supportive effect after only 10 minutes of active exercise^{116–118} and offers virtually no support after one hour.¹¹⁹ The mechanism behind the effect of ankle taping thus remains unclear. However, tape does not only restrict extensive joint motion but also enhances proprioceptive feedback mechanism and shortens the recruitment time of the dynamic ankle stabilisers.¹²⁰

Many athletes may have a skin reaction to tape and therefore skin protection may be used. There seems to be no difference in support between tape directly on the skin and tape with skin protections. Because of these skin problems, tape is used mostly by top athletes and not by recreational athletes.

Braces

The use of ankle orthoses has increased over the past decade. In contrast with tape the mechanical effect of a brace does not wear off with activity as shown by Shapiro *et al*.¹²¹

Not many clinical trials have been performed judging the effect of ankle bracing on the frequency of ankle inversion injuries. In a retrospective trial Rovere *et al*¹²² found a reduction of sprains in football using a lace-on cloth brace. Tropp *et al*¹²³ in a prospective study showed the effect of a semirigid ankle brace in reducing the frequency of injuries in soccer players, and in a prospective randomised study Sitler *et al*¹²³ found that ankle stabilisers significantly reduced the frequency of ankle injuries but not their severity in a group of basketball players at West Point, New York. The resistance to an eversion moment seems to be the same for ankle braces as that for newly applied tape.¹¹⁴ Like tape, braces have been shown to have a proprioceptive enhancing effect.¹²⁴

There have been discussions about the effect on performance of tape and braces. In sprinting a reduction in maximal speed has been found by most,¹²⁵ but not by all.¹²⁶ For vertical jumping most studies also show that a brace decreases ability.^{126–128} Opinions about bracing compared with taping vary, with some

studies showing a smaller effect and others a larger detrimental effect of tape compared with braces. It must be noted that most of the studies were done on subjects with stable ankles. **The use of an ankle brace or taping, or both, is recommended as they both give support against ankle inversion when the evertor muscles are relaxed and increase the maximal resistance to inversion with evertor muscles activated.** They do seem to be able to enhance proprioceptive awareness, and to decrease the frequency of inversion injuries and decrease the severity of the injuries in groups with functionally unstable ankles.

The role of shoes

Biomechanical studies support the use of high top shoes for ankle sprain prevention because they limit extreme range of motion, provide additional proprioceptive input, and decrease external stress on joints.¹²⁹ Clinical trials, however, have not shown that the effect of this biomechanically improved stability translates into a lower rate of ankle sprains in high top shoes. Further studies are needed.

Supportive treatment modalities

Passive physical therapy modalities are often recommended to promote healing in the early rehabilitation phase. The most commonly used are ultrasound, temperature contrast baths, short waves, and various current treatments including dynamic or interference current therapy, or electrogalvanic stimulations. Of these different types of passive physical therapy, only cryotherapy has been proved to be effective.¹³⁰

Non-steroidal anti-inflammatory drugs have been studied in prospective randomised double blind trials, and found to be more effective than placebo for ankle tenderness, swelling, and pain, though the difference was not striking and seemed to disappear during an extended follow up.^{3 130}

In subjects with chronic post-injury swelling, moist heat packs, warm whirlpool baths, contrast baths, electrogalvanic stimulation, and intermittent pneumatic compression have been tried.⁴ Prospective randomised studies have shown that only the last alternative has an independent ability to reduce the amount of swelling.^{131 132}

Table 1 Classification of operative treatments for chronic ankle ligament injury

Non-anatomical reconstruction	
Non-anatomical reconstruction	
Endogenous	
Peroneal tendon	
Watson-Jones	
Evans	
Chrisman-Snook	
Other	
Plantaris	
Partial Achilles tendon	
Free autogenous graft	
Exogenous	
Carbon fibre	
Bovine xenograft	
Anatomical repair	
Direct suture	
Imbrication and repair to bone	
Local tissue augmentation

Figure 4 Anatomical reconstruction of chronic ankle ligament instability. (A) Elongated ligaments are divided 3 to 5 mm from insertion of the fibula. (B) Bone surface of the distal end of the fibula is roughened to form a trough to promote ligament healing. Holes are drilled through the distal fibula. (C) Mattress sutures are used to fix the distal stump of the ligaments and the capsule to the fibula. The sutures are tightened while the foot is held in dorsiflexion and eversion. (D) The proximal ends of the ligaments are imbricated over the distal portion.

Surgery for chronic instability

Chronic ankle instability, as demonstrated by pain, recurrent giving way, and positive stress tests, is often treated surgically. A combination of mechanical and functional instability is the most commonly reported indication for surgery. More than 50 procedures or modifications have been described for treating chronic ankle instability, and these can be loosely grouped as non-anatomical reconstructions or anatomical repairs (table 1).

Non-anatomical reconstructions use other structures or materials to substitute for the injured ligaments. Structures commonly used for grafting are fascia late and the peroneus brevis tendons in procedures such as Watson-Jones, Evans, and Chrisman-Snook. Numerous modifications of these classic procedures have been described. The Chrisman-Snook¹⁰⁹ modification of the Elmslie procedure is the most widely used non-anatomical reconstruction. In four clinical series with a total of 100 ankles, more than 90% of patients had good or excellent results, and stability was more than 95% of the normal ankle. Reported complications included neuromas in 0 to 16% of patients, restricted dorsiflexion in about 20%, and restricted inversion to some degree in all patients. An advantage of the Chrisman-Snook procedure is that less lateral weakness is produced because only half of the peroneus brevis tendon is used for the graft. A criticism of the procedure has been that it results in restricted subtalar motion. Furthermore, **these tenodesis procedures are non-anatomical and do not restore normal biomechanics.**¹³³ Although short term results are excellent, long term results may not be as

good. Thus Karlsson *et al*⁹³ found that the Evans static tenodesis was satisfactory in fewer than 50% of cases after a mean follow up period of 14 years.

Anatomical reconstructions, in which the tissues of the damaged ligaments are used, have become increasingly popular as they permit reconstruction without sacrificing any normal tissue.^{30 93 134} Broström in 1966³⁰ reported that direct suture repair of chronic ankle ligament injuries was possible even many years after the initial injuries and that the ends of the ligament could be found. Others reported that the elongated ligaments had healed encased in a fibrous scar tissue. Several authors have reported successful imbrication or shortening and replantation of the ligaments to obtain a more anatomical reconstruction. We have obtained good results with a modified Broström technique (Peterson procedure) that includes shortening of the ligament, repair through bony tunnels, and imbrication with local tissue^{93 134} (fig 4). This anatomical technique repairs both the ATFL and CFL. The non-anatomical reconstructions, except for the Chrisman-Snook modification of the Elmslie procedure, repair only the ATFL. Repair of the CFL is important, because insufficiency of this ligament may be a factor in the development of subtalar instability. The post-operative treatment after an anatomical reconstruction is a splint for eight days to secure wound healing and, thereafter, a removable walking boot for five weeks. The patient starts with dorsiflexion and plantar flexion exercises after eight days or as early as possible out of the boot two to three times a day. Return to sport is usually possible in three months.

Anatomical repair of both the ATFL and the CFL through bony tunnels produces good long term results and is recommended as the initial procedure in most cases. If anatomical repair fails, a tenodesis procedure, such as the Chrisman-Snook reconstruction, is a good alternative. A non-anatomical reconstruction is also indicated in patients with moderate arthritis or lax joints.

- 1 Makhani JS. Diagnosis and treatment of acute rupture of the various components of the lateral ligaments of the ankle. *Am J Orthop* 1962;4:224-30.
- 2 McCulloch PB, Holden P, Robson DJ, *et al*. The value of mobilisation and nonsteroidal anti-inflammatory analgesia in the management of inversion injuries of the ankle. *Br J Clin Pract* 1985;39:69-72.
- 3 Viljakka T, Rokkanen P. The treatment of ankle sprain by bandaging and antiphlogistic drugs. *Ann Chir Gynaecol* 1983;72:66-70.
- 4 Balduini FC, Vegso JJ, Torg JS, *et al*. Management and rehabilitation of ligamentous injuries to the ankle. *Sports Med* 1987;4:364-80.
- 5 Brand RL, Black HM, Cox JS. The natural history of inadequately treated ankle sprain. *Am J Sports Med* 1977;5:248-9.
- 6 Fiore RD, Leard JS. A functional approach in the rehabilitation of the ankle and rear foot. *Athletic Training* 1980;15:231-5.
- 7 Garrick JG. The frequency of injury, mechanism of injury and epidemiology of ankle sprains. *Am J Sports Med* 1977;5:241-2.
- 8 Garrick JG, Requa RK. The epidemiology of foot and ankle injuries in sports. *Clin Sports Med* 1988;7:29-36.
- 9 Glick JM, Gordon RB, Nashimoto D. The prevention and treatment of ankle injuries. *Am J Sports Med* 1976;4:136-41.
- 10 Hardaker WT, Margello S, Goldner JL. Foot and ankle injuries in theatrical dancers. *Foot Ankle* 1985;6:59-69.

- 11 Lassiter TE Jr, Malone TR, Garrett WE Jr. Injury to the lateral ligaments of the ankle (review). *Orthop Clin North Am* 1989;20:629-40.
- 12 McConkey JP. Ankle sprains, consequences and mimics. *Medicine and Sports Science* 1987;23:39-55.
- 13 Broström L. Sprained ankles III: Clinical observations in recent ligament ruptures. *Acta Chir Scand* 1965;130:560-9.
- 14 Ekstrand J, Tjørrø H. The incidence of ankle sprains in soccer. *Foot Ankle* 1990;11:41-4.
- 15 Van Dijk CN. *On diagnostic strategies in patients with severe ankle sprain*. Amsterdam, Holland: University of Amsterdam, 1994.
- 16 Broström L. Sprained ankles I: anatomic lesions in recent sprains. *Acta Chir Scand* 1964;128:483-95.
- 17 Rasmussen O. Stability of the ankle joint. Analysis of the function and traumatology of the ankle ligaments. *Acta Orthop Scand Suppl* 1985;211:1-75.
- 18 Renstrom P, Wertz M, Incavo S, et al. Strain in the lateral ligaments of the ankle. *Foot Ankle* 1988;9:59-63.
- 19 Colville MR, Marder RA, Boyle JJ, et al. Strain measurement in lateral ankle ligaments. *Am J Sports Med* 1990;18:196-200.
- 20 Kjaersgaard-Andersen P, Wethelund JO, Helvig P, et al. The stabilizing effect of the ligamentous structures in the sinus and canalis tarsi on movements in the hindfoot. An experimental study. *Am J Sports Med* 1988;16:512-6.
- 21 Cass JR, Morrey BF, Chao KY. Three-dimensional kinematics of ankle instability following serial sectioning of lateral collateral ligaments. *Foot Ankle* 1984;5:142-9.
- 22 Cawley PW, France EP. Biomechanics of the lateral ligaments of the ankle: an evaluation of the effects of axial load and single plane motions on ligament strain patterns. *Foot Ankle* 1991;12:92-9.
- 23 Dias LS. The lateral ankle sprain: an experimental study. *J Trauma* 1979;19:266-9.
- 24 Ozeki S, Yasuda K, Kaneda K. *Simultaneous measurement of strain changes and determination of zero strain in the collateral ligaments of the human ankle*. 36th Annual Meeting, Orthopaedic Research Society, 1990.
- 25 Shybut GT, Hayes WC, White AA. *Normal patterns of ligament loading among the lateral collateral ankle ligaments*. 29th Annual Meeting, Orthopaedic Research Society, 1983.
- 26 Stormont DM, Morrey BF, An KN, et al. Stability of the loaded ankle. Relation between articular restraint and primary and secondary static restraints. *Am J Sports Med* 1985;13:295-300.
- 27 Rasmussen O, Kromann-Andersen C, Boe S. Deltoid ligament. Functional analysis of the medial collateral ligamentous apparatus of the ankle joint. *Acta Orthop Scand* 1983;54:36-44.
- 28 Smith RW, Reischl SF. Treatment of ankle sprains in young athletes. *Am J Sports Med* 1986;14:465-71.
- 29 Brand RL, Collins MD. Operative management of ligamentous injuries to the ankle. *Clin Sports Med* 1982;1:117-30.
- 30 Broström L. Sprained ankles V - treatment and prognosis in recent ligament ruptures. *Acta Chir Scand* 1966;135:537-50.
- 31 Glasgow M, Jackson A, Jamieson AM. Instability of the ankle after injury to the lateral ligament. *J Bone Joint Surg [Br]* 1980;62:196-200.
- 32 Kjaersgaard-Andersen P, Frich LH, Madsen F, et al. Instability of the hindfoot after lesion of the lateral ankle ligaments: investigations of the anterior drawer and adduction maneuvers in autopsy specimens. *Clin Orthop* 1991;266:170-9.
- 33 Leonard MH. Injuries of the lateral ligaments of the ankle. A clinical and experimental study. *J Bone Joint Surg [Am]* 1949;31:373-7.
- 34 Lindstrand A. New aspects in the diagnosis of lateral ankle sprains. *Orthop Clin North Am* 1976;7:247-9.
- 35 Kannus P, Renström P. Current concepts review: treatment for acute tears of the lateral ligaments of the ankle. *J Bone Joint Surg [Am]* 1991;73:305-12.
- 36 Gerner-Smidt M. *Ankle fractures in children*. Copenhagen, Denmark: University of Copenhagen, 1963.
- 37 Hølmer P, Søndergaard L, Konradsen L, et al. Epidemiology of sprains in the lateral ankle and foot. *Foot Ankle* 1994;15:72-4.
- 38 Meyer JM, Garcia J, Hoffmeyer P, et al. The subtalar sprain. A roentgenographic study. *Clin Orthop* 1988;226:169-73.
- 39 Taillard W, Meyer JM, Garcia J, et al. The sinus tarsi syndrome. *Int Orthop* 1981;5:117-30.
- 40 Taga I, Shino K, Inoue M, et al. Articular cartilage lesions in ankles with lateral ligament injury. *Am J Sports Med* 1993;21:120-6.
- 41 Grana WA. Chronic pain persisting after ankle sprain. *Journal of Musculoskeletal Medicine* 1990;7:35-49.
- 42 Sammarco GJ. Peroneal tendon injuries. *Orthop Clin North Am* 1994;25:109-22.
- 43 Hyslop GH. Injuries to the deep and the superficial peroneal nerves complicating ankle sprain. *Am J Surg* 1941;51:436-8.
- 44 Stoff MD, Greene AF. Common nerve palsy following inversion ankle injury: a report of two cases. *Physical Therapy* 1982;62:1463-4.
- 45 Nitz AJ, Dobner JJ, Kersey D. Nerve injury and grades II and III ankle sprains. *Am J Sports Med* 1985;13:177-82.
- 46 Kleinrensink GJ, Stoekart R, Meulstee J, et al. Lowered motor conduction velocity of the peroneal nerve after inversion trauma. *Med Sci Sports Exer* 1994;26:877-83.
- 47 Norfray JF, Groves HC, Rogers LF, et al. Common calcaneal avulsion fracture. *Am J Roentgenol* 1980;134:119-23.
- 48 Heckman JD, McLean MR. Fractures of the lateral process of the talus. *Clin Orthop* 1985;199:108-13.
- 49 Noble J, Royle SG. Fracture of the lateral process of the talus: computed tomographic scan diagnosis. *Br J Sports Med* 1992;26:245-6.
- 50 Lindenfeld TN. The differentiation and treatment of ankle sprains. *Orthopedics* 1988;11:203-6.
- 51 Stepanuk M. Arthrography of the ankle joint in the diagnosis of acute ankle sprains. *Journal of the American Orthopedic Association* 1977;76:52936.
- 52 Cass JR, Morrey BF. Ankle instability: current concepts, diagnosis, and treatment. *Mayo Clin Proc* 1984;59:165-70.
- 53 Van der Ent FWC. *Lateral ankle ligament injury*. Rotterdam: Elinkwijk, Utrecht, 1984.
- 54 Nilsson S. Sprains of the lateral ankle ligaments. An epidemiological and clinical study with special reference to different forms of conservative treatment. Part I. *Journal of Oslo City Hospital* 1982;32:3-29.
- 55 Prins JG. Diagnosis and treatment of injury to the lateral ligament of the ankle. A comparative clinical study. *Acta Chir Scand* 1978;Suppl. 486.
- 56 Stiell IG, McKnight RD, Greenberg GH, et al. Interobserver agreement in the examination of acute ankle injury patients. *Am J Emerg Med* 1992;10:14-17.
- 57 van Dijk CN, Lim LSL, Bossuyt PMM, et al. Physical examination is sufficient for the diagnosis of sprained ankles. *J Bone Joint Surg [Br]* 1996;78:958-62.
- 58 Stiell IG, McKnight RD, Greenberg GH, et al. Implementation of the Ottawa ankle rules [see comments]. *JAMA* 1994;271:827-32.
- 59 Jackson DW, Ashley RL, Powell JW. Ankle sprains in young athletes. Relation of severity and disability. *Clin Orthop* 1974;101:201-15.
- 60 Clark BL, Derby AC, Power GRI. Injuries of the lateral ligament of the ankle. Conservative vs. operative repair. *Can J Surg* 1965;8:358-63.
- 61 Evans GA, Hardcastle P, Frenyo AD. Acute rupture of the lateral ligament of the ankle. To suture or not to suture? *J Bone Joint Surg [Br]* 1984;66:209-12.
- 62 Freeman MAR. Treatment of ruptures of the lateral ligament of the ankle. *J Bone Joint Surg [Br]* 1965;47:661-8.
- 63 Grønmark T, Johnsen O, Kogstad O. Rupture of the lateral ligament of the ankle: a controlled clinical trial. *Injury* 1980;11:215-8.
- 64 Klein J, Schreckenberger C, Roddecker K. Operative or conservative treatment of recent rupture of the fibular ligament in the ankle. A randomized clinical trial. *Unfallchirurgie* 1988;91:154-60.
- 65 Korkala O, Rusanen M, Jokipii P, et al. A prospective study of the treatment of severe tears of the lateral ligament of the ankle. *Int Orthop* 1987;11:13-17.
- 66 Møller-Larsen F, Wethelund JO, Jurik AG, et al. Comparison of three different treatments for ruptured lateral ankle ligaments. *Acta Orthop Scand* 1988;59:564-6.
- 67 Niedermann B, Andersen A, Andersen SB, et al. Rupture of the lateral ligaments of the ankle: operation or plaster cast? A prospective study. *Acta Orthop Scand* 1981;52:579-87.
- 68 Sommer HM, Arza D. Functional treatment of recent ruptures of the fibular ligament of the ankle. *Int Orthop* 1989;13:157-60.
- 69 Van Moppes FI, Van den Hoogenband CR. *Diagnostic and therapeutic aspects of inversion trauma of the ankle joint*. Amsterdam, Holland: Free University, 1982.
- 70 Freeman MA. Instability of the foot after injuries to the lateral ligament of the ankle. *J Bone Joint Surg [Br]* 1965;47:669-77.
- 71 Korkala O, Laittamus L, Tanskanen P. Lateral ligament injuries of the ankle. Results of primary surgical treatment. *Ann Chir Gynaecol* 1982;71:161-3.
- 72 Konradsen L, Hølmer P, Søndergaard L. Early mobilizing treatment for grade III ankle ligament injuries. *Foot Ankle* 1991;12:69-73.
- 73 Eifff MP, Smith AT, Smith GE. Early mobilization versus immobilization in the treatment of lateral ankle sprains. *Am J Sports Med* 1994;22:83-8.
- 74 Kaikkonen A, Kannus P, Järvinen M. A performance test protocol and scoring scale for the evaluation of ankle injuries. *Am J Sports Med* 1994;22:462-9.
- 75 Schrier I. Treatment of lateral collateral ligament sprains of the ankle: a critical appraisal of the literature. *Clin J Sport Med* 1995;5:187-95.
- 76 Sommer HM, Schreiber T. Early functional conservative therapy of a fresh fibular rupture of the capsular ligament from a socioeconomic point of view. *Sportverletz Sportschaden* 1993;7:40-6.
- 77 Leanderson J, Wredmark T. Treatment of acute ankle sprain. *Acta Orthop Scand* 1995;66:529-31.
- 78 Leach RE, Schepesis AA. Acute injury to ligaments of the ankle. In: Evans CM, ed. *Surgery of the musculoskeletal system*. Vol 4. New York: Churchill Livingstone, 1990: 3887-913.
- 79 Burry HC. Soft tissue injury in sport. *Exerc Sport Sci Rev* 1975;3:275-301.
- 80 Järvinen M, Letho M. The effect of early mobilization and immobilization on the healing process following muscle injuries. *Sports Med* 1993;15:78-89.
- 81 Järvinen M. *Healing of a crush injury in rat striated muscle with special reference to treatment of early mobilization and immobilization*. Turku, Finland: University of Turku, 1976.

- 82 Letho M. *Collagen and fibronectin in healing skeletal muscle injury: an experimental study in rats under variable conditions of physical exercise*. Turku, Finland: University of Turku, 1983.
- 83 Montgomery JB, Steadman JR. Rehabilitation of the injured knee. *Clin Sports Med* 1985;4:333-43.
- 84 Akeson WH, Amiel D, Woo SL-Y. Immobility effects on synovial joints: the pathomechanics of joint contracture. *Biorheology* 1980;17:95-100.
- 85 Alves J, Alday R, Ketcham D, Lentell G. A comparison of the passive support provided by various ankle braces. *J Orthop Sports Phys Ther* 1992;15:10-18.
- 86 Jozsa L, Järvinen M, Kannus P, et al. Fine structural changes in the articular cartilage of the rat's knee following short-term immobilisation in various positions: a scanning electron microscopical study. *Int Orthop* 1987;11:129-33.
- 87 Jozsa L, Thoring J, Järvinen M, et al. Quantitative alterations in intramuscular connective tissue following immobilization: an experimental study in the rat calf muscles. *Exp Mol Pathol* 1988;49:267-78.
- 88 Jozsa L, Reffy A, Järvinen M, et al. Cortical and trabecular osteopenia after immobilization. A quantitative histological study of the rat knee. *Int Orthop* 1988;12:169-72.
- 89 Noyes FR. Functional properties of knee ligaments and alterations induced by immobilization. *Clin Orthop* 1977;123:210-42.
- 90 Ogata K, Whiteside LA, Andersen DA. The intra-articular effect of various postoperative managements following knee ligament repair: an experimental study in dogs. *Clin Orthop* 1980;150:271-6.
- 91 Woo SL, Inoue M, McGurk-Burleson E, et al. Treatment of the medial collateral ligament injury. II: Structure and function of canine knees in response to differing treatment regimens. *Am J Sports Med* 1987;15:22-9.
- 92 Renström P, Kannus P. Injuries of the foot and ankle. In: DeLee JC, Drez DJ, eds. *Orthopaedic sports medicine: principles and practice*. Vol 2. Philadelphia: Saunders, 1994: 1705-67.
- 93 Karlsson J, Bergsten T, Lansinger O, et al. Reconstruction of the lateral ligaments of the ankle for chronic lateral instability. *J Bone Joint Surg [Am]* 1988;70:581-8.
- 94 Mann G, Eliashuv O, Perry C. Recurrent ankle sprain: literature review. *Israel Journal of Sports Medicine* 1994;4: 104-13.
- 95 Tropp H. *Functional instability of the ankle joint*. Linköping, Sweden: Linköping University, 1985.
- 96 Freeman MAR, Dean MRE, Hanham IWF. The etiology and prevention of functional instability of the foot. *J Bone Joint Surg [Br]* 1965;47:678-85.
- 97 Harper MC. The lateral ligamentous support of the subtalar joint (review). *Foot Ankle* 1991;11:354-8.
- 98 Konradsen L, Ravn JB. Prolonged peroneal reaction time in ankle instability. *Int J Sports Med* 1991;12:290-2.
- 99 Karlsson J, Lansinger O. Lateral instability of the ankle joint. *Clin Orthop* 1992;276:253-61.
- 100 Löfvenberg R, Kärrholm J, Sundelin G. Prolonged reaction time in patients with chronic lateral instability of the ankle. *Am J Sports Med* 1995;23:414-7.
- 101 Isakov E, Mizrahi J, Solzi P, et al. Response of the peroneal muscles to sudden inversion of the ankle during standing. *International Journal of Sports Biomechanics* 1986;2:100-9.
- 102 Brunt D, Andersen JC, Huntsman B, et al. Postural responses to lateral perturbation in healthy subjects and ankle sprain patients. *Med Sci Sports Exerc* 1992;24:171-6.
- 103 Johnson MB, Johnson CL. Electromyographic response of peroneal muscles in surgical and nonsurgical injured ankles during sudden inversion. *Journal of Sports Physical Therapy* 1993;18:497-501.
- 104 Glencross D, Thornton E. Position sense following joint injury. *J Sports Med Phys Fitness* 1981;21:23-7.
- 105 Garn SN, Newton RA. Kinesthetic awareness in subjects with multiple ankle sprains. *Phys Ther* 1988;68:1667-71.
- 106 Jerosch J, Hoffstetter I, Bork H, et al. The influence of orthoses on the proprioception of the ankle joint. *Knee Surg Sports Traumatol Arthrosc* 1995;3:39-46.
- 107 Gross MT. Effects of recurrent lateral ankle sprains on active and passive judgement of joint position. *Phys Ther* 1987;67: 1505-9.
- 108 Leanderson J, Eriksson E, Nilsson C, et al. Proprioception in classical ballet dancers: a prospective study of the influence of an ankle sprain on proprioception in the ankle joint. *Am J Sports Med* 1996;24:370-4.
- 109 Chrisman OD, Snook GA. Reconstruction of lateral ligament tears of the ankle. *J Bone Joint Surg [Am]* 1969;51:904-12.
- 110 Clanton TO. Instability of the subtalar joint (review). *Orthop Clin North Am* 1989;2:583-92.
- 111 Karlsson J, Renström P. Subtalar ankle instability. *Scand J Sports Med* (in press).
- 112 Grimby G. Clinical aspects of strength and power training. In: Komi PV, eds. *Strength and power in sport*. Oxford: Blackwell Scientific, 1992: 338-56.
- 113 Karlsson J, Lansinger O. Chronic lateral instability of the ankle in athletes. *Sports Med* 1993;16:355-65.
- 114 Ashton-Miller JA, Ottaviani RA, Hutchinson C, et al. What best protects the inverted weightbearing ankle against further inversion. *Am J Sports Med* 1996;24:800-9.
- 114a Garrick JG, Requa RK. The role of external support in prevention of ankle sprains. *Med Sci Sports* 1973;5:200-3.
- 115 Lindenberger U, Reese D, Andreasson G, et al. *The effect of prophylactic taping of ankles*. Gothenburg, Sweden: Chalmers Technical University, 1985.
- 116 Fumich RM, Ellison AE, Guerin GJ et al. The measured effect of taping on combined foot and ankle motion before and after exercise. *Am J Sports Med* 1981;9:165-70.
- 117 Rarick GL, Bigley G, Karst R. The measurable support of the ankle joint by conventional methods of taping. *J Bone Joint Surg [Am]* 1962;44:1183-90.
- 118 Larsen E. Taping the ankle for chronic instability. *Acta Orthop Scand* 1984;55:551-3.
- 119 Myburgh KH, Vaughan CL, Isaacs SK. The effect of ankle guards and taping on joint motion before, during, and after a squash match. *Am J Sports Med* 1984;12:441-6.
- 120 Karlsson J, Bergsten T, Lansinger O, et al. Surgical treatment of chronic lateral instability of the ankle joint. A new procedure (review). *Am J Sports Med* 1989;17:268-73;discussion 273-4.
- 121 Shapiro MS, Kabo JM, Mitchell PW, et al. Ankle sprain prophylaxis: an analysis of the stabilizing effects of braces and tape. *Am J Sports Med* 1994;22:78-82.
- 122 Rovere GD, Clarke TJ, Yates CS. Retrospective comparison of taping and ankle stabilizers in preventing ankle injuries. *Am J Sports Med* 1988;16:228-33.
- 123 Sittler M, Ryan J, Wheeler B, et al. The efficacy of a semi-rigid ankle stabilizer to reduce acute ankle injuries in basketball. A randomized clinical study at West Point. *Am J Sports Med* 1994;22:454-61.
- 124 Nishikawa T, Grabiner MD. *Ankle braces influence the reflex amplitude of muscle in response to stretch*. 42nd Annual Meeting, Orthopaedic Research Society, 1996.
- 125 Greene TA, Wright CR. A comparative support evaluation of three orthoses before, during and after exercise. *J Orthop Sports Phys Ther* 1990;11:453-66.
- 126 Burks RT, Marcus R, Bean BG, Barker HB. Analysis of athletic performance with prophylactic ankle devices. *Am J Sports Med* 1991;19:104-6.
- 127 Coffman JL, Mize NL. A comparison of ankle taping and the Aircast sport-stirrup on athletic performance. *Athletic Training* 1989;24:123-8.
- 128 Gross MT, Everts JR, Roberson SE. Effect of DonJoy ankle ligament protector and Aircast sport-stirrup orthoses on functional performance. *J Orthop Sports Phys Ther* 1994;19:150-6.
- 129 Barrett J, Bilisko T. The role of shoes in the prevention of ankle sprains. *Sports Med* 1995;20:227-80.
- 130 Cote DJ, Prentice WE Jr, Hooker DN, et al. Comparison of three treatment procedures for minimizing ankle sprain swelling. *Phys Ther* 1988;68:1072-6.
- 131 Airaksinen O, Kolari PJ, Herve R, et al. Treatment of post-traumatic oedema in lower legs using intermittent pneumatic compression. *Scand J Rehabil Med* 1988;20:25-8.
- 132 Airaksinen O. Changes in posttraumatic ankle joint mobility, pain, and edema following intermittent pneumatic compression therapy. *Arch Phys Med Rehabil* 1989;70:341-4.
- 133 Tohyama H, Beynon BD, Renstrom P, Pope MH. *Laxity and stiffness of normal and reconstructed ankles using Chrisman-Snook procedure*. 61st Annual Meeting, American Academy of Orthopaedic Surgeons.
- 134 Althoff B, Peterson L, Renstrom P. A simple reconstructive procedure of chronic ligament injuries of the ankle. *Laker-tidningen* 1981;78:2857-61.