- competition cyclists? Report of two cases. *Vascular Surgery* 1985;19:258–63.
- 3 Abraham P, Leftheriotis G, Bourre Y, Chevalier J, Saumet J. Echography of external iliac artery endofibrosis in cyclists. *Am J Sports Med* 1993;21:861–3.
  - 4 Johnstone MT, Creager SJ, Scales KM, Cusco JA, Byron KL, Creager MA. Impaired endothelium-dependent vasodilation in patients with insulin dependent diabetes mellitus. *Circulation* 1993;88:2510–16.
  - 5 Ramsey MW, Goodfellow J, Jones CJH, Luddington LA, Lewis MJ, Henderson AH. Endothelial control of arterial distensibility is impaired in congestive heart failure. *Circulation* 1995;92:3212–19.
  - 6 Luscher T, Lie J, Stanson A, Houser O, Hollier L, Sheps S. Arterial fibromuscular dysplasia. *Mayo Clin Proc* 1987;62:931–52.
  - 7 Astrand P, Cuddy T, Saltin B, Stenberg J. Cardiac output during submaximal and maximal work. *J Appl Physiol* 1964;19:268–74.
  - 8 Griffith T. Modulation of blood flow and tissue perfusion by endothelium-derived relaxing factor. *Exp Physiol* 1994;79:873–913.
  - 9 Joannides R, Haefeli WE, Lilylinder VR, El Hassan B, Thuilliez C, Luscher T. Nitric oxide is responsible for flow dependent dilatation of human peripheral conduit arteries in vivo. *Circulation* 1995;91:1314–19.
  - 10 Goodfellow J, Ramsey M, Luddington LA, Jones CJH, Owens DR, Lewis MJ, *et al.* Flow-related endothelial dilatation is impaired in non-insulin dependent diabetics. *BMJ* 1996;312:744–5.

Commentary

As a racing cyclist myself, I am only too aware of the frequency with which competitors land on their buttocks during competitive spills, particularly while riding in track events. Fortunately, these accidents usually only produce bruises, abrasions, and multiple splinters when wooden tracks are being used. It is indeed intriguing to think that there may be damage to the intima of the external iliac artery as a direct result of such trauma. It is fascinating to see a paper presenting a case of external iliac artery stenosis in a runner, albeit an insulin dependent diabetic competitor. As a clinician, I shall now consider more seriously the possibility of arterial causes when attempting to diagnose leg pain in competitors, particularly cyclists. Once again, these cases indicate the importance of taking a good clinical history and, where possible, assessing the athlete after exercise.

GRAHAME J BROWN