Physiological implications of altitude training for endurance performance at sea level: a review

Damian M Bailey, Bruce Davies

Summary

Acclimatisation to environmental hypoxia initiates a series of metabolic and musculoskeletal adaptations that influence oxygen transport and utilisation. Whilst it is clear that adequate acclimatisation, or better still, being born and raised at altitude, is necessary to achieve optimal physical performance at altitude, scientific evidence to support the potentiating effects after return to sea level is at present equivocal. Despite this, elite athletes continue to spend considerable time and resources training at altitude, misled by subjective coaching opinion and the inconclusive findings of a large number of uncontrolled studies. Scientific investigation has focused on the optimisation of the theoretically beneficial aspects of altitude acclimatisation, which include increases in blood haemoglobin concentration, elevated buffering capacity, and improvements in the structural and biochemical properties of skeletal muscle. However, not all aspects of altitude acclimatisation are beneficial; cardiac output and blood flow to skeletal muscles decrease, and preliminary evidence has shown that hypoxia in itself is responsible for a depression of immune function and increased tissue damage mediated by oxidative stress. Future research needs to focus on these less beneficial aspects of altitude training, the implications of which pose a threat to both the fitness and the health of the elite competitor.

Paul Bert was the first investigator to show that acclimatisation to a chronically reduced inspiratory partial pressure of oxygen (P_{iO_2}) invoked a series of central and peripheral adaptations that served to maintain adequate tissue oxygenation in healthy skeletal muscle,¹ physiological adaptations that have been subsequently implicated in the improvement in exercise performance during altitude acclimatisation. However, it was not until half a century later that scientists suggested that the additive stimulus of environmental hypoxia could potentially compound the normal physiological adaptations to endurance training and accelerate performance improvements after return to sea level. This has stimulated an exponential increase in scientific research, and, since 1984, 22 major reviews have summarised the physiological implications of altitude training for both aerobic and anaerobic performance at altitude and after return to sea level. Of these reviews, only eight have specifically focused on physical performance changes after return to sea level,²⁻⁹ the most comprehensive of which was recently written by Wolski *et al.*⁹

Few reviews have considered the potentially less favourable physiological responses to moderate altitude exposure, which include

decreases in absolute training intensity,¹⁰ decreased plasma volume,¹¹ depression of haemopoiesis and increased haemolysis,¹² increases in sympathetically mediated glycogen depletion at altitude,¹³ and increased respiratory muscle work after return to sea level.¹⁴ In addition, there is a risk of developing more serious medical complications at altitude, which include acute mountain sickness, pulmonary oedema, cardiac arrhythmias, and cerebral hypoxia.¹⁵ The possible implications of changes in immune function at altitude have also been largely ignored, despite accumulating evidence of hypoxia mediated immunosuppression.¹⁶

In general, altitude training has been shown to improve performance at altitude, whereas no unequivocal evidence exists to support the claim that performance at sea level is improved. Table 1 summarises the theoretical advantages and disadvantages of altitude training for sea level performance.

This review summarises the physiological rationale for altitude training as a means of enhancing endurance performance after return to sea level. Factors that have been shown to affect the acclimatisation process and the subsequent implications for exercise performance at sea level will also be discussed.

Studies were located using five major database searches, which included Medline, Embase, Science Citation Index, Sports Discus, and Sport, in addition to extensive hand searching and cross referencing. All published English studies, dating back from the present day to 1956, that included physiological measurements during exercise before and after hypoxic training were incorporated in the overall analysis. Ninety one investigations were selected, which included 772 hypoxically trained experimental and 209 normoxically trained control subjects.

The investigations were subdivided according to whether a normoxically trained control group was incorporated into the experimental design. Other classifications were made depending on the characteristics of the hypoxic stimulus, which included type (normobaric or hypobaric hypoxia; continuous or intermittent), duration, and magnitude (calculated ambient PO_2), and timing of physiological testing after the descent to sea level.

The continued popularity of altitude training has been influenced by two factors. Firstly, hypoxia in itself increases blood haemoglobin (Hb) concentration, which has been shown to improve endurance performance. Secondly, several of the best endurance runners in the world have originated from East African countries that are based at altitude (1500–2000 m). Is it possible that either living and/or training at

School of Applied Sciences, University of Glamorgan, Pontypridd, Mid-Glamorgan, United Kingdom
D M Bailey
B Davies

Correspondence to:
D M Bailey.

Accepted for publication
8 May 1997

Table 1 Physiological changes during altitude acclimatisation in native lowlanders; time course and theoretical implications for exercise performance at sea level

Physiological advantages	Response time	Physiological disadvantages	Response time
Increased free fatty acid mobilisation	Weeks	Increased ventilation	Immediate
Increased haemoglobin	Days	Decreased cardiac output	Days
Increased capillarity	Months/ years?	Decreased blood flow	Days
Increased oxidative enzyme activity	Weeks	Immunosupression	Immediate?/ days
Increased mitochondrial volume	Weeks	Increased oxidative stress and tissue damage	Immediate
		Increased dehydration	Immediate
		Jet lag	Immediate
		Decreased training intensity	Immediate
		Acute mountain sickness	Days
		Sunburn due to increased ultraviolet B (290-320nm)	Immediate
		Catecholamine mediated glycogen depletion	Days-weeks
		Increased haemolysis	Weeks

altitude may contribute to their running success?

Physiological rationale for altitude training

AUTOLOGOUS BLOOD REINFUSION AND ENDURANCE PERFORMANCE

One of the most documented physiological adaptations to a reduced P_iO₂ is the increased release of erythropoietin, which causes a transient increase in red blood cell mass.¹⁷ The implications of secondary polycythaemia to both submaximal and maximal indices of endurance performance have been clearly shown by studies that have artificially induced erythrocythaemia after either autologous blood reinfusion¹⁸ or subcutaneous injections of recombinant human erythropoietin.^{19, 20} Table 2 summarises the major research findings. It has been reported that absolute maximal oxygen uptake ($\dot{V}O_{2MAX}$) values are increased by about 200 ml/min per g/dl increase in Hb, irrespective of the methods by which polycythaemia is induced.²¹

However, the use of blood doping as an ergogenic aid is considered unethical and potentially dangerous and is banned by the International Olympic Committee.²²

PHYSIOLOGICAL ADAPTATIONS OF THE NATIVE HIGHLANDER: A SUPERIOR ATHLETE?

Figure 1 illustrates the apparently disproportionate running success of the native highlander. This figure represents data obtained

from athletes who were born and raised at a median altitude of 2000 m above sea level. This phenomenon has prompted several comparative investigations into what, if any, physiological adaptations mediated by hypoxia could contribute to their superiority in distance running events. Much interest has focused on the four steps of the oxygen transport system, namely alveolar ventilation, lung diffusion, circulatory oxygen transport, and tissue oxygen extraction. Studies have shown that the native highlander is characterised by a larger pulmonary diffusion capacity³² and adaptations in the structural and metabolic organisation of skeletal muscle that result in a tighter coupling between ATP hydrolysis and oxidative phosphorylation.³³ These are the major factors that facilitate oxygen transport and utilisation. The significance of these adaptations has been elucidated in a series of investigations that have reported higher values for $\dot{V}O_{2MAX}$,³⁴ power output,³⁵ arterial oxygen saturation,³⁶ and cerebral oxygen delivery³⁷ during maximal exercise and decreased blood lactate^{33, 35} and ammonia concentrations³⁸ for a given submaximal work rate.

To what extent these physiological adaptations are acquired as the result of inheritance or hypobaric hypoxia is not well defined. The influence of genetic factors on quantitative oxygen transport was recently investigated in a unique study by Beall *et al.*³⁹ They identified a major gene that enhances arterial oxygen saturation in sedentary Tibetan natives. The physiological significance of this was shown by Niermeyer *et al.*,⁴⁰ who concluded that genetic adaptations to hypobaric hypoxia resulted in improved oxygenation and conferred resistance to subacute infantile mountain sickness. These adaptations were more pronounced in a cohort of Tibetan newborns whose ancestors have resided at altitude for 50 000 to 100 000 years, in comparison with Han newborns whose ancestors had resided at altitude for only 45 years.⁴¹ In general, these findings would suggest that a lifetime or perhaps generations of altitude exposure are responsible for the biological distinctiveness of the high altitude population.

Altitude training and sea level endurance performance in native lowlanders

Table 3 summarises the effects of altitude training on sea level endurance performance. The weight of scientific evidence does not support the potentiating effects of altitude training. However, it is becoming clearer that a number of methodological deficiencies may preclude the potential synergistic effects of hypoxia and physical exercise, the physiological implications of which will be discussed in the following sections.

Intensity and duration of the hypoxic stimulus and associated haematological adaptation

There is still much controversy about the optimal altitude and duration required for athletes to train in an attempt to optimise endurance performance at sea level. Much attention has

Table 2 Effects of autologous blood reinfusion on $\dot{V}O_{2MAX}$

Author/reference	Volume of blood reinfused (ml)	Change in Hb after reinfusion (%)	Change in $\dot{V}O_{2MAX}$ after reinfusion (%)
Ekblom ¹⁹	1350	+9*	+8*
Celsing ²¹	2250	+11	+7**
Buick ²³	900	+8**	+5**
Spriet ²⁴	1200	NR	+7*
Williams ²⁵	920	+7**	NR
Goforth ²⁶	760	+4*	+11**
Robertson ²⁷	750	+28*	+13*
Robertson ²⁸	475	+16*	+10*
Thompson ²⁹	1000	+12*	+11*
Sawka ³⁰	600	+10*	+11*
Robertson ³¹	475	+16*	+10**

* Significantly different from before reinfusion (P < 0.05).
** Significantly different from before reinfusion (P < 0.01).
NR, not reported.

Figure 1 Percentage of Olympic medals for middle and long distance running events won by native highlanders since 1960.

Table 3 Effects of hypoxic training on sea level endurance performance

Hypoxic stimulus	Altitude (m)	Exposure time (days)	Time tested after altitude (days)	Submaximal improvement	Change in $\dot{V}O_{2\text{MAX}}$ (%)	Control group
Potentiating effects						
CH ⁴²	1300–2500	28	1	Yes†	+4†	No
CH ⁴³	1900	21	1/14	Yes†	NS/NS	No
CH ⁴⁴	2100–2700	14	2	Yes†	NS	No
CH ⁴⁵	2300	23	3/21	Yes†	+8†/+10†	No
CH ⁴⁶	2500	28	7	ND	+6†	No
IH ⁴⁷	3049–4268	23	3–4	Yes†	NS	No
CH ⁴⁸	3800	35	14	ND	+14†	No
IH ⁴⁹	4020	21/28	1	Yes†	+8†/+26‡	No
CH ⁵⁰	1250–2500	28	7	Yes†	+4†	Yes
IH ⁵¹	2300	21–28	1	Yes†	NS	Yes
IH ⁵²	2300	28	1–2	Yes†	ND	Yes
IH ⁵³	4000	70	1	Yes†	NS	Yes
No potentiating effects						
CH ⁴⁵	2300	14	1	No	NS	No
CH ⁵⁴	1695–2700	7	4	ND	NS	No
CH ⁵⁵	2240	20	4/22	ND	+6/+9*	No
CH ⁵⁶	2300	42	4–5	No	NS	No
CH ⁵⁷	2300	70	5	ND	NS	No
CH ⁵⁸	2800	10	2–4	No	+7*	No
CH ⁵⁹	3090	17	1	ND	NS	No
CH ⁶⁰	3110	21	7	ND	–5*	No
CH ⁶¹	4000	48–63	2–15	No	NS	No
IH ⁶²	4000	21	1	ND	NS	No
CH ³⁸	2000	14	6/12	ND	NS/NS	Yes
CH ⁶³	1600–1800	18–28	7	ND	NS	Yes
CH ⁶⁴	1640	28	20	No	NS	Yes
CH ⁶⁵	1700–2000	28	7	No	NS	Yes
CH ⁶⁶	2300	21	1	No	NS	Yes
IH ⁶⁷	2250	28	1	ND	+17.5*	Yes
IH ⁶⁷	3450	28	1	ND	+10.0*	Yes
IH ⁶⁸	2500	28	1	No	NS	Yes
IH ⁶⁹	2500	35	1	ND	NS	Yes
CH ⁷⁰	2600	11	1	ND	NS	Yes
IH ⁷¹	3100	19	6	No	NS	Yes
IH ⁷²	3345	42	1	ND	NS	Yes
IH ⁷³	4020	15	1	ND	NS	Yes
IH ⁷⁴	4100–5700	21	1	ND	NS	Yes
CH ⁷⁵	4300	28	1–5	ND	NS	Yes

CH, Chronic hypobaric; IH, intermittent hypobaric; ND, no data.
* Level of significance not reported.
† Significantly different from pre-altitude value ($P<0.05$).
‡ Significantly different from pre-altitude value ($P<0.01$).
NS, not significantly different from pre-altitude value ($P>0.05$).

focused on the erythropoietic response to hypoxia and subsequent haematological adaptation. Considering the inverse relationship between PO_2 and resting Hb concentration,⁷⁶ it would seem logical that the higher the athlete

can train the better. However, other factors that inhibit exercise performance are exacerbated with a reduction in PO_2 . Acute mountain sickness presents at altitudes above 2000 to 3000 m,⁷⁷ with the possibility of the elite athlete suffering physiological symptoms at even lower altitudes.¹⁵ Prolonged exposure to altitudes above 4500 m has been shown to result in a reduction in muscle mass, the underlying physiological mechanisms for which have been recently reviewed by Kayser.⁷⁸ Finally, the effects of training at a lower PO_2 may result in a reduction in work rate, so that detraining may override the potential benefits of altitude acclimatisation.⁷⁹

Hypoxia and detraining

A recent study has shown that $\dot{V}O_{2\text{MAX}}$ is significantly reduced at an altitude as low as 610 m above sea level in elite endurance athletes.⁸⁰ This is a phenomenon peculiar to about 50% of trained subjects, with $\dot{V}O_{2\text{MAX}}$ values of above 65 ml/kg per min or 4 litres/min.⁸¹ These elite athletes develop more severe levels of arterial hypoxaemia during maximal and submaximal exercise than sedentary controls both under normoxic and hypoxic conditions.^{82 83} Several mechanisms have been proposed to explain these findings, which include hypoventilation, venoarterial shunting, ventilation-perfusion inequality, and an alveolar-capillary diffusion limitation.^{84 85}

These observations led early investigators to hypothesise that altitude exposure may result in a detraining response.⁷⁹ Daniels and Oldridge⁵⁷ have shown the importance of training intensity at altitude and its effects on sea level performance. They suggested that intermittent exposures to altitudes of 2300 to 3300 m and sea level optimised the balance between hypoxic acclimatisation and training intensity. Despite the experimental limitations of a single group design, two world records and 12 personal best times were recorded by athletes on return to sea level, which presented a reasonable endorsement for such an approach. However, from our experience, it is equally possible to have expected similar improvements in a control group training at sea level.⁶⁴

The detraining effect induced during chronic exposure to hypobaric hypoxia has been quantified in a sequence of studies by Levine *et al.*^{42 50 68} In their most recent study,⁵⁰ 39 competitive runners were randomly assigned to four weeks of (a) living high (2500 m) and training low (1250 m), (b) living high (2500 m) and training high (2500 m), or (c) living low (150 m) and training low (150 m). They showed that, although $\dot{V}O_{2\text{MAX}}$ values significantly improved 5 km race performance times by 4% in the two altitude trained groups, the running velocity that corresponded to $\dot{V}O_{2\text{MAX}}$ and the ventilatory threshold at sea level were significantly improved only in the group that lived high and trained low. An unusual finding was that 5 km performance time was 31 seconds slower in the sea level control group, which would suggest that the training stimulus was not absolutely controlled during the experimental period. Nevertheless, it was

concluded that the potentiating effects of altitude training were due to a high altitude acclimatisation effect (improved haematology) and a low altitude training effect (increased training intensity). Thus the authors advocated the practice of living high and training low as the optimal approach to altitude training. This has popularised the use of "altitude houses" recently developed in Finland which are portable hypobaric chambers used by elite athletes, who alternate living and sleeping at simulated altitude with normobaric training.⁸⁶ However, the effectiveness of this procedure should at present be considered equivocal, and further scientific investigation is warranted to endorse this approach to altitude training.

Concept of a critical P_{O_2} and haematological adaptation

Few athletes can afford the costs inherent in a "live high, train low" approach to altitude training. Therefore is it possible that a "threshold" altitude exists that optimises the benefits of haematological acclimatisation and minimises the negative effects of detraining? Weil *et al.*⁸⁷ have presented the most comprehensive evidence indicating the existence of such a threshold, albeit in sedentary highland natives (B Levine, personal communication). They identified a biphasic relationship between the arterial partial pressure of oxygen (P_{aO_2}) and red blood cell mass, and shown a clear inflection point at a "critical" P_{aO_2} of 67 mm Hg, equivalent to an interpolated arterial oxygen saturation of 92%. This point corresponds to the steeper portion of the oxygen-Hb dissociation curve. The equivalent P_{O_2} would equate to about 135 mm Hg, which is comparable with an altitude of 2200–2500 m above sea level required to stimulate sufficient haemopoiesis at rest to influence endurance performance.⁴ However, it has been shown that the decrement in $\dot{V}O_{2MAX}$ measured in hypobaric hypoxia is directly proportional to $\dot{V}O_{2MAX}$ measured in normoxia.⁸⁸ This would suggest that elite athletes are more prone to developing arterial hypoxaemia and may gain more benefit haematologically by training at lower altitudes in comparison with sedentary controls. This contention was supported by Ingjer *et al.*,⁴³ who showed that three weeks of altitude training at 1900 m in elite cross country skiers was sufficient to elevate Hb by 5% ($P < 0.02$) and decrease blood lactate concentration during a standardised submaximal test, despite no changes in $\dot{V}O_{2MAX}$. However, it should be noted that these authors did not measure their subjects' plasma volumes, and their comments that the polycythaemia was independent of a haemoconcentration remains only speculative. The scarcity of training studies conducted at moderate altitudes of 1500 to 2000 m in elite athletes does not allow definitive conclusions to be made.

Optimal duration

Few data are available on the optimal time an athlete should spend training at altitude. On the basis of subjective coaching opinion as opposed to objective scientific evidence, it

would appear that three weeks are sufficient to gain a performance advantage at sea level.⁸⁹ However, the longer the duration of the hypoxic stimulus the greater the erythropoietic response and associated haematological adaptation.¹⁷ This was shown by Berglund,⁹⁰ who summarised the haematological changes during previous altitude training studies conducted between 1829 and 3048 m. He identified a "true" increase in Hb concentration of 1% per week, which was independent of a haemoconcentration. Thus, assuming that the detraining response could be minimised and polycythaemia did not approach pathological values, the longer the athlete spends at altitude, the greater the potential benefit for endurance performance.

Iron status during altitude training

Hypoxia in itself increases iron demand and mobilisation,^{91, 92} such that endurance athletes training at altitude may be prone to iron deficiency. Lack of this critical erythropoietic factor has been shown to inhibit complete haematological adaptation.⁹³ Despite its importance, few studies have actually reported iron status of athletes during their hypoxic exposure. Suboptimal iron stores may account for the vast majority of training studies that have failed to show increases in Hb concentration and endurance performance on return to sea level after the hypoxic exposure. The differences in iron status may also characterise the highly individualised haematological responses observed during altitude training.⁴³

Interval between descent and event

There is some evidence to suggest that endurance performance is affected by the timing of the descent to sea level after a sojourn to altitude. The general consensus amongst top coaches would suggest that endurance performance is optimised after 14 days at sea level after a bout of altitude training,⁸⁹ yet there is no scientific evidence to support this claim. Suslov⁹⁴ characterised the undulating nature of endurance performance after altitude training. His research was based on over 1000 competitive track results obtained from middle and long distance runners after different periods of altitude training (1300–2500 m) and repeated sea level $\dot{V}O_{2MAX}$ tests conducted after training at 1800 m. He identified a decrease in competition performance during the first two days at sea level and the first phase of enhanced work capacity occurring between days 3 and 7, followed by a decrease between days 8 and 10. Performance was shown to continue to improve between days 12 and 13, with the best results achieved on days 18 to 20. He also identified an additional upsurge in performance between days 36 and 48 after altitude. He failed to identify the physiological mechanisms responsible for this phenomenon.

Few studies have tested subjects on more than one occasion after return to sea level. Asahina *et al.*⁹⁵ and Faulkner *et al.*⁹⁶ did not show any significant changes in $\dot{V}O_{2MAX}$ values after either 3 or 22 days at sea level. Ingjer *et al.*⁴³ showed that after a group of elite cross country

Figure 2 Number of hypoxic training studies conducted with or without a normoxically trained control group since 1950.

skiers had trained for three weeks at an altitude of 1900 m, submaximal blood lactate values were lower than pre-altitude values on day 1 but not day 14 at sea level. The authors concluded that a 0.8 g/dl increase in Hb concentration measured on day 1 was responsible for the observed improvement in submaximal exercise. However, their failure to quantify plasma volume and blood flow changes weakens the validity of their haematological findings. Svedenhag *et al*³⁸ studied a group of Swedish middle distance runners who trained for a period of two weeks at altitude (2000 m) and were tested after 6 and 12 days on return to sea level. They did not identify any significant changes in $\dot{V}O_{2\text{MAX}}$, maximal oxygen deficit, and submaximal blood lactate values compared with pre-altitude values or between days 6 and 12 at sea level. However, they showed a significant reduction in heart rate, Borg rating of perceived exertion, and plasma ammonia concentration during a standardised submaximal treadmill test, which was more apparent after 12 days at sea level.

The physiological mechanisms responsible for these subtle changes in performance at sea level remain elusive. Intermittent altitude training has been shown to increase the hypoxic ventilatory response in a group of sedentary subjects, whereas an equivalent training programme at sea level had no effect.^{14 95} Acute exposure to altitude in the native lowlander may potentiate the hypoxic ventilatory response because of an increased peripheral chemoreceptor sensitivity, which would subsequently increase the work performed by the respiratory muscles. This has not been quantified in the elite athlete but may be implicated in the performance decrements shortly after return to sea level. Plasma volume has been shown to decrease by 25% during chronic exposure to hypobaric hypoxia¹¹ and may take as long as two months to normalise.⁹¹ After return to sea level, this may remain depressed for six days,⁹⁶ which may also negatively affect performance. Altitude training may also involve considerable travelling time, and the

negative impact of jet lag on exercise performance cannot be ignored.⁹⁷

Measurement of the altitude effect independent of training

Figure 2 shows that, since 1956, only 27 (30%) of the 91 hypoxic training studies reviewed have incorporated a normoxically trained control group. This makes it impossible to determine whether the physiological changes that occur after a bout of altitude training can be attributed to an improvement in physical conditioning or to the additive effects of hypoxia itself.

To our knowledge, the altitude training studies conducted by Asano *et al*,⁵³ Terrados *et al*,^{51 52} and Levine *et al*⁵⁰ would appear to be the only investigations employing a control group that have reported statistically significant improvements in aerobic performance after return to normoxia. Asano *et al*⁵³ studied ten elite middle to long distance male runners, who trained for a ten week period at the same relative exercise intensity at either sea level or a simulated altitude of 4000 m. After training, there were no improvements in $\dot{V}O_{2\text{MAX}}$ at sea level, yet 10 km personal best running times improved by about 6% ($P < 0.05$). Using a one legged training model, Terrados *et al*^{51 52} attributed the potentiating effects of intermittent hypobaric training to increases in citrate synthase activity and myoglobin content. The findings of Levine *et al*⁵⁰ have already been described in this review.

Whilst previous investigations have dealt primarily with aerobic responses to altitude training, there is some evidence to suggest that anaerobic performance is improved on return to sea level.^{44 98 99} Mizuno *et al*⁴⁴ showed that exercise time to exhaustion after altitude training improved by 17% ($P < 0.05$) when compared with pre-altitude values, which they attributed to a 6% increase ($P < 0.05$) in muscle buffer capacity. However, the validity of these findings is questionable because of the lack of a normoxically trained control group. A well controlled investigation by Martino *et al*,⁹⁸ which incorporated a performance matched control group based at sea level, investigated the effects of three weeks of altitude training at 2800 m on anaerobic measures of swimming performance. Sea level sprint performance time over 100 m was 2.4 seconds quicker in the altitude trained group than the control group ($P < 0.05$). The largest improvements in the altitude trained group were noted in an upper body Wingate test. Peak power output increased by 27.9 W more than the control group ($P < 0.05$). In a recent investigation, Nummela *et al*⁹⁹ showed that ten days of living high (~2200 m) and training low (sea level) resulted in greater improvements in 400 m running time ($P < 0.05$) and running velocity at a fixed concentration of blood lactate ($P < 0.05$) when compared with an equivalent programme of sea level training.

However, the vast majority of altitude training studies have not identified performance improvements at sea level. Whilst a decrease in absolute training intensity may be implicated,¹⁰

a decrease in muscle perfusion may also play a contributory role; oxygen transport, determined as a product of blood flow and arterial oxygen concentration is regulated during changes in P_{aO_2} . Reductions in blood flow during the inhalation of a hyperoxic gas mixture regulates oxygen delivery to the working muscles, such that total oxygen delivery is similar to that observed in normoxia¹⁰⁰. Auto-regulation of this mechanism has been investigated at altitude and after return to sea level.^{101 102} Whilst chronic exposure to hypobaric hypoxia increased arterial oxygen content as the result of an increase in Hb concentration, sympathetically mediated arterial vasoconstriction and a reduction in total cardiac output caused a reduction in blood flow, thus preventing an increase in oxygen transport.¹⁰¹ This decrease in muscle perfusion has been shown to persist after return to sea level. Using ¹³³Xe, blood flow to the vastus lateralis was shown to decrease by up to 39% ($P < 0.001$) during submaximal exercise after a three month expedition to 8398 m.¹⁰²

Favier *et al.*⁷² suggested that the negative findings reported in the literature could, in part, be attributed to the fact that subjects were not fully acclimatised to hypobaric hypoxia. In a unique experiment they used three groups of sedentary high altitude residents, who trained for 30 minutes a day at a constant work rate on a bicycle ergometer, during a six week period. Group 1 trained at a P_{O_2} that was equivalent to an interpolated altitude of 3345 m at 70% of $\dot{V}_{O_2MAX}$ determined in hypoxia. The remaining two groups trained under normoxic conditions at either the same relative work rate (70% of the normoxic $\dot{V}_{O_2MAX}$) or the same absolute work rate (70% of the hypoxic $\dot{V}_{O_2MAX}$) as the hypoxically trained group. An incremental test to exhaustion was performed by all groups in normoxia and hypoxia immediately before and after training in an attempt to ascertain the physiological responses to submaximal and maximal exercise. The authors showed that $\dot{V}_{O_2MAX}$ values improved similarly in all groups. However, they suggested that a lower reduction in base excess and bicarbonate stores observed in the hypoxically trained group could only potentially benefit anaerobic metabolism and, although time to exhaustion was not measured, facilitate exercise performance.

Hypoxia and immune function

Changes in total leucocyte, granulocyte, monocyte, lymphocyte, natural killer cell, and T cell count, helper/suppression cell ratio, cell proliferation in response to mitogens, and serum immunoglobulin levels have all been implicated in some form of immunosuppression, which may subsequently cause underperformance in the athlete at sea level.¹⁰³ The additive stress of a reduction in the inspiratory P_{O_2} , in conjunction with the extensive training loads employed by athletes at altitude, may explain why some investigators have reported physiological evidence for a less favourable modulation of immune function in vivo during acute and chronic exposure to hypobaric hypoxia.^{63 104–106}

Human studies have shown that chronic exposure to hypobaric hypoxia results in a suppression of cell mediated immunity, whereas B cell function remains unimpaired.¹⁶ Animal studies have further shown that murine host defences against bacterial pathogens are also impaired in hypoxia. The contributory immunomodulatory role of endogenous glucocorticoids and neuropeptides, which are increased at altitude, may contribute to the observed alterations in immune competence. In an experiment that employed elite distance runners and matched controls, we showed that plasma glutamine concentrations decreased significantly in comparison with pre-altitude values after 20 days of endurance training at an altitude of 1640 m above sea level ($P_{O_2} = 135$ mm Hg) (D M Bailey *et al.*, unpublished work). A reduction in glutamine concentration has been identified in "overtrained" athletes and may be a contributory factor leading to immunosuppression and underperformance.¹⁰⁷ It is difficult to comment on the physiological mechanisms responsible for these changes, but there is evidence that suggests that chronically elevated levels of circulating catecholamine levels decrease the rate of glutamine transport out of muscle incubated in vitro (Parry-Billings M, unpublished data). In addition to this, Wagenmakers¹⁰⁸ has proposed an alternative mechanism, again related to elevated catecholamine levels observed at altitude.¹¹ He suggested that hypoxia induced glycogen depletion would result in a reduction in the availability of tricarboxylic acid cycle intermediates, in particular 2-oxoglutarate. This is required for the activation of the branched chain amino acid aminotransferase reaction, which ultimately produces glutamine. The implications of the immunosuppressive influence of hypobaric hypoxia for endurance performance warrants further investigation in order to elucidate potential mechanisms that may modulate performance after return to sea level.

Reactive oxygen species at altitude

There is a limited body of evidence suggesting that oxidative injury mediated by free radicals is increased at altitude.^{109–111} Simon-Schnass¹⁰⁹ identified significant increases in indirect indices of free radical mediated lipid peroxidation at altitude, which included increased pentane excretion and thiobarbituric acid reactive substances, decreased erythrocyte filterability, and increased leucocyte and granulocyte counts. Daily supplementation with an antioxidant such as tocopherol (vitamin E) equivalent to 300–400 mg has been shown to improve endurance performance, by theoretically limiting tissue damage.^{109 110} An accelerated production of the highly toxic hydroxyl radical may occur as a consequence of an increased production of free iron derived from altitude induced and training induced destruction of red blood cells.¹¹¹ Thus it would appear that hypobaric hypoxia significantly increases oxidative stress, which has been shown to negatively influence energy metabolism and membrane integrity.

Summary and future research

Physiological acclimatisation to a chronically reduced P_{iO_2} is a prerequisite to achieve optimal physical performance in environmental hypoxia. However, scientific evidence to support the claim that either continuous or intermittent hypoxic training will enhance sea level performance remains at present equivocal. Future research should focus on methodological technicalities that optimise the balance between the favourable and less favourable responses to hypoxia and potential mediators of performance after return to sea level. Preliminary evidence showing that the additive stress of hypobaric hypoxia may provoke an adverse immune response and further potentiate free radical mediated oxidative injury has important implications which, if confirmed by scientific rigor, would present a threat to both the fitness and health of the elite competitor.

Our research was supported by the British Olympic Association and the British Athletics Federation. We would also like to acknowledge Mr N Papps of Ordnance Survey, UK, and Mr S Greenberg for their keen assistance in collecting data contained in fig 1.

- Bert P. *Barometric pressure: researches in experimental physiology*. Columbus, OH: College Book Co, 1943. (Translated by MA Hitchcock and RA Hitchcock.)
- Smith MH, Sharkey BJ. Altitude training: who benefits? *Physician and Sportsmedicine* 1984;12:48–62.
- Terrados N. Altitude training and muscular metabolism. *Int J Sports Med* 1992;13:S206–9.
- Levine BD, Stray-Gundersen J. A practical approach to altitude training: where to live and train for optimal performance enhancement. *Int J Sports Med* 1992;13:S209–12.
- Levine BD, Roach RC, Houston CS. Work and training at altitude. In: Sutton JR, Coates G, Houston CS, eds. *Hypoxia and mountain medicine*. Texas, VT: Queen City Printers, 1992:192–201.
- Sutton JR. Exercise training at high altitude: does it improve endurance performance at sea-level? *Sports Science Exchange* 1993;6:4.
- Hahn AG. The effect of altitude training on athletic performance at sea-level: a review. *Sports* 1993;13:1–7.
- Saltin B. Exercise and the environment: focus on altitude. *Res Q Exerc Sport* 1996;67:1–10.
- Wolski LA, McKenzie DC, Wenger HA. Altitude training for improvements in sea level performance. Is there scientific evidence of benefit? *Sports Med* 1996;22:251–63.
- Stine TA, Levine BD, Taylor S, Schultz W, Stray-Gundersen J. Quantification of altitude training in the field [abstract]. *Med Sci Sports Exerc* 1992;24:S103.
- Young AJ, Young PM. Human acclimatization to high terrestrial altitude. In: Pandolf KB, Sawka MN, Gonzalez PR, eds. *Human performance physiology and environmental medicine at terrestrial extremes*. Indianapolis: Benchmark Press, 1988:497–543.
- Szygula Z. Erythrocytic system under the influence of physical training and exercise. *Sports Med* 1990;10:181–97.
- Young A. Energy substrate utilization during exercise in extreme environments. *Exerc Sport Sci Rev* 1990;18:65–118.
- Levine BD, Friedman DB, Engfred K, Hanel B, Kjaer M, Clifford PS, Secher N. The effect of normoxic or hypobaric hypoxic endurance training on the hypoxic ventilatory response. *Med Sci Sports Exerc* 1992;24:769–75.
- Shepherd RJ. Problems of high altitude. In: Shepherd RJ, Astrand PO, eds. *Endurance in sport*. Oxford: Blackwell Science, 1992:471–8.
- Meehan RT. Immune suppression at high altitude. *Ann Emerg Med* 1987;16:974–9.
- Schmidt W, Spielvogel H, Eckardt KU, Quintela A, Penalzoza R. Effects of chronic hypoxia and exercise on plasma erythropoietin in high-altitude residents. *J Appl Physiol* 1993;74:1874–8.
- Gledhill N. The influence of altered blood volume and oxygen transport capacity on aerobic performance. *Exerc Sports Sci Rev* 1985;13:75–93.
- Eklund B, Berglund B. Effect of erythropoietin administration on maximal aerobic power. *Scand J Sports Med* 1991;1:88–93.
- Metra M, Cannella G, La Canna G. Improvement in exercise capacity after correction of anemia in patients with end stage renal failure. *Am J Cardiol* 1991;68:1060–6.
- Celsing F, Svedenahg J, Pilhstedt P, Eklund B. Effects of anaemia and stepwise-induced polycythemia on maximal aerobic power in individuals with high and low haemoglobin concentrations. *Acta Physiol Scand* 1987;31:47–54.
- American College of Sports Medicine. Position stand on the use of blood doping as an ergogenic aid. *Med Sci Sports Exerc* 1996;28:i-viii.
- Buick FJ, Gledhill N, Froese AB, Spriet L, Meyers EC. Effect of induced erythrocythemia on aerobic work capacity. *J Appl Physiol* 1980;48:636–42.
- Spriet LL, Gledhill N, Froese AB, Wilkes DL, Meyers EC. The effects of induced erythrocythemia on central circulation and oxygen transport during maximal exercise. *Med Sci Sports Exerc* 1980;12:122–3.
- Williams MH, Wesseldine S, Somma T, Schuster R. The effect of induced erythrocythemia upon 5-mile treadmill run time. *Med Sci Sports Exerc* 1981;13:169–75.
- Goforth HW, Campbell Jr NL, Hodgdon JA, Sucec AA. Hematologic parameters of trained distance runners following induced erythrocythemia. *Med Sci Sports Exerc* 1982;14:174.
- Robertson RJ, Gilcher R, Metz KF, Skrinar GS, Allison TG, Bahnson HT, et al. Effect of induced erythrocythemia on hypoxia tolerance during physical exercise. *J Appl Physiol* 1982;53:490–5.
- Robertson RJ, Gilcher R, Metz KF, Caspersen CJ, Allison TG, Abbott RA, et al. Hemoglobin concentration and aerobic work capacity in women following induced erythrocythemia. *J Appl Physiol* 1984;57:568–75.
- Thompson JM, Stone JA, Ginsburg AD, Hamilton P. O_2 transport during exercise following blood reinfusion. *J Appl Physiol* 1982;53:1213–19.
- Sawka MN, Dennis RC, Gonzalez RR, Young AJ, Muza SR, Martin JW, et al. Influence of polycythemia on blood volume and thermoregulation during exercise-heat stress. *J Appl Physiol* 1987;62:912–18.
- Robertson RJ, Gilcher R, Metz KF, Caspersen CJ, Allison TG, Abbott RA, et al. Effect of simulated altitude erythrocythemia in women on hemoglobin flow rate during exercise. *J Appl Physiol* 1988;64:1644–9.
- Dempsey JA, Reddan WG, Birnbaum ML. Effects of acute through life-long hypoxic exposure on exercise pulmonary gas exchange. *Respir Physiol* 1971;13:62–89.
- Hochachka PW, Stanley C, Matheson GO, McKenzie DC, Allen PS, Parkhouse WS. Metabolic and work efficiencies during exercise in Andean natives. *J Appl Physiol* 1991;70:1720–30.
- Sun SF, Droma TS, Zhang JG, Tao JX, Huang SY, McCullough RG, et al. Greater maximal O_2 uptake and vital capacities in Tibetan than Han residents of Lhasa. *Respir Physiol* 1990;79:151–62.
- Ge RL, Chen QH, Wang LH, Gen D, Yang P, Kubo K. Higher exercise performance and lower VO_2 max in Tibetan than Han residents at 4,700 m altitude. *J Appl Physiol* 1994;77:684–91.
- Favier R, Spielvogel H, Desplanches D, Ferretti G, Kayser B, Hoppeler H. Maximal exercise performance in chronic hypoxia and acute normoxia in high-altitude natives. *J Appl Physiol* 1995;78:1868–74.
- Huang SY, Sun S, Droma T, Zhuang J, Tao JX, McCullough RG, et al. Internal carotid arterial flow velocity during exercise in Tibetan and Han residents of Lhasa (3,658 m). *J Appl Physiol* 1992;73:2638–42.
- Svedenahg J, Saltin B, Johansson C, Kaijser L. Aerobic and anaerobic exercise capacities of elite middle-distance runners after two weeks of training at moderate altitude. *Scand J Med Sci Sports* 1991;1:205–14.
- Beall CM, Blangero J, Williams-Blangero S, Goldstein MC. Major gene for percent of oxygen saturation of arterial hemoglobin in Tibetan highlanders. *Am J Phys Anthropol* 1994;95:271–6.
- Niermeyer S, Yang P, Drolkar S, Zhuang J, Moore LG. Arterial oxygen saturation in Tibetan and Han infants born in Lhasa, Tibet. *N Engl J Med* 1995;333:1248–52.
- Ward MP, Milledge JS, West JB. High altitude medicine and physiology. 2nd ed. London: Chapman and Hall Medical, 1995:344.
- Levine BD, Stray-Gundersen J, Duhaime G, Snell PG, Friedman DB. Living high-training low: the effect of altitude acclimatization/normoxic training in trained runners. *Med Sci Sports Exerc* 1991;23:[abstract145].
- Ingjer F, Myhre K. Physiological effects of altitude training on elite male cross-country skiers. *J Sports Sci* 1992;10:37–47.
- Mizuno M, Juel C, Bro-Rasmussen T, Mygind E, Schibye B, Rasmussen B, Saltin B. Limb skeletal muscle adaptation in athletes after training at altitude. *J Appl Physiol* 1990;68:496–502.
- Faulkner JA, Daniels JT, Balke B. Effects of training at moderate altitude on physical performance capacity. *J Appl Physiol* 1967;23:85–9.
- Stray-Gundersen J, Mordecai N, Levine BD. O_2 transport response to altitude training in runners. *Med Sci Sports Exerc* 1996;27(S5):[abstract1131].
- Loeppky JA, Bynum WA. Effects of periodic exposure to hypobaric and exercise on physical work capacity. *J Sports Med Phys Fitness* 1970;10:238–47.
- Klausen K, Robinson S, Michael ED, Myhre LG. Effect of high altitude on maximal working capacity. *J Appl Physiol* 1966;21:1191–4.
- Banister EW, Woo W. Effects of simulated altitude training on aerobic and anaerobic power. *Eur J Appl Physiol* 1978;38:55–69.
- Levine BD, Friedmann B, Stray-Gundersen J. Confirmation of the “high-low” hypothesis: living at altitude-training near sea level improves sea level performance. *Med Sci Sports Exerc* 1996;28:[abstract742].
- Terrados N, Melichna J, Sylven C, Jansson E, Kaijser L. Effects of training at simulated altitude on performance

- and muscle metabolic capacity in competitive road cyclists. *Eur J Appl Physiol* 1988;57:203-9.
- 52 Terrados N, Jansson E, Sylvén C, Kaijser L. Is hypoxia a stimulus for synthesis of oxidative enzymes and myoglobin? *J Appl Physiol* 1990;68:2369-72.
 - 53 Asano K, Sub S, Matsuzaka A, Hirakoba K, Nagai J, Kawaoka T. The influences of simulated high altitude training on work capacity and performance in middle and long distance runners. *Bulletin of Institute of Health and Sports Sciences* 1986;9:195-202.
 - 54 Klausen T, Mohr T, Ghisler U, Nielsen OJ. Maximal oxygen uptake and erythropoietic responses after training at moderate altitude. *Eur J Appl Physiol* 1991;62:376-9.
 - 55 Asahina K, Ikai M, Agawas Y, Kuroda Y. A study on acclimatization to altitude in Japanese athletes. *Schweizerische Zeitschrift fuer Sportmedizin* 1966;14:240-5.
 - 56 Faulkner JA, Kollias J, Favour CB, Buskirk ER, Balke B. Maximum aerobic capacity and running performance at altitude. *J Appl Physiol* 1968;24:685-91.
 - 57 Daniels J, Oldridge N. The effects of alternate exposure to altitude and sea-level on world-class middle-distance runners. *Med Sci Sports* 1970;2:107-12.
 - 58 Balke B, Nagle FJ, Daniels J. Altitude and maximum performance in work and sports activity. *JAMA* 1965;194:176-9.
 - 59 Dill DB, Adams WC. Maximal oxygen uptake at sea-level and at 3,090-m altitude in high school champion runners. *J Appl Physiol* 1971;30:854-9.
 - 60 Reeves JT, Grover RF, Cohn JE. Regulation of ventilation during exercise at 10,200 ft in athletes born at low altitude. *J Appl Physiol* 1967;22:546-54.
 - 61 Buskirk ER, Kollias J, Akers RF, Prokop EK, Reategui EP. Maximal performance at altitude and on return from altitude in conditioned runners. *J Appl Physiol* 1967;23:259-66.
 - 62 Vallier JM, Chateau P, Guezennec CY. Effects of physical training in a hypobaric chamber on the physical performance of competitive triathletes. *Eur J Appl Physiol* 1996;73:471-8.
 - 63 Rusko HK, Kirvesniemi H, Paavola L, Vahasoyrinki P, Kyrö KP. Effect of altitude training on sea level aerobic and anaerobic power of elite athletes. *Med Sci Sports Exerc* 1996;28(S5):[abstract 739].
 - 64 Bailey DM, Davies B, Romer L, Gandy G. Physiological implications of moderate altitude training (1640 metres) on sea-level endurance performance in elite distance runners. *Brit J Sports Med* 1996;30:370[abstract].
 - 65 Telford RD, Graham KS, Sutton JR, Hahn AG, Campbell DA, Creighton SW, et al. Medium altitude training and sea-level performance. *Med Sci Sports Exerc* 1996;28(S5):[abstract 741].
 - 66 Adams WC, Bernauer EM, Dill DB, Bomar Jr JB. Effects of equivalent sea-level and altitude training on VO₂ Max and running performance. *J Appl Physiol* 1975;39:262-6.
 - 67 Roskamm H, Landry F, Samek L, Schlager M, Weidemann H, Reindell H. Effects of a standardized ergometer training program at three different altitudes. *J Appl Physiol* 1969;27:840-7.
 - 68 Levine BD, Engfred K, Friedman DB, Kjaer M, Saltin B, Clifford PS, Secher NH. High altitude endurance training: effect on aerobic capacity and work performance. *Med Sci Sports Exerc* 1990;22:[abstract 209].
 - 69 Friedman DB, Levine BD, Hanel B, Engfred K, Clifford PS, Secher NH. Endurance training and the hypoxic ventilatory response. *Med Sci Sports Exerc* 1990;22(S99):[abstract 592].
 - 70 Rahkila P, Rusko H. Effects of high altitude training on muscle enzyme activities and physical performance characteristics of cross-country skiers. In: Komi P, ed. *Exercise and sport biology*. Champaign, IL: Human Kinetics, 1982:143-51.
 - 71 Hahn AG, Telford RD, Tumilty DM, McBride ME, Campbell DP, Kovacic JC, et al. Effect of supplementary hypoxic training on physiological characteristics and ergometer performance of elite rowers. *Excel* 1992;8:127-38.
 - 72 Favier R, Spielvogel D, Desplanches D, Ferretti G, Kayser B, Grunfelder A, et al. Training in hypoxia vs. training in normoxia in high-altitude natives. *J Appl Physiol* 1995;78:2286-93.
 - 73 Davies CT, Sargeant AJ. Effects of hypoxic training on normoxic maximal aerobic power output. *Eur J Appl Physiol* 1974;33:227-36.
 - 74 Desplanches D, Hoppeler H, Linoissier MT, Denis C, Claassen H, Dormois D, et al. Effects of training in normoxia and normobaric hypoxia on human muscle ultrastructure. *Eur J Physiol* 1993;425:263-7.
 - 75 Hansen JE, Vogel JA, Stelter GP, Consolazio CF. Oxygen uptake in man during exhaustive work at sea-level and high altitude. *J Appl Physiol* 1967;23:511-22.
 - 76 Winslow RM, Monge CC. *Hypoxia, polycythemia and chronic mountain sickness*. Baltimore: John Hopkins University Press, 1987.
 - 77 Milledge JS. High altitude. In: Harries M, Williams C, Stanish W, Micheli L, ed. *Oxford textbook of sports medicine*. London: Oxford University Press, 1994:217-30.
 - 78 Kayser B. Nutrition and energetics of exercise at altitude. Theory and possible practical implications. *Sports Med* 1994;17:309-23.
 - 79 Saltin B. Aerobic and anaerobic work capacity at 2,300 metres. *Medical Thoracics* 1967;24:205-10.
 - 80 Gore CJ, Hahn AG, Watson DB, Norton KI, Campbell DP, Scroop GS, et al. VO₂ Max and arterial O₂ saturation at sea-level and 610 m. *Med Sci Sports Exerc* 1996;27(S5):[abstract 42].
 - 81 Anselme F, Caillaud C, Courret I, Prefaut C. Exercise induced hypoxemia and histamine excretion in extreme athletes. *Int J Sports Med* 1992;13:80-1.
 - 82 Lawler J, Powers SK, Thompson D. Linear relationship between VO₂ Max and VO₂ Max decrement during exposure to acute hypoxia. *J Appl Physiol* 1988;64:146-9.
 - 83 Koistinen P, Takala T, Martikkala V, Leppälousto J. Aerobic fitness influences the response of maximal oxygen uptake and lactate threshold in acute hypobaric hypoxia. *Int J Sports Med* 1995;26:78-81.
 - 84 Rowell LB, Taylor HL, Wang Y. Saturation of arterial blood with oxygen during maximal exercise. *J Appl Physiol* 1964;19:284-6.
 - 85 Dempsey JA. JB Wolfe memorial lecture. Is the lung built for exercise? *Med Sci Sports Exerc* 1986;18:143-55.
 - 86 Nummela A, Joutte P, Rusko H. Effect of living high and training low on sea level anaerobic performance in runners. *Med Sci Sports Exerc* 1996;28(S5):[abstract 740].
 - 87 Weil JV, Jamieson G, Brown DW, Grover RF. The red cell mass-arterial oxygen relationship in normal man. *J Clin Invest* 1968;47:1627-39.
 - 88 Shephard RJ, Bouhelle E, Vandewalle H, Monod H. Peak oxygen intake and hypoxia. *Int J Sports Med* 1988;9:279-83.
 - 89 Dick FW. Training at altitude in practice. *Int J Sports Med* 1992;13:S203-5.
 - 90 Berglund B. High-altitude training. Aspects of haematological adaptation. *Sports Med* 1992;14:289-303.
 - 91 Rejnafarje C, Lozano R, Valdivieso J. The polycythemia of high altitudes: iron metabolism and related aspects. *Blood* 1959;14:433-55.
 - 92 Hannon JP, Shields JL, Harris CW. Effects of altitude acclimatization on blood composition of women. *J Appl Physiol* 1969;26:540-7.
 - 93 Stray-Gundersen, Alexander C, Hochstein A, deLomos D, Levine BD. Failure of red cell volume to increase to altitude exposure in iron deficient runners. *Med Sci Sports Exerc* 1992;24(suppl):S90.
 - 94 Suslov FP. Basic principles of training at high altitude. In: *New studies in athletics, The IAAF Quarterly Magazine for NSA* 1994;2:45-9.
 - 95 Benoit H, Costes F, Castells J, Busso T, Denis C, Lacour JR, Geyssant A. Endurance training in normobaric hypoxia: ventilatory response and ability to exercise. *Int J Sports Med* 1992;13:90[abstract].
 - 96 Dill DB, Braithwaite K, Adams WC, Bernauer EM. Blood volume of middle distance runners: effect of 2300 m altitude and comparison with non-athletes. *Med Sci Sports Exerc* 1974;6:1-7.
 - 97 Reilly T. Circadian rhythms. In: Harries M, Stanish W, Micheli L, eds. *Oxford textbook of sports medicine*. London: Oxford University Press, 1994:238-54.
 - 98 Martino M, Myers K, Bishop P. Effects of 21 days training at altitude on sea-level anaerobic performance in competitive swimmers. *Med Sci Sports Exerc* 1996;27(S5):[abstract 37].
 - 99 Nummela A, Joutte P, Rusko H. Effect of living high and training low on sea level anaerobic performance in runners. *Med Sci Sports Exerc* 1996;28(S5):[abstract 740].
 - 100 Hogan MC, Welch HG. The effect of altered arterial oxygen tensions on muscle metabolism in dog skeletal muscle during fatiguing work. *Am J Physiol* 1986;251:C216-C222.
 - 101 Wolfel EE, Groves BM, Brooks GA, Butterfield GE, Mazzeo RS, Moore LG, et al. Oxygen transport during steady-state submaximal exercise in chronic hypoxia. *J Appl Physiol* 1996;70:1129-36.
 - 102 Boutellier U, Marconi C, di Prampero PE, Cerretelli P. Effects of chronic hypoxia on maximal performance. *Clinical Respiratory Physiology* 1982;18(S4):39-44.
 - 103 Shephard RJ, Pang NS. Potential impact of physical activity and sport on the immune system: a brief review. *Brit J Sports Med* 1994;28:247-55.
 - 104 Uchakin P. Immunological adaptations to altitude training [abstract]. In: *The third world congress for sport science* 1995: 209-210.
 - 105 Klokner M, Kharazmi A, Galbo H, Bygberg I, Pedersen BK. Influence of in vivo hypobaric hypoxia on function of lymphocytes, neutrophils, natural killer cells and cytokines. *J Appl Physiol* 1993;74:1100-6.
 - 106 Meehan RT, Duncan U, Neale LS, Taylor G, Muchmore H, Scott N, et al. Operation Everest II: alterations in the immune system at high altitudes. *J Clin Immunol* 1988;8:397-406.
 - 107 Parry-Billings M, Blomstrand E, McAndrew N, News-holme E. A communicational link between skeletal muscle, brain, and cells of the immune system. *Int J Sports Med* 1990;11(S2):S122-8.
 - 108 Wagenmakers AJM. Amino acid metabolism, muscular fatigue and muscle wasting. Speculations on adaptations at high altitude. *Int J Sports Med* 1992;13:S110-13.
 - 109 Simon-Schnass I. Risk of oxidative stress during exercise at high altitude. In: Sen CK, Packer L, Hanninen O, eds. *Exercise and oxygen toxicity*. Amsterdam: Elsevier, 1994;1: 191-210.
 - 110 Nagawa T, Kita H, Aoki J, Maeshima T, Shiozawa K. *Asian Medicine* 1968;11:619-33.
 - 111 Biselli R, Pecci G, Oliva C, Fattorossi A, D'Amelio R, Barsotti P. Effects of hypobaric hypoxia (HH) on polymorphonuclear granulocytes (PMNL) respiratory activity in HH-acclimatized and non-acclimatized rats [abstract]. *Int J Sports Med* 1992;13:94.