causing a crush injury, the remainder sustaining a high energy impact on striking the ground. The fact that none of our subjects wore a body protector may have contributed to the severity of the injuries. These protective garments are commercially available and are manufactured according to British Equestrian Trade Association (BETA) standards.¹ The use of helmets with retaining straps by all of our patients probably explains the absence of head injuries in the group, together with the fact that the energy of impact was concentrated at the pelvic region. Lack of riding experience was not a contributory factor in the group studied here, in contrast with the findings of Silver and Parry² who found that one third of serious spinal injuries caused by falls from horses were associated with inadequate rider experience. The predominance of male riders in this population may be explained by the fact that our catchment area includes Kildare county which is an internationally renowned centre for horse breeding and training in which men are very involved at amateur and professional levels. A study from Newmarket in England similarly had a much greater ratio of men to women (almost seven to one) for horse related injuries.⁴ Doctors should inform their patients who ride horses about the hazards of equestrian sports and should emphasise the im-

portance of using both a helmet and a body protector, as these may reduce the severity of the injury caused by a fall from a horse.

- 1 Parry JL. Injuries in equestrian sports. In: Renstrom PA, ed. *The encyclopaedia of sports medicine: clinical practice of sports injury prevention and care*. Oxford: Blackwell Scientific Publications, 1994:662.
- 2 Silver JR, Parry JM. Hazards of horse riding as a popular sport. *Br J Sports Med* 1991;25:105-10.
- 3 Christey GL, Nelson DE, Rivara FP, Smith SM, Condie C. Horse back riding injuries among children and young adults. *J Fam Pract* 1994;39:148-52.
- 4 Edixhoven P, Sinha SC, Dandy DJ. Horse injuries. *Injury* 1981;12:279-82.
- 5 Anonymous. From the Centers for Disease Control. Injuries associated with horse-back riding, United States 1987-88. *JAMA* 1990;264:18-19.
- 6 Hobbs GD, Yealy DM, Rivas J. Equestrian injuries: a five-year review. *J Emerg Med* 1994;12:143-5.
- 7 Nelson DE, Bixby-Hammet D. Equestrian injuries in children and young adults. *Am J Dis Child* 1992;146:611-14.
- 8 Bixby-Hammett D. Paediatric equestrian injuries. *Paediatrics* 1992;89:1173-6.
- 9 Pounder D. The grave yawns for the horseman. *Med J Aust* 1984;141:632-5.
- 10 Brueton RN. A review of 40 acetabular fractures: the importance of early surgery. *Injury* 1993;24:171-4.
- 11 Johnson EE, Matta JM, Mast JW, Letournel E. Delayed reconstruction of acetabular fractures 21-120 days following injury. *Clin Orthop* 1994;305:20-30.
- 12 Robertson C, Redmond A. The management of major trauma. In: *Oxford handbooks in emergency medicine*. 2nd ed. Oxford: Oxford University Press, 1994:102-5.
- 13 Burgess AR. Fractures of the pelvis. In: Rockwood CA, Green DP, Bucholz RW, eds. *Fractures in adults*. Philadelphia: JB Lippincott Company, 1991:1414.
- 14 Tile M. Fractures of the acetabulum. In: Rockwood CA, Green DP, Bucholz RW, eds. *Fractures in adults*. Philadelphia: JB Lippincott Company, 1991:1445.

Commentary

It will have been the experience of many doctors providing care for the injured rider that pelvic fractures are a cause for serious concern, and that a high index of suspicion must exist if a lethal outcome is to be avoided. This excellent paper draws attention to the severity of injury if a rider is crushed by a horse (weighing approximately 500 kg) or, when thrown, is exposed to a high energy impact, particularly on contact with hard ground. The spectre of a closed massive haemorrhage (up to 5 litres) without immediate local signs is a daunting one. The inclusion of a table showing examples and the mechanism of such injuries is most helpful; it may well increase our vigilance when observing such accidents in the future. It is a matter for concern that, despite the quality of care of these skilled orthopaedic surgeons at the Meath Hospital, three riders could not ride again and three were unable to return to their previous occupation. The nine equestrians are described as all wearing hats with a retention strap and that they escaped head injuries. It is encouraging to learn of their good sense, perhaps reflecting the wise tuition and many years of riding experience. There is no doubt that head injuries represent the greatest threat to horse riders, and it cannot be too strongly emphasised that suitably harnessed personal protective headgear of the highest standard must be worn at all times when mounted. A new British Standard BS EN 1384:96 came into effect on 15th January 1997 and is the English language version of EN 1384:1996 published by the European Committee for

Standardisation (CEN). It supersedes BS 4472:1988 and BS 6473:1984 which are withdrawn. In the United Kingdom this is now the minimum standard recognised by the British Horse Society and the Pony Club with the PAS 015 (Product Approval Specification) helmet recommended as of a higher standard. Professional jockeys wear helmets that are of this latter standard. It must be emphasised that if a hat is subjected to an impact it should be replaced, as external damage is inadequate in detecting significant impairment of future protection. Although it is conjectured that body protectors might have ameliorated these severe injuries, there is no clear evidence that this would have been the case. However, it is considered that such body protectors (class I, II or III) may well prevent soft tissue damage and perhaps minor fractures and injuries to the renal, hepatic, and splenic territories. The choice of body protectors should be influenced by the type of activity, comfort, weight, and freedom from restriction when riding. The one example of a rider trapped by the stirrup draws attention to the need for, in particular, recreational riders to have suitably heeled boots with matching stirrups; trainers should never be worn for riding. It should also be added that anticipation, fitness to ride, and learning how to fall might make all the difference. There is much to learn from this important paper, and it merits the close attention of all those engaged in or caring for those engaged in high risk sports.

JOHN LLOYD PARRY