Original articles

Evidence based prevention of acute injuries during physical exercise in a WHO safe community

T Timpka, K Lindqvist

Abstract**Objective**—To evaluate a community based programme for evidence based prevention of injuries during physical exercise.**Design**—Quasi-experimental evaluation using an intervention population and a non-random control population.**Participants**—Study municipality (population 41 000) and control municipality (population 26 000) in Sweden.**Main outcome measures**—Morbidity rate for sports related injuries treated in the health care system; severity classification according to the abbreviated injury scale (AIS).**Results**—The total morbidity rate for sports related injuries in the study area decreased by 14% from 21 to 18 injuries per 1000 population years (odds ratio 0.87; 95% confidence interval (CI) 0.79 to 0.96). No tendency towards a decrease was observed in people over 40. The rate of moderately severe injury (AIS 2) decreased to almost half (odds ratio 0.58; 95% CI 0.50 to 0.68), whereas the rate of minor injuries (AIS 1) increased (odds ratio 1.22; 95% CI 1.06 to 1.40). The risk of severe injuries (AIS 3–6) remained constant. The rate of total sports injury in the control area did not change (odds ratio 0.93; 95% CI 0.81 to 1.07), and the trends in the study and control areas were not statistically significantly different.**Conclusion**—An evidence based prevention programme based on local safety rules and educational programmes can reduce the burden of injuries related to physical exercise in a community. Future studies need to look at adjusting the programme to benefit all age groups.

(Br J Sports Med 2001;35:20–27)

Keywords: injuries; prevention; evaluation; community; intervention; safety promotion

Only a few decades ago, sport and physical exercise were usually omitted from the public health agenda.¹ Today, as a result of the spread of a sedentary lifestyle, the situation has changed. Morbidity from cardiovascular disease has been confirmed to be inversely related to the level of leisure time physical activity.^{2,3} Recent studies have suggested that the inverse relation between exercise and disease also holds for other organ systems,⁴ and that this

relation persists after adjustment for familial factors.⁵ An example of causal mechanisms is the positive effect of exercise on bone density at the hip in postmenopausal women.⁶ There is plenty of evidence from a public health perspective to encourage physical activity among the young and to promote maintenance of exercise with increasing age. However, in parallel with the benefits, there is the risk of injury.⁷ Community based studies in Scandinavia have suggested that every sixth unintentional injury is associated with leisure time physical exercise.^{8,9} Factors known to influence the risk of injury are the nature of the exercise (vigorousness, rules), the equipment and the environment (clothes and footwear, facilities, weather conditions), and individual characteristics (age, experience, technique).¹⁰

Comprehensive programmes for prevention of injuries related to physical exercise have been called for.¹¹ One strategy for injury prevention is to collect and analyse local injury data and then supply the results to local organisations with an interest in decreasing injury rates in the community. The first community development programme aimed at preventing injuries was initiated in 1978 in the municipality of Falköping, Sweden.¹² The aim of the present study is to evaluate the effect of using such a model, the WHO safe community programme, on sports related injuries. Sport is defined as a leisure time physical activity requiring vigorous bodily exertion and carried out according to a traditional form or set of rules. Using a quasi-experimental design, the specific aim is to analyse changes in community level morbidity rate, taking into account injury severity and anatomical location. Only injuries requiring medical attention are included in the evaluation.

The evaluated WHO safe community programme is implemented in Motala municipality in the western part of Östergötland County in Sweden. An assessment of the general structure and process of the programme has previously been reported.¹³ In the community analysis stage, performed in 1983/1984, the local epidemiology of injuries during physical exercise was studied, and the local social structure and values were analysed.^{9,14,15} An interorganisational sports safety council was formed with representatives from the municipality's leisure time administration, the Educational Association of the Swedish Sporting Organisations (SISU), the

Linköping University,
Faculty of Health
Sciences, Department
of Health and
Environment, Division
of Preventive and
Social Medicine,
S-58185 Linköping,
Sweden
T Timpka
K Lindqvist

Correspondence to:
Dr Timpka, Division of
Preventive and Social
Medicine, Department of
Health and Environment,
Faculty of Health Sciences,
Linköping University, S-581
85 Linköping, Sweden
tooti@ida.liu.se

Accepted 14 November 2000

Intercompany Sports Association, the health care administration, schools, and the local alliance of soccer clubs. Injuries sustained during team sports were found to predominate, soccer accounting for 39% of the injuries, followed by handball/basketball (10%). About 60% of the soccer injuries were consequences of aggressive play—that is, tackles, kicks, and blows. It was also found that intercompany team competitions took place with few practice sessions before matches, no time allotted for warming up, and few coaches and trained instructors. Stage two, the programme design and initiation (1985–1987) stage, included organising management of the interventions and setting local planning goals. Injuries incurred during physical exercise, more than injuries in other environments, were seen to be related to a person's conscious choice of leisure time activity. As a person's decision to use leisure time for physical exercise was seen as essential for community health, the interventions had to restrict minimally the choice of physical exercise as a leisure time activity. The implementation stage (1987–1988) included population based interventions at three levels.

- Community level: all physical education teachers in the study area participated in an injury prevention course organised by SISU; high risk groups were investigated.
- Sports organisation level: SISU was given the primary responsibility for coordinating interventions. In team sports, the emphasis was put on safety norms and rules. On the basis of experience from ice hockey,¹⁶ workshops for coaches and referees were used to define a fair play programme to discourage foul play. In addition, the administrators of the Intercompany Sports Association introduced compulsory use of shin pads during intercompany soccer matches as well as warming up programmes before kick off. For individual sports, a specific attempt was made to increase supervision of novices. For instance, a rule was introduced prescribing supervision of young riders during all interaction with horses.
- Group level: the alliance of soccer clubs arranged courses for local coaches on how injuries can be avoided by proper physical preparation.¹⁷

Individual interventions were mediated through the introduction of a sports medicine service in the primary care system. During the intervention year, 120 patients were examined by a specially trained doctor or physiotherapist. Based on the findings, individual advice was given on rehabilitation and how to avoid injury in the future. At stage four, programme maintenance-consolidation (1989–1995), the intervention activities were further integrated into existing community networks. The final stage, evaluation (1995–1999), focused on assessing and reporting results from the programme.

Methods

A quasi-experimental design¹⁸ was used, with measurements taken before and after implementation in the programme implementation area

and in a neighbouring control municipality in Östergötland county. The main intervention effect was studied using prospective registration of all acute health care episodes during the study period. As the study areas were not randomly chosen, environmental indicators were studied. To avoid bias, the research team was composed of evaluators who had not taken part in the implementation and programme managers with experience of the programme process.

DEFINITIONS

The ICD-8 was used to classify the nature and external cause of the injury.¹⁹

A sport related injury was defined as an injury requiring treatment in the health care system sustained during a leisure time physical activity that required more or less vigorous bodily exertion and carried out according to a traditional form or set of rules. Sport related injuries occurring at school were included. The abbreviated injury scale (AIS-80)²⁰ was used to classify the severity of the injury into three levels: AIS 1, minor; AIS 2, moderate; AIS 3–6, severe-fatal.²¹

DATA COLLECTION

The study period before implementation covered the 52 weeks from 1 October 1983 to 30 September 1984. The period after implementation covered the 52 weeks from 1 January 1989 to 31 December 1989.

Data were collected for all patients presenting at a health care unit located in the study and control areas during the registration periods. A report form with the time of contact and standard personal data was filled in by staff at the care unit. On the same form was recorded whether unintentional injury was a possible reason for the visit.

Two specially trained nurses made an injury classification from the medical records after the care episode. Injury severity was only recorded in the study area. When necessary, the attending doctor was asked to help to classify the injury. The registration model was based on experiences from earlier prevention programmes in Sweden.²²

The routine for data collection was tested in a pilot project. Before the start of each study period, the staff at the relevant health care units were carefully instructed, and the routine was practised over two weeks.

VALIDITY AND RELIABILITY TESTS

To identify confounding social trends, population variables possibly related to the sports injury rate were selected with the guidance of social change theory.^{23–24} Data on population age and sex mix, sites of residency, education, income, and employment were collected retrospectively from national registers (Statistics Sweden) for the study periods. Whereas physical exercise is often performed as a scheduled activity among adults, children practice their favourite sports both in improvised play and by participation in organised training and competition. In the light of previous experience,²⁵ no registration of risk exposure among participants was performed. Instead, membership

and activity data from local sports clubs in the study area were followed.

The study area had four health care centres and a county annex hospital with a casualty department, whereas the control area shared the annex hospital and had two health care centres, one of which had an emergency unit. The use of acute health care services at these units was followed by recording data from all attendances during both registration periods. Both the study and control areas are situated 50 km from Linköping University Hospital. Systematic differences between the areas with regard to injured people attending for care outside the registration areas were sought by analysing all attendances at the emergency departments of the university hospital during September 1984. Fatal injuries other than those recorded in the study were researched by analysing local police records.

To estimate the quality of the specific injury registration procedure, secondary sampling of all acute health care attendances in the study area was undertaken during the third week of the registration period before implementation and in both the study and control areas during the third week of the registration period after implementation.

STATISTICAL ANALYSIS

Statistical analyses were based on the cumulative rate of sports injuries in the community. Changes in the relative risk of injury associated with the factor "exposure to the safe community programme" were estimated by odds ratio.²⁶

Results

ENVIRONMENTAL INDICATORS

Age and sex mix in both areas were close to the national average. Residential and income characteristics also remained stable (table 1). The educational level in both areas was slightly below the national average and showed a tendency to increase. The proportion of the population in the study area who were members of a sports club was 39% in both 1983/1984 and 1989, whereas their activity level increased by 12%, from 730 000 participations in organised sport in 1983/1984 to 820 000 participations in 1989.

QUALITY OF REGISTRATIONS

During the registration period before implementation, which also included injuries other than sports injuries, identity data were missing for 18 of the 4926 injured patients (0.4%) in the study area and 23 (0.9%) of the 2694 injured patients in the control area. During the

period after implementation, which also includes non-sport related injuries, 10 of the 4287 injured patients (0.2%) in the study area could not be identified in the medical records. For six of the 2746 injured patients (0.2%) in the control area, identity data were missing.

In the registration control during the period before implementation, five (5%) of the 102 secondarily observed injuries were found not to have been registered in the study area: three had mistakenly not been registered as injuries and two others were found not to have been recorded. During the period after implementation in the study area, four (5%) of 84 secondarily observed injuries had not been registered; these had not been recorded. In the control area, seven (14%) of 51 secondarily observed injuries had not been registered: three had mistakenly not been registered as injuries and four others had not been recorded.

A lower proportion of all injured residents from the study area (11/422; 3%) than from the control area (28/253; 12%) was found to have been directly provided with acute care at the university hospital during the month of the control study.

INJURY RATES

In the study area, the total rate of sports related injuries decreased by 14% (table 2), corresponding to a change from 21 to 18 injuries per 1000 population years. No tendency to decrease was observed in people over 40. In the control area, there was no evident change in total morbidity, although a tendency towards a decrease was observed (from 16 to 15 injuries per 1000 population years). The trends in the study and control areas were not statistically significantly different. Taking injury severity into account, the risk of moderately severe injury in the study area was reduced to almost half (odds ratio 0.58; 95% confidence interval (CI) 0.50 to 0.68), while the minor injuries increased (odds ratio 1.22; 95% CI 1.06 to 1.40). At the same time, the risk of severe or fatal injuries remained constant.

TYPE OF SPORT

In the study area, total rates decreased only for injuries sustained in the two dominant team sports (table 3). In both of these, soccer (odds ratio 0.72; 95% CI 0.46 to 1.14) and handball/basketball (odds ratio 0.61; 95% CI 0.36 to 1.06), injuries to the upper extremities showed the strongest tendency to decrease. For moderate injuries, the rates were reduced by more than one third for the dominant team sports, soccer, handball/basketball, and ice hockey, and for horse riding and skiing among the individual sports. For all sports, the rates for severe or fatal injuries remained constant.

ANATOMICAL LOCATION

In the study area, the total rate decreased for injuries to the upper extremities and the trunk (table 4). The decrease was most pronounced for upper extremity injuries sustained during horse riding, which were reduced to one quarter (odds ratio 0.21; 95% CI 0.06 to 0.74). For

Table 1 Sociodemographic trends in the study area (Motala) and control area (Mjölby) over the registration periods

	Population characteristics (%)			
	Study area		Control area	
	1984 (n=41 400)	1989 (n=41 700)	1984 (n=25 900)	1989 (n=26 100)
Urban residents	82	82	79	81
Gainfully employed	49	50	49	51
Average income*	93	93	93	93

*Percentage of national mean.

Table 3 Relative risk between 1983/1984 and 1989 estimated by odds ratio (95% confidence interval) for sport related injuries in the study area (Motala) over the registration periods classified according to type of physical exercise and degree of severity

	Degree of severity					
	Minor (AIS 1)		Moderate (AIS 2)		Serious (AIS 3-6)	
	1983/1984	1989	1983/1984	1989	1983/1984	1989
Basketball/Volleyball/Handball						
Injured	49	36	32	15	0	1
Non-injured	41 383	41 674	41 400	41 695		
Odds ratio	0.73 (0.47-1.12)		0.47 (0.25-0.86)			
Racket sports						
Injured	17	23	13	10	0	1
Non-injured	41 415	41 687	41 419	41 700		
Odds ratio	1.34 (0.72-2.52)		0.76 (0.33-1.74)			
Motor sports						
Injured	6	17	21	11	1	2
Non-injured	41 426	41 693	41 411	41 699		
Odds ratio	2.82 (1.11-7.14)		0.52 (0.25-1.08)			
Soccer						
Injured	109	136	208	124	6	7
Non-injured	41 323	41 574	41 224	41 586	41 426	41 703
Odds ratio	1.24 (0.96-1.60)		0.59 (0.47-0.74)		1.16 (0.39-3.45)	
Ice hockey						
Injured	12	11	22	9	0	2
Non-injured	41 420	41 699	41 410	41 701		
Odds ratio	0.91 (0.40-2.06)		0.41 (0.19-0.88)			
Bandy						
Injured	28	42	15	15	0	2
Non-injured	41 404	41 668	41 417	41 695		
Odds ratio	1.49 (0.92-2.40)		0.99 (0.49-2.03)			
Gymnastics						
Injured	27	34	23	19	0	1
Non-injured	41 405	41 676	41 409	41 691		
Odds ratio	1.25 (0.75-2.07)		0.82 (0.45-1.51)			
Horse riding						
Injured	11	26	36	11	3	0
Non-injured	41 421	41 684	41 396	41 699		
Odds ratio	2.35 (1.16-4.75)		0.30 (0.15-0.60)			
Running						
Injured	22	17	25	16	0	0
Non-injured	41 410	41 693	41 407	41 694		
Odds ratio	0.77 (0.41-1.45)		0.64 (0.34-1.19)			
Skiing						
Injured	19	13	29	8	4	2
Non-injured	41 413	41 697	41 403	41 702	41 428	41 708
Odds ratio	0.68 (0.34-1.38)		0.27 (0.13-0.60)		0.50 (0.09-2.71)	
Skating						
Injured	13	17	13	8	0	1
Non-injured	41 419	41 693	41 419	41 702		
Odds ratio	1.30 (0.63-2.67)		0.61 (0.25-1.47)			
Others						
Injured	50	72	40	36	0	2
Non-injured	41 382	41 638	41 392	41 674		
Odds ratio	1.43 (1.00-2.05)		0.89 (0.57-1.40)			
All						
Injured	363	444	477	282	14	21
Non-injured	41 069	41 366	40 955	41 428	41 418	41 689
Odds ratio	1.22 (1.06-1.40)		0.58 (0.50-0.68)		1.49 (0.76-2.93)	

AIS, Abbreviated injury scale.

Table 4 Relative risk between 1983/1984 and 1989 estimated by odds ratio (95% confidence interval) for sport related injuries in the study area (Motala) over the registration periods classified according to diagnosis and location of injury and degree of severity

Location or cause	Degree of severity					
	Minor (AIS I)		Moderate (AIS 2)		Serious (AIS 3-6)	
	1983/1984	1989	1983/1984	1989	1983/1984	1989
Diagnosis*	1983/1984	1989	1983/1984	1989	1983/1984	1989
I						
Head						
Injured	40	34	25	13	5	2
Non-injured	41 392	41 676	41 407	41 697	41 427	41 708
Odds ratio	0.84 (0.53-1.33)		0.52 (0.26-1.01)		0.40 (0.08-2.05)	
II						
Upper extremities						
Injured	66	57	121	54	1	3
Non-injured	41 366	41 653	41 311	41 656		
Odds ratio	0.86 (0.60-1.22)		0.44 (0.32-0.61)			
III						
Spine, chest, and pelvis						
Injured	22	10	27	17	0	0
Non-injured	41 410	41 700	41 405	41 693		
Odds ratio	0.45 (0.21-0.95)		0.63 (0.34-1.15)			
IV						
Lower extremities						
Injured	115	143	206	177	7	16
Non-injured	41 317	41 567	41 226	41 533	41 425	41 694
Odds ratio	1.24 (0.97-1.58)		0.85 (0.70-1.04)		2.27 (0.93-5.52)	
V						
Lacerations and injuries at multiple sites; crush injuries						
Injured	117	198	97	20	1	0
Non-injured	41 315	41 512	41 335	41 690		
Odds ratio	1.68 (1.34-2.12)		0.20 (0.13-0.33)			
VI						
Other						
Injured	3	2	1	1	0	0
Non-injured	41 429	41 708				
Odds ratio	0.66 (0.11-3.96)					
All						
Injured	363	444	477	282	14	21
Non-injured	41 069	41 266	40 955	41 428	41 418	41 689
Odds ratio	1.22 (1.06-1.40)		0.58 (0.50-0.68)		1.49 (0.76-2.93)	

*Specification of diagnosis (International statistical classification of diseases, injuries, and causes of death, 1965 revision): I, 800-804, 830, 850-854, 870-874, 879, 910, 930-933, 935; II, 810-819, 831-834, 840-842, 880-887, 912-915; III, 805-809, 846-848, 860-869, 875-878, 911, 934, 936-939; IV, 820-829, 835-839, 843-845, 890-897, 916-917; V, 900-907, 918, 920-929; VI, 940-999.

AIS, Abbreviated injury scale.

indications that the elimination of high risk situations by adhering to rules and norms for conduct in organised sports was successful. Rules that have been successful for preventing injuries have previously been reported for specific sports—for example, ice hockey—where compulsory use of the face mask by young players has virtually eliminated eye injuries.³³ The greatest risk reduction observed in this study was for injuries to the upper extremities. A sports related injury to the upper extremities is typically caused by a collision or fall, events that are disadvantageous in most sports. Risk reduction can therefore be a consequence of the influence of the fair play programme on these specific, and for the nature of the physical exercise, unfavourable events. This observation is in accordance with previous studies,³⁴ which have suggested that rules discouraging play that may result in collisions or falls can be recommended in programmes to prevent sports related injury without deviating from the spirit of the sport. Avoidance of high risk situations by enforcing rules does not have to be restricted to team sports. An example from an individual sport is the supervision rule for young and novice horse riders, which was associated with a decrease in relative risk of riding injuries to the upper extremities to one quarter.

The decrease in moderate soccer injuries was a major contribution to the total decline in injury rate. This decrease is probably a result of several combined factors. At a population level, the risk of sustaining a soccer injury is highest among children and teenagers.¹⁴ Even though injuries to the lower extremities are the most common in adult soccer,³⁵ in this study the injuries to the upper extremities decreased the most. One explanation is that the nature of soccer injuries differs between youths and adults. Compared with adults, children more seldom sustain knee and groin injuries, whereas injuries to the upper extremities are less common in adults. The decrease in injuries to the upper extremity can be explained by the fact that the rules for fair play in school and youth soccer have had a rapid effect. Although the physical preparation and warm up programmes reduced the injuries to the lower extremities in adult soccer, the effect was slower and spread beyond the time frame of this study. To be able to optimally adjust prevention of soccer injuries to different age groups, further longitudinal community level investigations are still required.

The lack of proper definitions is a hindrance to the study of sports related injuries. The fact that this study primarily had a public health, and not a sport competitor, focus is shown in the choice of scale for classification of injury severity. The AIS is validated with regard to clinical parameters and not measurements of injury severity usually used for sports—that

is, the period of inability to perform physical exercise.³⁶ The scale is nevertheless well established for emergency room use and was, considering the study objectives, the obvious alternative for severity classification. There are few reasons to doubt that the AIS has sufficient validity for classification of acute sports injuries. However, an under-representation of overuse injuries is likely to have occurred.

The sociodemographic mix in the study and control areas are representative of general European populations, and the baseline sports related injury patterns correspond to previous studies.^{37,38} However, it has to be questioned how much of the decrease in morbidity can be attributed to the programme and to other factors. The study design was quasi-experimental, therefore external factors may have systematically influenced the results. Moreover, the individual specific risk exposure was not included in the analyses. As community level injury risk was the object of this study and participation in organised sports was controlled at the population level, the morbidity rate was considered to suffice. No evidence of change in exercise patterns or relevant environmental factors was observed during the study period. However, development of protective equipment for soccer players, for example, increased rapidly during the study period. The tendency toward a decrease in injury rates in the control area can thus be attributed to a general trend in prevention of sports related injury during the time of the study. However, both the total decrease and the decrease in non-minor injuries in the study area was of an extent that cannot be explained by general trends or chance.

In conclusion, a prevention programme based on evidence based promotion of fair play, supervision of novices, and compulsory use of protective equipment was found to reduce the rate and severity of injuries related to physical exercise in a local community. The effect was most pronounced for moderate injuries (AIS 2) and in the prevalent team sports. No effect was observed in the older age groups and on severe injuries (AIS 3–6). Community based programmes for injury prevention during physical exercise can be effective but need to be adjusted to include all age groups and types of physical exercise.

This study was supported by grants from the Swedish Centre for Sports Research (CIF), the Swedish National Institute of Public Health, the Swedish MTO programme, and the Östergötland County Council.

- 1 Terris M. Approaches to an epidemiology of health. *Am J Public Health* 1975;65:1037–45.
- 2 O'Connor G, Hennekens CH, Willett WC, et al. Physical exercise and reduced risk of nonfatal myocardial infarction. *Am J Epidemiol* 1995;142:1147–56.
- 3 Wannamethee SG, Shaper AG, Walker M. Changes in physical activity, mortality, and incidence of coronary heart disease in older men. *Lancet* 1998;351:1603–8.
- 4 Lissner L, Bengtsson C, Björkelund C, et al. Physical activity levels and changes in longevity. A prospective study of Swedish women. *Am J Epidemiol* 1996;143:54–62.
- 5 Kujala UM, Kaprio J, Sarna S, et al. Relationship of leisure-time physical activity and mortality. The Finnish twin cohort. *JAMA* 1998;279:440–4.

- 6 Kelley GA. Aerobic exercise and bone density at the hip in postmenopausal women. *Prev Med* 1998;27:798–807.
- 7 Walther SD, Sutton JR, McIntosh JM, et al. The aetiology of sports injuries. *Sports Med* 1985;2:47–58.
- 8 De Loës M, Goldie J. Incidence rate of injuries during sport activity and physical exercise in a rural Swedish municipality: incidence rates in 17 sports. *Int J Sports Med* 1988;9:461–7.
- 9 Lindqvist K. Epidemiology of accidents in a Swedish municipality. *Accid Anal Prev* 1989;21:33–43.
- 10 Peterson L, Renström O. *Injuries in sport*. Gothenburg: Tiden Publishing Co, 1983.
- 11 Gerberich S. Sports injuries: implications for prevention. *Public Health Rep* 1985;100:570–1.
- 12 Schelp L. *Epidemiology as a basis for evaluation of a community intervention program on accidents*. Thesis. Karolinska Institute, Stockholm, Sweden, 1987.
- 13 Lindqvist K, Timpka T, Schelp L. Ten years of experiences from a participatory community-based injury prevention program in Motala, Sweden. *Public Health* 1996;110:339–46.
- 14 Lindqvist K, Timpka T, Bjurulf P. Injuries during leisure physical activity in a Swedish municipality. *Scand J Soc Med* 1996;24:282–92.
- 15 Lindqvist K, Brodin H. One-year economic consequences of accidents in a Swedish municipality. *Accid Anal Prev* 1996;28:209–19.
- 16 Vaz E. *The professionalization of young hockey players*. Lincoln, NE: University of Nebraska Press, 1982.
- 17 Ekstrand J, Gillquist J. Soccer injuries and their mechanisms: a prospective study. *Med Sci Sports Exerc* 1983;15:267–70.
- 18 Cook TD, Campbell DT. *Quasi-experimentation*. Boston: Houghton Mifflin, 1979.
- 19 Swedish National Board of Health and Welfare. *Classification of diseases 1968. Systematic list (International statistical classification of diseases, injuries, and causes of death, 1965 revision, adapted for indexing hospital records and morbidity statistics)*. (In Swedish.) Stockholm: Swedish National Board of Health and Welfare, 1982.
- 20 Petrucelli E, States JD, Hames LN. The Abbreviated Injury Scale: evolution, usage, and future adaptability. *Accid Anal Prev* 1981;13:29–35.
- 21 Baker SP, O'Neill B, Haddon W, et al. The Injury Severity Score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 1974;14:187–96.
- 22 Schelp L, Svanström L. A model for registration and mapping of accident cases in health care. *Scand J Prim Health Care* 1987;5:91–9.
- 23 Carley M. *Social measurement and social indicators: issues of policy and theory*. London: George, Allen & Unwin, 1981.
- 24 Rothman J. *Planning and organizing for social change: action principles from social research*. New York: Columbia University Press, 1974.
- 25 Turtz A, Crost M. Sport-related injuries in children. *Am J Sports Med* 1986;4:294–9.
- 26 Miettinen OS. Estimability and estimation in case-referent studies. *Am J Epidemiol* 1976;103:226–35.
- 27 Armitage P, Berry G. *Statistical methods in medical research*. Oxford: Blackwell Science Ltd, 1994.
- 28 Fletcher GF, Balady G, Blair SN, et al. Statement on exercise: benefits and recommendations for physical activity programs for all Americans. *Circulation* 1996;94:857–92.
- 29 NIH Consensus Conference. Physical activity and cardiovascular health. NIH consensus development panel on physical activity and cardiovascular health. *JAMA* 1996;276:241–6.
- 30 van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med* 1992;14:82–99.
- 31 Jorgensen U. Epidemiology of injuries in typical Scandinavian team sports. *Br J Sports Med* 1984;2:59–63.
- 32 Avery JG, Harper P, Ackroyd S. Do we pay dearly for our sport and leisure activities? *Public Health* 1990;104:417–23.
- 33 Biasca N, Simmen HP, Trentz O. Review of typical ice hockey injuries. *Unfallchirurg* 1995;98:283–5.
- 34 Roberts WO, Brust JD, Leonard B, et al. Fair-play rules and injury reduction in ice hockey. *Arch Pediatr Adolesc Med* 1996;150:140–5.
- 35 Ekstrand J. Soccer injuries and their prevention. Linköping University medical dissertations no 130. Linköping, Sweden: Linköping University, 1982.
- 36 Watson AWS. Sports injuries: incidence, causes, prevention. *Physical Therapy Reviews* 1997;2:135–51.
- 37 De Loës M. Risk exposure and incidence rates of acute injuries from sports and physical exercise in a total population. Thesis. Department of Social Medicine, Karolinska Institute, and the Department of Orthopedic Surgery, Karolinska Hospital, Sundbyberg and Stockholm, Sweden, 1989.
- 38 Sahlin Y. Sports accidents in childhood. *Br J Sports Med* 1990;24:40–4.

Take home message

In these days when physical exercise is promoted in public health campaigns, it is of concern that the effectiveness of broad sports injury prevention is lacking. The WHO safety community programme reduced the total incidence of sports related injuries by 14% in a local community, and the non-minor injury rate was reduced to almost half. Evidence based programmes can prevent severe sports related injuries in a community and should be used in parallel with promotion of physical exercise.

British Association of Sport and Exercise Medicine in association with the National Sports Medicine Institute

Education programme 2001

Injury management and medicine: head, neck and upper limb Lilleshall Hall National Sports Centre, Shropshire	11–16 February
Diploma preparation course Exact date and location to be confirmed	March
General sports medicine Lilleshall Hall National Sports Centre, Shropshire	22–27 April
Current concepts Topic, date, cost, and location to be confirmed	1 May
Injury management and medicine: thorax, groin, pelvis and hip Lilleshall Hall National Sports Centre	8–13 July
General sports medicine Lilleshall Hall National Sports Centre	23–28 September
Practical sport and medicine meeting Club La Santa, Lanzarote Families and non-delegates welcome; booking deadline 31 July	4–11 October
Diploma preparation course Exact date and location to be confirmed	October
BASEM Congress Date, cost, and location to be confirmed	Oct/Nov
Injury Management and Medicine: lower limb Lilleshall Hall National Sports Centre	18–23 November
Current Concepts Topic, date, and location to be confirmed	2 December

For further details of these courses please contact Mr Barry Hill, The National Sports Medicine Institute, c/o Medical College of St Bartholomew's Hospital, Charterhouse Square, London EC1M 6BQ.

Tel: 020 7251 0583 (ext 237); fax: 020 7251 0774; email: barry.hill@nsmi.org.uk

Web site: www.nsmi.org.uk