EDITOR'S
CHOICE

Airway dysfunction in swimmers

Valérie Bougault,^{1,2} Louis-Philippe Boulet²

¹Faculté des Sciences du Sport, Droit et Santé de Lille 2, Lille, France

²Centre de recherche, Institut universitaire de cardiologie et de pneumologie de Québec, Québec, Canada

Correspondence to

Valerie Bougault, Faculté des Sciences du Sport, Université Droit et Santé de Lille 2, Lille 59790, France; valerie.bougault@univ-lille2.fr

Received 29 November 2011

Accepted 11 December 2011

Published Online First

12 January 2012

ABSTRACT

Elite competitive swimmers are particularly affected by airway disorders that are probably related to regular and intense training sessions in a chlorinated environment. Upper and lower airway respiratory symptoms, rhinitis, airway hyper-responsiveness, and exercise-induced bronchoconstriction are highly prevalent in these athletes, but their influence on athletic performance is still unclear. The authors reviewed the main upper and lower respiratory ailments observed in competitive swimmers who train in indoor swimming pools, their pathophysiology, clinical significance and possible effects on performance. Issues regarding the screening of these disorders, their management and preventive measures are addressed.

INTRODUCTION

Regular physical exercise is recognised as an important component to attain optimal health and swimming is among the preferred activities in developed countries. Buoyancy properties of water and the warm humid atmosphere of indoor swimming pools are particularly attractive for infants, the older generation or people with various ailments. Although there are recognised benefits on health, there is evidence that people who regularly attend indoor chlorinated swimming pools are at risk of developing airway disorders, associated with repeated inhalation of chlorine byproducts.^{1–4}

Chlorine, an inexpensive and easy-to-use product, is widely used in swimming pools worldwide as a chemical disinfectant, serving as the principal barrier to microbial contaminants.⁵ During chlorination process, chlorine reacts with organic matter brought by swimmers to form disinfection byproducts such as chloramines. The concentrations of these chlorine byproducts are regularly controlled in the pools' water and regulated by the WHO.⁵ However, gases such as nitrogen trichloride (NCl₃) are also formed during the reaction. NCl₃ is responsible for the strong smell reported by swimmers in chlorinated swimming pools and is strongly irritating for the airways.⁵ According to guidelines from the WHO, the maximum cut-off for NCl₃ in pools' atmosphere is 0.5 mg/m³,⁵ but its levels are not routinely measured. Other more expensive disinfectants may be used to replace chlorine, but data are still limited on their effects on the respiratory tract.

As the ventilation rate during swimming determines the quantity of inhaled chlorine byproducts, elite swimmers, training many hours in the pools, are likely to be more affected by these products. Exposure to these products may also account for the high prevalence of upper and lower airway disorders reported in this group.⁶ In keeping with this possibility, is the recent finding of significant

airway inflammation and remodelling on bronchial biopsies of swimmers similar to what is found in those with mild asthma.⁷

In this report, we discuss the prevalence of upper and lower airway disorders in swimmers, underlying mechanisms of development and persistence, their general management and the future research needed to help understand their clinical significance in order to prevent potential long-term damage to the airways.

ADVERSE RESPIRATORY HEALTH EFFECTS OF SWIMMING IN CHLORINATED SWIMMING POOLS

Airway symptoms are often reported by children, lifeguards, pool workers and swimmers, who are regularly exposed to a chlorinated swimming pool environment.^{1,4,8,9}

Upper airways

Prevalence and diagnosis

The most disturbing symptoms among swimmers seem to originate from the upper airways. Nasal obstruction, rhinorrhoea, sneezing and nasal itching are reported by up to 74% of competitive elite swimmers, during the training season.^{9–11} The diagnosis of nasal disorders is often made through a physical examination and medical history. In the case of recurrent obstruction or rhinorrhoea, nasal cytology, endoscopy and/or rhinomanometry may provide additional information on the nature of the disease.¹¹

Mechanisms of rhinitis

Exercise itself may cause rhinitis in athletes independently of atopic status.¹² There may be a decrease in nasal resistance after exercise,¹³ following the stimulation of the autonomic nervous system during physical activity. The latter may act as an active decongestant (sympathetic activation), although it may also induce exercise-induced rhinorrhoea (parasympathetic influence).^{13,14}

In a recent study on nasal cytology, 44% of young competitive swimmers with rhinitis had a pre-existing allergic component. In those with non-allergic rhinitis, 63% had a predominant nasal neutrophilic inflammation.¹¹ After the use of a nose-clip during a 30-day swimming period, nasal symptoms were significantly reduced, particularly in those with neutrophilic inflammation. This highlights the irritating role of chlorine on the nasal mucosa.¹¹ The mechanisms and consequences of such an irritating effect are still unclear, but epithelial damage may occur, as observed in the lower airways.³

Nasal symptoms may also be due to an acute viral infection or a postinfectious state. There is a high incidence of viral infections in elite swimmers

Table 1 Main recommendations to limit chlorine-byproduct exposure in bathers and swimmers attending indoor chlorinated swimming pools⁴⁵

	Measures
Swimmers' behaviour	<p>To respect bare feet zones.</p> <p>To use a swimsuit reserved exclusively for swimming.</p> <p>To wear a swim cap.</p> <p>To respect personal hygiene precautions before entering the pool.</p> <p>To take off makeup and other cosmetics.</p> <p>To take a shower with soap before entering the pool.</p> <p>Pool accessory equipment (buoys, water lines, goggles, balls and so on) must be appropriately maintained at all times and used exclusively in the pool.</p>
Premises' organisation	<p>To avoid crossing between dirty and clean areas (only one direction).</p> <p>Footbath should contain water disinfected by chlorine at a residual concentration of 5 mg/l.</p> <p>Ventilation equipment should be regularly maintained.</p> <p>Overflow tank should be easily accessible and equipped with efficient mechanically driven ventilation.</p> <p>A stripping system should be installed.</p>
Water treatment	To respect the maximal cut-off proposed by the WHO – freshwater provision should be of 30 l/bather/day.
Management of air quality	<p>Flow rate of fresh air should be of at least 60 m³/h.</p> <p>Maximum cut-off for NCl₃ in the air should be reduced to 0.3 mg/m³.</p>
Recommendations, baby swimming	<p>Infant swimmers and their supervisors should have a pool for their exclusive use.</p> <p>Babies should wear disposable diapers adapted to water.</p> <p>Before swimming session:</p> <p>Water temperature should be 32°C.</p> <p>Air temperature should be set according to water temperature.</p> <p>Water should have been double renewed before the beginning of the session.</p> <p>The maximum cut-off for NCl₃ in the water should be 0.2 mg/m³.</p> <p>Maximum turbidity cut-off should be 0.2 FNU.</p> <p>Ventilation should not be stopped or reduced the night before the session.</p> <p>When possible, a fresh air ventilation of swimming pool and hall should be done at least 1 h before the session.</p> <p>After swimming session:</p> <p>Completely renew the water.</p> <p>Disinfect the pool with chlorine at the maximal cut-off authorised during a few hours.</p>
Recommendations for workers and swimmers	To regularly measure the level of NCl ₃ in the air.

This table was developed according to the recommendations from the 'Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail'.⁴⁵
FNU, Formazine Nephelometric Unit.

during heavy training programmes.¹⁵ Intense training is known to affect, transiently, the immune system involved in tissue damage repair¹⁶ and increase the risk of developing upper respiratory tract infections due to viruses.^{16 17}

Clinical consequences and possible effects on performance

Nasal symptoms experienced by competitive swimmers during the training season impair their quality of life and may worsen their performance.⁹ Symptoms due to allergic rhino conjunctivitis, quality of life and performance may improve with appropriate medication and environmental measures.¹⁸

Pyne *et al* failed to provide significant evidence for an association between upper airway disorders and competitive performance in elite swimmers. However, the authors judiciously emphasised that the mean differences observed in performance time between healthy and affected swimmers (ie, 0.1 to 0.5 s in 100 to 200 m events) were greater than differences between winning a medal or missing a medal altogether.¹⁵

Lower airway diseases

Prevalence and diagnosis

Screening in competitive swimmers has demonstrated that up to 76% have symptomatic or asymptomatic airway hyper-responsiveness (AHR) and/or exercise-induced bronchoconstriction (EIB).^{4 6 19 20} Asthma is generally diagnosed clinically on the basis of symptoms such as wheezing, dyspnoea, chest

tightness, phlegm production and cough, associated with objective evidence of variable airway obstruction or AHR.²¹ The prevalence of asthma before the beginning of the competitive career seems to be similar or slightly increased in swimmers, compared with other endurance athletes and healthy subjects.^{4 6 22} This suggests that swimmers do not engage in swimming due to their asthma, but rather develop respiratory disorders during their athletic career. Moreover, the development of AHR generally occurs in young adults rather than in adolescent competitive swimmers.²³ Such AHR may be the result of the cumulative effects of years of intense swimming training, the repeated inhalation of large volumes of air containing chlorine byproducts possibly inducing airway inflammatory and structural changes.⁴

Symptoms

Asthma-like symptoms are commonly reported by swimmers, children, lifeguards and bathers who regularly attend chlorinated swimming pools.^{1 4 8 10 19} Cough is the most common exercise-related symptom and a study reported that 18% of swimmers had to cease a training session at least once due to the severity of the symptoms associated with the strong odour of the pool environment.⁴

Pulmonary/airway function changes

Many competitive swimmers demonstrate AHR to direct or indirect stimuli,^{4 10 19 20 24} independently of the presence or not

of swimming-related symptoms.⁴²⁰ In accordance with the criteria used by the International Olympic Committee-Medical Commission's (IOC-MC) Independent Asthma Panel,²⁵ about 60% of young competitive swimmers had AHR to at least one bronchial provocation test.^{20 26} Little information is currently available on the comparison between synchronised swimmers, free-water swimmers, swimmers, divers and water-polo players. However, the very high training level encountered in Olympic swimmers and synchronised swimmers, training up to 40 h per week, may explain the high prevalence of AHR in these two swimming specialities.^{27 28} In swimmers, AHR is usually found in the presence of a normal resting flow-volume curve, emphasising the necessity to use a bronchial provocation or a reversibility test to assess airway function.⁴²⁰ Furthermore, in some athletes, provocation tests with direct stimuli such as methacholine may be negative while EIB can be confirmed by indirect tests such as eucapnic voluntary hyperpnoea (EVH) in these athletes.^{4 26}

Mechanisms

As for endurance athletes, increased ventilation sustained for many hours per day during many weeks of swimming training can probably affect the airway epithelium.²⁹ Epithelial damage, in addition to the osmotic stress, may lead to the release of inflammatory mediators and possibly initiate a repair and rehydration process involving fibrogenic cytokines.²⁹ Since an indoor swimming pool environment is warm and humid, dehydration of the airways should be less than for other sports, thus protecting against EIB and AHR. However, further evidence suggests that in chlorinated indoor swimming pools, repeated inhalation of chloramines formed during the reaction of chlorine with organic matter brought in the pool by swimmers can contribute over the years to allergic diseases and asthma in swimmers.^{3 4} Elite swimmers, breathing large volumes of air containing chlorine derivatives, immediately above the water surface, are particularly at risk. In young children, the possibility that asthma develops following repeated attendance of chlorinated swimming pools remains controversial^{30 31} and needs further study to better assess the role of chlorine in the development of airway disorders.

Epithelial damage and chlorine hypothesis

Indoor chlorinated swimming pool attendance induces a rapid and transient disruption of the airway epithelium in recreational swimmers, children or passively exposed attendees, without concomitant development of asthma symptoms.³² Inhaled chlorine byproducts certainly have the potential to cause structural changes in the airway epithelium, thus permitting allergens easier access to antigen-presenting cells, possibly promoting allergen sensitisation.³³ Allergies, AHR, asthma and other airway disorders may then possibly develop as a consequence of repeated exposure to these byproducts.³³

The mechanisms responsible for epithelial damage and AHR in swimmers remain incompletely understood. The increased oxidative stress observed in swimmers' airways and reduced antioxidant capacity may promote the release of inflammatory mediators and sensitisation of airway smooth muscle, contributing to the development of AHR in swimmers.³⁴

Inflammation and remodelling of the airways

Similar inflammation and remodelling changes have been observed in the bronchial mucosa and/or sputum of competitive swimmers, when compared with nonathletes with mild asthma.^{3 7} Bronchial remodelling was observed in swimmers independently of AHR to methacholine or EVH.⁷ These

changes could presumably constitute a normal adaptation process to intense training and be partly reversible after the end of the swimming career, although this remains to be confirmed.³⁶ Despite the fact that clinical consequences of these changes remain uncertain, it seems advisable to propose preventive measures such as reduction of chloramines' exposure in pool environments. We do not know, however, if preventive treatments should be offered to swimmers, in order to try to reduce airway inflammation and the possibility of structural changes.

Non-invasive measures of airway inflammation, such as with exhaled nitric oxide (eNO) or induced sputum analysis, may be useful to detect active airway disease,³⁷ but these tests may not be sensitive enough to detect a more subtle inflammation of the airway mucosa. Neutrophil cell counts in induced sputum of competitive swimmers have been shown to correlate with the number of training hours per week.³⁵ However, no significant difference in induced sputum inflammatory cell counts was observed between competitive swimmers and controls, when performed 12 h or more after the last training session.³⁵ Furthermore, no change in eNO was observed after swimming training in competitive swimmers.³⁸

Clinical consequences and possible effects on performance

When treated, Olympic swimmers with asthma perform their sport as well or better than other athletes without asthma.²⁸ The consequences of EIB on performance seem obvious, but for asymptomatic AHR, they remain unclear. In a certain proportion of the general population, asymptomatic AHR is associated with the subsequent development of asthma, particularly in subjects with atopy.³⁹ In swimmers, AHR has been shown to be transient.^{36 40} The clinical consequences of AHR in this group are also unclear and need to be further studied.

What do the swimmers report to the physician?

It has been suggested that competitive swimmers, particularly the youngest, may consider their upper and lower airway symptoms as a 'normal phenomenon' due to the intensity of exercise or the presence of a strong chemical odour in the swimming pool rather than to rhinitis or asthma.⁴ Consequently, many of them do not mention about symptoms to their general physician, and do not get the opportunity to consult a respiratory physician.

Should swimmers be screened for respiratory disorders?

Whether we should screen competitive swimmers for rhinitis, allergy, and AHR or asthma is still debated, mainly due to the costs involved. Currently, questionnaires are available on allergies and respiratory symptoms of athletes, and may constitute an inexpensive prescreening assessment.⁴¹ Spirometry and a bronchodilator reversibility test are also easy to perform and may be ordered on a yearly basis in high-level swimmers, despite a high baseline forced expiratory volume in one second. In athletes reporting symptoms during swimming, the recommendations of the IOC-MC Independent Asthma Panel should be followed and a test should be performed to prove AHR or EIB, according to the described methodologies.²⁵

GENERAL MANAGEMENT AND PREVENTION

Management

Currently, rhinitis and asthma in swimmers are managed as in the general population.^{42 43} No data, however, are available on the benefits of treatment with nasal anticholinergic agents or corticosteroids for rhinitis in swimmers, a disorder that is

mainly characterised by a neutrophilic rather than eosinophilic inflammation. The management of asthma is mostly based upon the use of fast-acting bronchodilators for intermittent symptoms or for the prevention of EIB, in addition to inhaled corticosteroids (ICS). Inhaled long-acting β_2 -agonists or leucotrienes receptor antagonists may be added if asthma control is not achieved by a low dose of ICS.⁴³

Prevention

Considering its simplicity to use, efficiency and low cost, chlorine will probably continue to be used in the future. Reducing exposure to inhaled chlorine byproducts, however, could potentially help reduce the risk of airway disorder in swimmers. A summary of recommendations to enhance the quality of water and air in the indoor swimming pools is presented in table 1.

Should we replace chlorine by other disinfectants?

Alternative chemical disinfectants to chlorine such as ozone and chlorine dioxide are increasingly being used.⁵ Some studies reported that the use of a copper-silver disinfectant does not induce epithelial damage in swimmers or favour the development of asthma in school children.^{32–44} On the other hand, copper-silver disinfection cannot remove organic material and does not seem to ensure the total elimination of viral pathogens from water, even if its use is combined with low levels of chlorine.⁴⁵ Each alternative disinfection measure produces its own set of byproducts and/or is partially effective as a means of disinfection.⁵ Therefore, there is a need to better understand the chemistry, toxicology and epidemiology of chemical disinfectants and their associated byproducts of disinfection. However, whatever the product uses, it should not compromise the microbiological quality of swimming pools' water.

A need to change bathers' behaviour

Because NCl_3 levels in the air above swimming pool water is influenced by ambient ventilation and water chemistry, a proper ventilation of the pool environment and reduction of human protein matter in the water may reduce respiratory health problems of swimmers.⁵ Efficient ventilation in indoor swimming pools would increase the turnover of ambient air and remove concentrated chloramines.

An efficient measure to reduce chloramines production would also be a modification of swimmer's behaviour (table 1). In addition, adhering to a maximal occupancy rate in the swimming pools, expressed as the number of bathers per unit of volume, would help to keep cleaner air and water of swimming pools.⁴⁶ Altogether, these measures could significantly reduce the accumulation of chloramines in the swimming pools' ambient air and contribute to a better respiratory health of the swimmers and bathers.

CONCLUSION

Particular attention should be paid to the respiratory health of competitive swimmers, its management and prevention. Simple measures could be taken to reduce airway dysfunction in this population with minimal financial costs. Further studies are needed to understand the combined effects of exercise and inhalation of chlorine byproducts and the direct effect of chloramines as potential inducers of epithelial damage and airway inflammation. The impact of such disorders on athletic performance and their effects on the health of swimmers during and after the end of their training career should be evaluated.

What this study adds

This review underlines the impact of swimming in a chlorinated environment on respiratory health of people regularly attending swimming pools. This review may help clinicians and trainers to better understand the mechanisms of airway disorders in swimmers and to become aware of the necessity for a proper management.

Contributors Both authors contributed equally to the redaction of this manuscript and the literature search. VB is the guarantor.

Competing interests None.

Provenance and peer review Commissioned; internally peer reviewed.

REFERENCES

1. **Jacobs JH**, Spaan S, van Rooy GB, *et al*. Exposure to trichloramine and respiratory symptoms in indoor swimming pool workers. *Eur Respir J* 2007;**29**:690–8.
2. **Momas I**, Brette F, Spinasse A, *et al*. Health effects of attending a public swimming pool: follow up of a cohort of pupils in Paris. *J Epidemiol Community Health* 1993;**47**:464–8.
3. **Bernard A**, Carbone S, Michel O, *et al*. Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools. *Occup Environ Med* 2003;**60**:385–94.
4. **Potts JE**. Adverse respiratory health effects of competitive swimming: the prevalence of symptoms, illness, and bronchial hyper-responsiveness to methacholine and exercise (dissertation). Vancouver (BC): University of British Columbia, 1994.
5. World Health Organization. Guidelines for safe recreational water environments. Vol 2. Swimming pools and similar environments. Geneva: WHO, 2006.
6. **Bougault V**, Turmel J, Levesque B, *et al*. The respiratory health of swimmers. *Sports Med* 2009;**39**:295–312.
7. **Bougault V**, Loubaki L, Joubert P, *et al*. Airway remodeling and inflammation in competitive swimmers with and without airway hyperresponsiveness. *J Allergy Clin Immunol* 2012;**129**:351–8, 358.e1.
8. **Lévesque B**, Duchesne JF, Gingras S, *et al*. The determinants of prevalence of health complaints among young competitive swimmers. *Int Arch Occup Environ Health* 2006;**80**:32–9.
9. **Bougault V**, Turmel J, Boulet LP. Effect of intense swimming training on rhinitis in high-level competitive swimmers. *Clin Exp Allergy* 2010;**40**:1238–46.
10. **Zwick H**, Popp W, Budik G, *et al*. Increased sensitization to aeroallergens in competitive swimmers. *Lung* 1990;**168**:111–15.
11. **Gelardi M**, Ventura MT, Fiorella R, *et al*. Allergic and non-allergic rhinitis in swimmers: clinical and cytological aspects. *Br J Sports Med* 2012;**46**:54–8.
12. **Silvers WS**, Poole JA. Exercise-induced rhinitis: a common disorder that adversely affects allergic and non-allergic athletes. *Ann Allergy Asthma Immunol* 2006;**96**:334–40.
13. **Forsyth RD**, Cole P, Shephard RJ. Exercise and nasal patency. *J Appl Physiol* 1983;**55**:860–5.
14. **Harris WE**, Giebaly K, Adair C, *et al*. The parasympathetic system in exercise-induced rhinorrhoea. *Rhinology* 1992;**30**:21–3.
15. **Pyne DB**, McDonald WA, Gleeson M, *et al*. Mucosal immunity, respiratory illness, and competitive performance in elite swimmers. *Med Sci Sports Exerc* 2000;**33**:348–53.
16. **Gleeson M**. Immune function in sport and exercise. *J Appl Physiol* 2007;**103**:693–9.
17. **Spence L**, Brown WJ, Pyne DB, *et al*. Incidence, etiology, and symptomatology of upper respiratory illness in elite athletes. *Med Sci Sports Exerc* 2007;**39**:577–86.
18. **Katellaris CH**, Carrozzi FM, Burke TV, *et al*. Effects of intranasal budesonide on symptoms, quality of life, and performance in elite athletes with allergic rhinoconjunctivitis. *Clin J Sport Med* 2002;**12**:296–300.
19. **Helenius IJ**, Ryttilä P, Metso T, *et al*. Respiratory symptoms, bronchial responsiveness, and cellular characteristics of induced sputum in elite swimmers. *Allergy* 1998;**53**:346–52.
20. **Bougault V**, Turmel J, Boulet LP. Bronchial challenges and respiratory symptoms in elite swimmers and winter sport athletes: Airway hyper-responsiveness in asthma: its measurement and clinical significance. *Chest* 2010;**138**:31–7S.
21. **Bousquet J**, Jeffery PK, Busse WW, *et al*. Asthma. From bronchoconstriction to airways inflammation and remodeling. *Am J Respir Crit Care Med* 2000;**161**:1720–45.
22. **Langdeau JB**, Turcotte H, Thibault G, *et al*. Comparative prevalence of asthma in different groups of athletes: a survey. *Can Respir J* 2004;**11**:402–6.

23. **Pedersen L**, Lund TK, Barnes PJ, *et al.* Airway responsiveness and inflammation in adolescent elite swimmers. *J Allergy Clin Immunol* 2008;**122**:322–7.
24. **Pedersen L**, Winther S, Backer V, *et al.* Airway responses to eucapnic hyperpnea, exercise, and methacholine in elite swimmers. *Med Sci Sports Exerc* 2008;**40**:1567–72.
25. **Fitch KD**, Sue-Chu M, Anderson SD, *et al.* Asthma and the elite athlete: summary of the International Olympic Committee's consensus conference, Lausanne, Switzerland, January 22–24, 2008. *J Allergy Clin Immunol* 2008;**122**:254–60.
26. **Castricum A**, Holzer K, Brukner P, *et al.* The role of the bronchial provocation challenge tests in the diagnosis of exercise-induced bronchoconstriction in elite swimmers. *Br J Sports Med* 2010;**44**:736–40.
27. **Bougault V**, Turmel J, Boulet LP. Prevalence of airway hyper-responsiveness in international level synchronised swimmers (Abstract). Oslo, Norway: XIth International Symposium for Biomechanics and Medicine in Swimming, 2010. http://www.nih.no/upload/BMS2010/Documents/BM2010_Program_Abstracts_final_lowres.pdf (accessed 24 Nov 2011).
28. **Fitch K**. Overview of the respiratory health of Olympic athletes (Abstract). Monaco, France: IOC World Conference on Prevention of Injury and Illness in Sport, 2011. http://video.nih.no/Congress2011/07c_13.45_Fitch_1/Ken%20Fitch.swf (accessed 1 Nov 2011).
29. **Anderson SD**, Kippelen P. Airway injury as a mechanism for exercise-induced bronchoconstriction in elite athletes. *J Allergy Clin Immunol* 2008;**122**:767–73.
30. **Font-Ribera L**, Villanueva CM, Nieuwenhuijsen MJ, *et al.* Swimming pool attendance, asthma, allergies, and lung function in the Avon Longitudinal Study of Parents and Children cohort. *Am J Respir Crit Care Med* 2011;**183**:582–8.
31. **Schoefer Y**, Zutavern A, Brockow I, *et al.* Health risks of early swimming pool attendance. *Int J Hyg Environ Health* 2008;**211**:367–73.
32. **Carbounelle S**, Francaux M, Doyle I, *et al.* Changes in serum pneumoproteins caused by short-term exposures to nitrogen trichloride in indoor chlorinated swimming pools. *Biomarkers* 2002;**7**:464–78.
33. **Bernard A**. Chlorination products: emerging links with allergic diseases. *Curr Med Chem* 2007;**14**:1771–82.
34. **Bougault V**, Morissette M, Murray N, *et al.* Oxidative stress in swimmers' airways (Abstract). *Eur Respir J* 2010;**36**(Suppl 54):5582.
35. **Bougault V**, Turmel J, St-Laurent J, *et al.* Asthma, airway inflammation and epithelial damage in swimmers and cold-air athletes. *Eur Respir J* 2009;**33**:740–6.
36. **Helenius I**, Ryttilä P, Sarna S, *et al.* Effect of continuing or finishing high-level sports on airway inflammation, bronchial hyper-responsiveness, and asthma: a 5-year prospective follow-up study of 42 highly trained swimmers. *J Allergy Clin Immunol* 2002;**109**:962–8.
37. **Lemière C**, Ernst P, Olivenstein R, *et al.* Airway inflammation assessed by invasive and non-invasive means in severe asthma: eosinophilic and noneosinophilic phenotypes. *J Allergy Clin Immunol* 2006;**118**:1033–9.
38. **Moreira A**, Delgado L, Haahtela T, *et al.* Training does not affect exhaled nitric oxide in competitive swimmers. *Allergy* 2008;**63**:623–4.
39. **Boulet LP**. Asymptomatic airway hyper-responsiveness: a curiosity or an opportunity to prevent asthma? *Am J Respir Crit Care Med* 2003;**167**:371–8.
40. **Bougault V**, Turmel J, Boulet LP. Airway hyperresponsiveness in elite swimmers: is it a transient phenomenon? *J Allergy Clin Immunol* 2011;**127**:892–8.
41. **Bonini M**, Braidò F, Baiardini I, *et al.* AQUA: Allergy Questionnaire for Athletes. Development and validation. *Med Sci Sports Exerc* 2009;**41**:1034–41.
42. **Bonini S**, Bonini M, Bousquet J, *et al.* Rhinitis and asthma in athletes: an ARIA document in collaboration with GA2LEN. *Allergy* 2006;**61**:681–92.
43. **Bateman ED**, Hurd SS, Barnes PJ, *et al.* Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J* 2008;**31**:143–78.
44. **Bernard A**, Nickmilder M, Voisin C, *et al.* Impact of chlorinated swimming pool attendance on the respiratory health of adolescents. *Pediatrics* 2009;**124**:1110–18.
45. **Bosch A**, Diez JM, Abad FX. Disinfection of human enteric viruses in water by copper-silver and reduced levels of chlorine. *Water Sci Technol* 1993;**27**:351–6.
46. Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail. Risques sanitaires liés aux piscines: évaluation des risques sanitaires liés aux piscines – partie 1: piscines réglementées (saisine n 2006/11). Ed Scientifique Eau Et Agents Biologiques, 2010. www.anses.fr (accessed 5 Oct 2011).