OPEN ACCESS

Classification and grading of muscle injuries: a narrative review

Bruce Hamilton,¹ Xavier Valle,^{2,3,4} Gil Rodas,³ Luis Til,² Ricard Pruna Grive,² Josep Antoni Gutierrez Rincon,² Johannes L Tol⁵

► To read the full version of this paper, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2014-093551>)

¹Medical Lead HPSNZ, National Training Centre, Millenium Institute of Sport and Health, Auckland, New Zealand

²F.C. Barcelona Medical Services, Barcelona, Spain

³Sports Medicine School, University of Barcelona, Barcelona, Spain

⁴Clinica Mapfre de Medicina del Tenis, Barcelona, Spain

⁵Sports Medicine Department, Aspetar, Qatar Orthopaedic and Sports Medicine Hospital, Doha, Qatar

Correspondence to

Dr Bruce Hamilton, High Performance Sport NZ, Millenium Institute of Sport and Health, Mairangi Bay, Auckland 0632, New Zealand; Bruce.hamilton@hpsnz.org.nz

Accepted 16 October 2014

ABSTRACT

A limitation to the accurate study of muscle injuries and their management has been the lack of a uniform approach to the categorisation and grading of muscle injuries. The goal of this narrative review was to provide a framework from which to understand the historical progression of the classification and grading of muscle injuries. We reviewed the classification and grading of muscle injuries in the literature to critically illustrate the strengths, weaknesses, contradictions or controversies. A retrospective, citation-based methodology was applied to search for English language literature which evaluated or utilised a novel muscle classification or grading system. While there is an abundance of literature classifying and grading muscle injuries, it is predominantly expert opinion, and there remains little evidence relating any of the clinical or radiological features to an established pathology or clinical outcome. While the categorical grading of injury severity may have been a reasonable solution to a clinical challenge identified in the middle of the 20th century, it is time to recognise the complexity of the injury, cease trying to oversimplify it and to develop appropriately powered research projects to answer important questions.

INTRODUCTION

Muscle injuries are among the most common injuries in sport, but there remain few high-quality studies evaluating their specific management.¹ A limitation to the comprehensive study of muscle injuries has been the lack of uniformity in their categorisation and description. Reflecting this observation, the Munich muscle injury classification group stated that ‘...little information is available in the international literature about muscle injury definitions and classification systems.’² The terms classify and grade do not refer to the same process. Injury ‘classification’ refers specifically to describing or categorising an injury (eg, by its location, mechanism or underlying pathology). By contrast, a ‘grade’ provides an indication of injury severity.³

While it would appear logical to initially classify a muscle injury according to a system of choice (eg, by location or mechanism), and then grade the injury severity within that classification (eg, grade I, II or III), this approach has not been uniformly applied. When referring to muscle injuries, the terms classification and grading have frequently been used interchangeably and ambiguously.^{4,5}

The following narrative review outlines the historical progression of the classification and grading literature for acute muscle injuries, predominantly from the English literature. We illustrate the strengths, weakness, inconsistencies and controversies in the

literature to better understand the paradigm in which muscle injury descriptors have been developed, thereby facilitating future understanding.

Methodology

A retrospective, citation-based methodology was applied to search for English language literature which evaluated or utilised a novel muscle classification or grading system. Peer-reviewed journal publications were the primary source, but prior to 1970 popular sports medicine textbook sources were also utilised. No systematic search strategy was used and one author (BH) independently screened and documented the literature.

Muscle injury classifications

By the turn of the 20th century, muscle injuries were being classified by both the causative or mechanistic forces and the anatomical location of the injury (see online supplementary table S1 for a complete summary).^{6–8} Specifically, authors categorised muscle injuries as either being derived from internal forces (secondary to violent exertion) or external forces (secondary to direct ‘violence’).^{6–8} Anatomically, it was recognised that the muscle may ‘rupture’ in distinct locations such as ‘where fibres meet the tendon’, the ‘body of the muscle’ or in the tendon.⁶ This early literature predates frequently cited classification systems, but most likely provided the foundation for their subsequent development^{9–12} as minor variations of this approach were common throughout the early 20th century.^{13–17}

In the 1960s, approaches to muscle injury classification expanded to include newly defined conditions such as myositis ossificans, and to incorporate mechanistic and anatomical descriptors in a single classification.^{9,12} This approach of incorporating the mechanism, injury location and distinct pathologies continues to be utilised.¹⁸ Indeed, the classification of muscle injuries by the causal mechanism (intrinsic vs extrinsic forces) and the anatomical location of the injury has remained largely unchanged with time.^{10,19,20} Although not all influential authors in the past have felt it clinically necessary to separately classify internally and externally derived injuries,¹¹ animal injury research and imaging techniques of the late 20th century have largely validated the clinically derived distinctions of ‘contusion’ (external force) and ‘strain’ (intrinsic force).

From the 1980s, availability of imaging in the form of ultrasound (US) and MRI allowed direct visualisation of muscle injury, resulting in enhanced anatomical accuracy and an expansion of the

Open access
Scan to access more
free content

CrossMark

To cite: Hamilton B, Valle X, Rodas G, et al. *Br J Sports Med* 2015;**49**:306.

imaging literature.^{10 20–22} Initial image-based muscle injury publications were descriptive in nature, articulating the appearance of images corresponding to popular clinical classification (and grading) approaches that were in use by the 1980s.²³ Early imaging reports included only small subject numbers, limited clinical information,^{10 20–24} and rarely acknowledged that the imaging descriptions of the clinical classifications had little pathophysiological or prognostic grounding.²⁵

In the early 21st century, there has been renewed interest in muscle classification. Askling *et al*^{26–28} have continued the history of classifying hamstring injuries by their distinct anatomical location (eg proximal/distal; semimembranosus/biceps femoris; figure 1A–C) and, for the first time, demonstrated a relationship between the anatomical location and time to return to play. The same group also proposed a subclassification of intrinsic force injury, specifically the ‘stretching type’ (type 2) or ‘high speed running type’ (type 1) related to the muscle involved.^{26–28} While stretch versus high force intrinsic injuries have previously been alluded to,⁹ Askling *et al*^{26–28} highlighted a relationship between the specific nature of the intrinsic force and clinical prognosis. Thus, classification of injuries based on their causation may have prognostic validity. Ultimately, larger study numbers may enable further grading of injury severity within each classification (ie, Stretch (type 2) or high speed (type 1)).

‘Central tendon’ disruption was identified as a potential contributor to prognosis in earlier imaging studies.²⁰ Comin *et al*²⁹ recently demonstrated a distinct prognosis when the ‘central’ tendon was disrupted in hamstring injuries. In 1966, Bass³⁰ proposed that anatomical classification was clinically relevant, but only recently have Askling *et al*³¹ and Comin *et al*²⁹ provided evidence that identifying the specific tissue involved may have clinical utility.

Historically, there was limited critique of the literature pertaining to muscle injury classification, but in the past 10 years, authors have critically evaluated the limitations of muscle injury classification.

Muscle injury grading

The ‘clinical era’ (c1900–1980)

The severity of any given injury may be determined by either directly identifying the underlying pathology (eg, with a muscle biopsy), indirectly by utilising a proxy representation of the pathology such as imaging or systemic (eg, serum) markers, or by the serial documentation of observed clinical outcomes related to specifically identified clinical, pathological, imaging or systemic variables.

Excluding a single biopsy report of a clinical ‘grade III’ injury in a patient suffering from systemic sclerosis, we are not aware of any human biopsy studies assessing muscle strain severity.³² While a number of animal models have assessed muscle injury, few reflect either the mechanism of injury or the symptoms experienced by athletes.^{33–39} As a result, clinicians employ indirect means of evaluating muscle damage severity.²⁵ Traditionally, this was achieved by identifying a range of symptoms and/or signs at injury presentation thought to reflect the severity of any underlying pathology, with early 20th century literature grading including variations of ‘mild’ or ‘severe’ categories.^{13 15 16 40 41} By the 1960s, there were a range of categorical grading systems for muscle injury with the severity determined by a subjective assessment of function loss, assumed to reflect either the number of muscle fibres involved or the injury location.^{13 15 16 40 41}

In 1966, the American Medical Association (AMA) subcommittee on the classification of sports injuries published the first comprehensive three grade system for acute muscle injuries.²³ This grading system, incorporating both clinical descriptors and a theoretical pathological correlation, provides the most detailed grading of the pre-imaging era (see online supplementary table 1 for details). While rarely cited, the AMA grading appears to have been highly influential in subsequent literature,^{42 43} and almost certainly forms the clinical basis for early imaging grading;^{10 44} recent literature appears to neglect this substantial work.^{2 45}

While more than 1500 muscle injuries were described in the literature prior to the 1980s,^{7 13 16 17 46 47} only Bass (1966), studying 72 football players, made any attempt to correlate any clinical findings to a distinct outcome.³⁰ As a result, there is no established prognostic validity to historical (clinical) grading systems of muscle injury, but despite this they have been recycled in various modified forms and continue to appear in the literature.⁴⁸ Traditional clinical grading of muscle injury is attractive for practitioners and patients, but the grading is based on expert opinion only and lacks any substantial empirical support.

The imaging era (c1985–2000)

From the 1980s, US and MRI allowed the indirect assessment of muscle anatomy and pathology with contemporary authors anticipating that this would provide greater objectivity to the complex clinical evaluation.⁴⁹ As with clinical grading systems, we are aware of no studies that validate imaging findings observed in muscle strain with a confirmed underlying pathology. Subsequently, as already noted with regard to the classification literature, early imaging grading literature describes the radiological appearance of a clinical presentation, which in itself lacks any pathophysiological or prognostic validity. Of the early imaging grading descriptions,^{10 24 44 49–52} only Pomeranz and Heidt,²⁰ evaluating 14 muscle injuries, made any attempt to establish a distinct prognosis between muscle grades. Pomeranz and Heidt²⁰ assessed muscle injury size with MRI and then carefully followed athletes during their rehabilitation, providing one of the earliest indications of a possible correlation between the extent of imaging findings and clinical prognosis.

Recently, limited data have supported the reliability,⁵³ and the prognostic validity of categorical imaging grading derived from clinical evaluations.⁵⁴ A landmark study involving 207 elite European footballers failed to show a statistically significant difference in prognosis between grade 1 and 2 injuries.⁵⁴ By contrast, a subsequent investigation with a larger cohort did establish that MRI can statistically differentiate prognosis in this group of athletes.⁵⁵ It is important to note that these studies do not reflect the natural history (ie, an injury course unmodified by treatment) of any of the clinically or radiologically determined injuries. Injured athletes in these cohorts may have been exposed to a range of intensive rehabilitation and invasive treatments which may significantly impact the natural history and prognosis for any given radiological appearance.⁵⁶ Furthermore, the timing of any imaging is critical for prognostication as MRI findings may remain after an athlete is clinically ready to return to competition.^{25 57} Finally, it is pertinent to recall that the imaging descriptions utilised (ie a ‘modification of Peetrons classification’^{10 54}) are based on historically derived clinical descriptions that have no established validity.

Hence, while data are emerging that in certain situations are image based, categorical grading systems may provide a valid prognosis, technical limitations and data reproduction demands that further evidence be collected.

Figure 1 Part (A) proximal hamstring origin. (B) Type I ('high-strain') hamstring injuries mainly occur within the long head of the biceps and typically involve the proximal muscle-tendon junction. (C) Type II (stretching) injuries typically occur close to the ischial tuberosity and affect the proximal free tendon of semimembranosus, reproduced with permission from Askling C, Schache A. *Brukner & Khan's clinical sports medicine*, Chapter 31: posterior thigh pain.⁸⁹

The modern era (c2000–)

Early in the 21st century, researchers began to address the limitations of existing classification and grading systems for muscle injuries. First, as illustrated above, there have been attempts to provide an evidential basis for correlating clinical and radiological grading with injury severity. Second, there has been recognition that imaging can provide continuous rather than categorical data, and that this may correlate with injury prognosis. Finally, there has been recognition of the benefits of combining clinical and radiological evaluations, and with this insight authors have begun to reconstruct classification and grading systems.

Establishing an evidence base

Despite Wise (1977) describing a clinical grading system for muscle injury incorporating both symptoms and signs,⁴³ until recently there was little empirical evidence for the prognostic value of either clinical variable.^{19 58} In 2003, Verrall *et al* illustrated that symptoms and signs such as the sudden onset of pain and localised tenderness, respectively, may accurately reflect underlying injury to the hamstring muscle. Furthermore, both patient reported pain severity and the clinician's 'estimate' of injury severity correlated with the return to play.⁵⁸ Similarly, time taken to walk pain free (more or less than 1 day) has been

noted to have a degree of prognostic merit for hamstring injuries,⁵⁹ and active range of knee joint extension has been correlated with hamstring injury severity and reinjury risk.^{19 60}

By contrast, Askling *et al*³¹ assessed passive straight leg raise and knee flexion strength in a cohort of injured sprinters and dancers, and found that neither of the clinical tests correlated with prognosis. Likewise, Verrall *et al*⁶¹ found that the initial clinical examination, including the categorical finding of swelling, bruising, tenderness and pain on hamstring contraction, had no value in predicting the likelihood of reinjury. Low numbers of subjects and conflicting clinical findings necessitate further data to enable a better understanding of the merits of clinical assessment in muscle injury prognostication.

The significant role of Ekstrand *et al*⁵⁴ in correlating imaging with prognosis has been noted above. An additional finding of note, that grade 0 (MRI negative injuries) had a significantly better prognosis than all other grades of injury, supports the findings of previous authors who highlighted the prognostic relevance of a positive versus negative MRI.^{25 58 62–64} Paradoxically, while US has been shown to be as sensitive as MRI in determining the presence of muscle injury,²⁵ a study involving 51 footballers illustrated no difference in hamstring injury prognosis based on a positive or negative initial US.⁶⁵ In the presence of a clinically diagnosed muscle injury, there remains uncertainty as to how to interpret negative imaging findings—specifically whether this reflects a muscle injury below the sensitivity of the imaging modality, or whether this is a true negative for muscle injury.^{2 66}

Anatomical details now visible on imaging, such as tendon involvement, may impact on muscle injury prognosis suggesting that historical categorical approaches to grading may be oversimplistic in nature.²⁹ Evidence is slowly accumulating, allowing the critical evaluation of clinical and radiological variables in the assessment and prognosis of muscle injury, but data quality and quantity remain limited.

Measuring continuous variables and prognosis

Since 2002, authors have correlated injury size on imaging, using a continuous scale, with clinical outcome (table 1).^{25 27 28 58 61–63 65 67 69 71 72} Of the continuous variables studied using MRI for hamstring injuries, lesion length, cross-sectional area and estimated volume all provide some predictive value—in essence, the larger the lesion, the longer the rehabilitation period required. By contrast, US has not consistently shown a relationship between muscle length and prognosis.^{25 65}

Askling *et al*^{28 69} found that the absolute (clinical and radiological) distance from the ischium in 18 hamstring injured sprinters correlated with prognosis. This finding was not reproduced in 15 dancers with ‘stretch’ type injuries of the hamstrings,²⁷ and previous studies have not found an association between injury location and return to play duration.⁵⁸ Furthermore, while continuously measured clinical variables such as pain at the initial injury correlate positively with return to play,^{58 63} measures of hip flexibility and knee flexion strength do not.³¹

With the total data using imaging analysis of continuous variables totalling just over 200 cases, there remain limited data with which to accurately predict an individual’s specific prognosis based on injury size. Furthermore, in the majority of the studies cited, bias cannot be excluded, as treating clinicians were not blinded to MRI or clinical findings. As a result, further study and larger subject numbers are required.

Combined approaches to classification and grading

The past 5 years have seen a range of publications touting ‘new’ muscle injury classification and grading systems, on occasion

varying little from previous approaches.^{2 16 45 73–79} However, only two manuscripts provide any clinical data to support the proposed systems.

In a novel approach, Cohen *et al*⁷⁶ evaluated hamstring injuries in 43 American football players, combining six radiological observations into a single injury score (see online supplementary table 1 for details). A combined score of greater than 10 points was found to have a worse prognosis. This comprehensive grading system, utilising currently available knowledge, illustrates a progressive approach and while the data have yet to be reproduced elsewhere, its clinical merit warrants further inspection.

In 2012, an experienced group of clinicians met in Munich to establish a comprehensive system for the classification and grading of muscle injuries.² While the authors retained the ‘direct’ and ‘indirect’ terminology first utilised as early as 1902 (then termed internal and external),⁶ that is where similarities with many previous classifications end. The authors expand previous definitions of muscle injury and pain, to incorporate terms such as ‘functional’, ‘structural’, ‘neuro-muscular muscle disorder’, ‘overexertion-related muscle disorder’ and ‘fatigue induced muscle disorder’ in an expansive system of subclassification. In support of this classification is an extensive clinical description including delineating factors from the history, examination and imaging. The authors also grade the ‘partial muscle tear’ into ‘minor partial muscle tear’ (3A) and ‘moderate partial muscle tear’ (3B), on the basis of symptoms, signs and imaging.

As with previous classification systems, there remain both a limited pathophysiological and pathoanatomical basis on which to base the detailed subclassification, and limited evidence for distinct clinical outcomes on the basis of either the classification or grading. However, the Munich group implemented an expansive research programme involving European football clubs to evaluate the validity of their system.⁸⁰ This study suggested a relationship between the injury category/grade and prognosis, particularly in differentiating the return to play duration between ‘functional’ and ‘structural’ disorders. Whether this terminological distinction reflects the previously identified importance of MRI positive versus negative injury remains to be determined.⁸⁰ However, the significance of this work, and the fact that for the first time in the history of muscle injuries, large volumes of data are being utilised to test a classification and grading system, should be recognised and commended. For the first time in over 100 years of muscle injury grading, authors are testing a proposed model.

Summary and future challenges

In reviewing the evolution of muscle injury classification and grading, several themes became apparent.

1. Variability in the nomenclature utilised to classify and grade muscle injuries has resulted in limited ability to compare the few studies available. Standardisation and enhanced anatomical detail⁸¹ of structural descriptions in manuscripts would enhance future discussion.
2. Authors have sometimes ignored, subtly adapted, or on occasion misrepresented existing muscle injury grading and classification systems, without recognising or addressing their limitations. This has resulted in widely used but unsubstantiated dogma established solely on expert opinion.
3. A historical ambivalence towards reporting clinical outcomes has meant that evidence is only beginning to appear relating clinical or imaging observations to functional outcome. To

Table 1 Clinical muscle injury research utilising continuous variables for the assessment of severity

Author	Grading/description	Outcome	Cited cases
Slavotinek <i>et al</i> ⁶³	Description based on MRI findings	Approximate volume of muscle involved; (r=0.46) percentage of abnormal muscle (r=0.70)	Association with RTP duration
Verrall <i>et al</i> ⁶⁸	Description based on MRI findings and symptoms	Subjective pain score Amount of pain	Association with MRI determined severity Positive correlation with RTP duration
Connell <i>et al</i> ²⁵	Description based on MRI and US findings	Injury cross-sectional area (%) Longitudinal length (r=0.58); Volume of intramuscular haematoma	US determined cross-sectional area associated with RTP duration; MRI positive correlation with RTP; haematoma, no correlation with RTP.
Gibbs <i>et al</i> ⁶²	Description based on MRI findings	Cross-sectional area (%); Length of lesion (cm)	Positive statistical correlation with RTP
Verrall <i>et al</i> ⁶¹	Description based on MRI findings	MRI transverse size (%); MRI volume	Larger lesion, increased risk of injury in subsequent season
Schneider-Kolsky <i>et al</i> ⁶⁷	Description based on MRI findings	Longitudinal length of lesion on coronal views (r=0.58); Cross-sectional area (%)	Positive correlation with RTP
Asklings <i>et al</i> ³¹	Description based on Clinical findings	Hip flexibility (Degrees/Borg CR-10 pain scale); Knee flexion strength (dynamometer)	No data on relationship to RTP
Koulouris <i>et al</i> ⁶⁸	Description based on MRI findings	Cross-sectional injured area (mm); Injury location (muscle, location); Injury longitudinal length (mm)	Non-significant impact on reinjury risk
Asklings <i>et al</i> ²⁸	Description based on MRI findings	Distance to ischial tuberosity (r=0.54); depth of injury (r=0.58); Volume of injury (r=0.61); Cross section of injury (r=0.70) Length of injury (r=0.51) Width of injury (r=0.39)	Positive correlation with RTP No statistical correlation with RTP No statistical correlation with RTP
Asklings <i>et al</i> ²⁷	Description based on MRI findings	Distance to ischial tuberosity; Length of injury; Width of injury; Depth of injury; Volume of injury	No statistical correlation with RTP
Balius <i>et al</i> ⁶⁹	Description based on US findings	Length of lesion	Positive significant association with RTP
Nescolarde <i>et al</i> ⁷⁰	Grading based on changes in localised BIA	Resistance; reactance (xc); phase angle PA	Decreases with increasing injury severity
Peterson <i>et al</i> ⁶⁵	Description based on US findings	Length of lesion	No association with RTP

*Refers to duplication of athletes from previous manuscript.

BIA, bioimpedance analysis; PA, phase angle; RTP, return to play; US, ultrasound.

date, there remains minimal pathological or prognostic validity to the majority of classification and grading systems utilised.

- While it may be reasonable to classify and subclassify the nature of an injury, given our current understanding of the variable healing times of different tissues, and the range of tissues involved in even a simple 'muscle' injury, it seems unlikely that any categorical grading of muscle injury severity will accurately predict an individual's healing time. While the categorical grading of injury severity may have been a reasonable solution to a clinical challenge identified in the middle of the 20th century, it is time to recognise the complexity of muscle injury, and to develop appropriately powered research projects to answer appropriate questions.

In the future, a range of novel techniques may provide further clues as to the underlying injury and prognosis, including serological biomarkers of injury,⁸² advanced MR imaging,^{83 84} diffusion tensor imaging⁸⁵⁻⁸⁷ and bio-impedance techniques.^{70 88} Given the incidence of muscle injuries, there remain limited injuries being incorporated into formal study protocols, and as a result there remains much to be done.

Understanding the history of muscle injury classification and grading provides a foundation for the development of appropriate questions.

What are the new findings?

- Classification and grading refer to distinct elements of muscle injury evaluation, but have been used interchangeably in the literature.
- Systems for clinical classification and grading have been present in the literature for over 100 years; in many ways, current approaches offer the clinician no more than did the first efforts.
- There is limited evidence to support either the pathological or prognostic validity of clinical and radiological grading systems.

Acknowledgements The authors would like to acknowledge the support of both Aspetar, Qatar Orthopaedic and Sports Medicine Hospital and FC Barcelona in the completion of this manuscript.

Contributors BH was involved in the concept design, research development and data collection, manuscript preparation and final approval. JLT was involved in the concept design, manuscript review and final approval. XV, GR, LT, RPG and JAGR were involved in the concept design and initiation, manuscript review and critique and final approval.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Mason D, Dickens V, Vail A. Rehabilitation for hamstring injuries (Review). *Cochrane Collab* 2012;2012:1–23.
- Mueller-Wohlfahrt H-W, Haensel L, Mithoefer K, et al. Terminology and classification of muscle injuries in sport: the Munich consensus statement. *Br J Sports Med* 2013;47:342–50.
- MedlinePlus: Merriam-Webster Medical Dictionary. 2013.
- Brunker P, Khan K. *Clinical sports medicine*. 3rd edn. Sydney: The McGraw-Hill Companies, 2007.
- Engelbrechtsen A, Myklebust G, Holme I, et al. Intrinsic risk factors for hamstring injuries among male soccer players; a prospective cohort study. *Am J Sports Med* 2010;38:1147–53.
- Crowley DD. Suturing of muscle and tendons. *Calif State J Med* 1902:48–54.
- Marsh H. Clinical lecture on displacements and injuries of muscles and tendons. *BMJ* 1896;2:181–6.
- Gilcreest E. Rupture of muscles and tendons. Particularly subcutaneous rupture of the biceps flexor cubiti. *JAMA* 1925;84:1819–22.
- O'Donoghue DH. *Treatment of injuries to athletes*. 1st edn. Philadelphia: W.B. Saunders Company, 1962.
- Peterson P. Ultrasound of muscles. *Eur Radiol* 2002;12:35–43.
- Ryan A. Quadriceps strain, rupture and charlie horse. *Med Sci Sports* 1969;1:106–11.
- Tucker WE, Armstrong JR. *Injury in sport: the physiology, prevention and treatment of injuries associated with sport*. 1st edn. London: Staples Press, 1964.
- Featherstone D. *Sports injuries*. 1st edn. Bristol: John Wright and Sons Ltd, 1957.
- Heald CB. *Injuries in sport: a general guide for the practitioner*. 1st edn. London: Oxford University Press, 1931.
- Page E. *Athletic injuries and their treatment*. London: Arco Publications, 1962.
- Thorndike A. *Athletic injuries: prevention, diagnosis and treatment*. 1st edn. Philadelphia: Lea and Febiger, 1938.
- Williams JGP. *Sports Medicine*. 1st edn. Baltimore: The Williams and Wilkins Company, 1962.
- Lee J, Healey J. Sonography of lower limb muscle injury. *Am J Roentgenol* 2004;182:341–51.
- Malliaropoulos N, Papacostas E, Kiritsi O, et al. Posterior thigh muscle injuries in elite track and field athletes. *Am J Sports Med* 2010;38:1813–19.
- Pomeranz SJ, Heidt RS Jr. MR imaging in the prognostication of hamstring injury. Work in progress. *Radiology* 1993;189:897–900.
- Connell D, Potter H, Sherman M, et al. Injuries of the pectoralis major muscle: evaluation with MR imaging. *Radiology* 1999;210:785–91.
- Jarvinen T, Kaariainen M, Jarvinen M, et al. Muscle strain injuries. *Curr Opin Rheumatol* 2000;12:155–61.
- Rachun A. *Standard Nomenclature of Athletic Injuries*. 1st edn. Chicago, Illinois: American Medical Association, 1966.
- Peterson P, Creteur P. Echographies et traumatismes musculaires aigus. In: Chevrot A, Kahn M, Morvan G, eds. *Imagerie Des Parties Molles De L'Appareil Locomoteur*. Sauramps Medical, 1993:229–35.
- Connell DA, Schneider-Kolsky ME, Hoving JL, et al. Longitudinal study comparing sonographic and MRI assessments of acute and healing hamstring injuries. *Am J Roentgenol* 2004;183:975–84.
- Asklings CM, Tengvar M, Saartok T, et al. Sports related hamstring strains-two cases with different etiologies and injury sites. *Scand J Med Sci Sports* 2000;10:304–7.
- Asklings CM, Tengvar M, Saartok T, et al. Acute first-time hamstring strains during slow-speed stretching: clinical, magnetic resonance imaging, and recovery characteristics. *Am J Sports Med* 2007;35:1716–24.
- Asklings CM, Tengvar M, Saartok T, et al. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med* 2007;35:197–206.
- Comin J, Malliaras P, Baquie P, et al. Return to competitive play after hamstring injuries involving disruption of the central tendon. *Am J Sports Med* 2013;41:111–15.
- Bass AL. Rehabilitation after soft tissue injury. *Br J Sports Med* 1966;162–72.
- Asklings C, Saartok T, Thorstensson A. Type of acute hamstring strain affects flexibility, strength, and time to return to pre-injury level. *Br J Sports Med* 2006;40:40–4.
- Rubin SJ, Feldman F, Staron RB, et al. Magnetic resonance imaging of muscle injury. *Clin Imaging* 1995;19:263–9.
- Garrett WE Jr. Injuries to the muscle-tendon unit. *Instr Course Lect* 1988;37:275–82.
- Garrett WE Jr. Muscle strain injuries: clinical and basic aspects. *Med Sci Sports Exerc* 1990;22:436–43.
- Garrett WE Jr. Muscle strain injuries. *Am J Sports Med* 1996;24(6 Suppl):S2–8.
- Garrett WE Jr, Cluff JC, Bassett FH, 3rd. Histochemical correlates of hamstring injuries. *Am J Sports Med* 1984;12:98–103.
- Garrett WE Jr, Seaber AV, Boswick J, et al. Recovery of skeletal muscle after laceration and repair. *J Hand Surg Am* 1984;9:683–92.
- Garrett WE Jr, Safran MR, Seaber AV, et al. Biomechanical comparison of stimulated and nonstimulated skeletal muscle pulled to failure. *Am J Sports Med* 1987;15:448–54.
- Nikolaou PK, Macdonald BL, Glisson RR, et al. Biomechanical and histological evaluation of muscle after controlled strain injury. *Am J Sports Med* 1987;15:9–14.
- Colson J. *Strapping and bandaging for football injuries*. Reprint 1960 ed. London: The Football Association, 1953.
- Smart M. *The principles of treatment of muscles and joints by graduated muscular contractions*. 1st edn. London: Oxford Medical Publications, 1933.
- Oakes BW. Hamstring muscle injuries. *Aust Fam Physician* 1984;13:587–91.
- Wise DD. Physiotherapeutic treatment of athletic injuries to the muscle-tendon complex of the leg. *Can Med Association J* 1977;117:635–9.
- Takebayashi S, Takasawa H, Banzai Y, et al. Sonographic findings in muscle strain injury: clinical and MR imaging correlation. *J Ultrasound Med* 1995;14:899–905.
- Chan O, Del Buono A, Best T, et al. Acute muscle strain injuries: a proposed new classification system. *Knee Surg Sports Traumatol Arthroscopy* 2012;20:2356–62.
- Anzel SH, Covey KW, Weiner AD, et al. Disruption of muscles and tendons. An analysis of 1014 cases. *Surgery* 1959;45:406–14.
- McMaster PE. Tendon and muscle ruptures. Clinical and experimental studies on the causes and location of subcutaneous ruptures. *JBJS* 1933:705–22.
- Maquieira J, Ghisi J, Kokalj A. Rectus abdominis muscle strains in tennis players. *Br J Sports Med* 2007;41:842–8.
- Fleckenstein JL, Shellock FG. Exertional muscle injuries: magnetic resonance imaging evaluation. *Top Magn Reson Imaging* 1991;3:50–70.
- De Smet AA. Magnetic resonance findings in skeletal muscle tears. *Skeletal Radiol* 1993;22:479–84.
- De Smet AA, Fisher DR, Heiner JP, et al. Magnetic resonance imaging of muscle tears. *Skeletal Radiol* 1990;19:283–6.
- Fleckenstein JL, Weatherall PT, Parkey RW, et al. Sports-related muscle injuries: evaluation with MR imaging. *Radiology* 1989;172:793–8.
- Hamilton B, Whiteley R, Almusa E, et al. Excellent reliability for MRI grading and prognostic parameters in acute hamstring injuries. *Br J Sports Med* 2014;48:1385–7.
- Ekstrand J, Healy JC, Walden M, et al. Hamstring muscle injuries in professional football: the correlation of MRI findings with return to play. *Br J Sports Med* 2012;46:112–17.
- Hallen A, Ekstrand J. Return to play following muscle injuries in professional footballers. *J Sports Sci* 2014;32:1229–36.
- Robinson M, Hamilton B. Medical interventions in the management of hamstring muscle injury. *Euro J Sports Sci* 2014;14:743–51.
- Reurink G, Goudswaard GJ, Tol JL, et al. MRI observations at return to play of clinically recovered hamstring injuries. *Br J Sports Med* 2014;48:1370–6.
- Verrall GM, Slavotinek JP, Barnes PG, et al. Diagnostic and prognostic value of clinical findings in 83 athletes with posterior thigh injury: comparison of clinical findings with magnetic resonance imaging documentation of hamstring muscle strain. *Am J Sports Med* 2003;31:969–73.
- Warren P, Gabbe BJ, Schneider-Kolsky M, et al. Clinical predictors of time to return to competition and of recurrence following hamstring strain in elite Australian footballers. *Br J Sports Med* 2010;44:415–19.
- Malliaropoulos N, Isinkaye T, Tsitsas K, et al. Reinjury after acute posterior thigh muscle injuries in elite track and field athletes. *Am J Sports Med* 2011;39:304–10.
- Verrall GM, Slavotinek JP, Barnes P, et al. Assessment of physical examination and magnetic resonance imaging findings of hamstring injury as predictors for recurrent injury. *J Orthop Sports Phys Ther* 2006;36:215–24.
- Gibbs NJ, Cross TM, Cameron M, et al. The accuracy of MRI in predicting recovery and recurrence of acute grade one hamstring muscle strains within the same season in Australian Rules football players. *J Sci Med Sport* 2004;7:248–58.
- Slavotinek JP, Verrall GM, Fon GT. Hamstring injury in athletes: using MR imaging measurements to compare extent of muscle injury with amount of time lost from competition. *Am J Radiol* 2002;179:1621–8.
- Verrall GM, Slavotinek JP, Barnes PG, et al. Clinical risk factors for hamstring muscle strain injury: a prospective study with correlation of injury by magnetic resonance imaging. *Br J Sports Med* 2001;35:435–9; discussion 440.

- 65 Peterson J, Thorborg K, Nielsen M, *et al.* The diagnostic and prognostic value of ultrasonography in soccer players with acute hamstring injuries. *Am J Sports Med* 2014;42:399–404.
- 66 Tol JL, Hamilton B, Best TM. Palpating muscles, massaging the evidence? An editorial relating to 'Terminology and classification of muscle injuries in sport: the Munich consensus statement'. *Br J Sports Med* 2013;47:340–1.
- 67 Schneider-Kolsky ME, Hoving JL, Warren P, *et al.* A comparison between clinical assessment and magnetic resonance imaging of acute hamstring injuries. *Am J Sports Med* 2006;34:1008–15.
- 68 Koulouris G, Connell D, Brukner PD, *et al.* Magnetic resonance imaging parameters for assessing risk of recurrent hamstring injuries in elite athletes. *Am J Sports Med* 2007;35:1500–6.
- 69 Balias R, Maestro A, Pedret C, *et al.* Central aponeurosis tears of the rectus femoris: practical sonographic prognosis. *Br J Sports Med* 2009;43:818–24.
- 70 Nescolarde L, Yanguas J, Lukaski H, *et al.* Localized bioimpedance to assess muscle injury. *Physiol Meas* 2013;34:237–45.
- 71 Koulouris G, Connell D. Imaging of hamstring injuries: therapeutic implications. *Eur Radiol* 2006;16:1478–87.
- 72 Askling CM, Malliaropoulos N, Karlsson J. High-speed running type or stretching-type of hamstring injuries makes a difference to treatment and prognosis. *Br J Sports Med* 2012;46:86–7.
- 73 Dixon J. Gastrocnemius vs. soleus strain: how to differentiate and deal with calf muscle injuries. *Curr Rev Musculoskelet Med* 2009;2:74–7.
- 74 Hancock C, Sanders T, Zlatkin M, *et al.* Flexor femoris muscle complex: grading systems used to describe the complete spectrum of injury. *Clin Imaging* 2009;33:130–5.
- 75 Lee J, Mitchell A, Healy J. Imaging of muscle injury in the elite athlete. *Br J Radiology* 2012;85:1173–85.
- 76 Cohen SB, Towers JD, Zoga A, *et al.* Hamstring injuries in professional football players: magnetic resonance imaging correlation with return to play. *Sports Health* 2011;3:423–30.
- 77 Rodas G, Pruna R, Til L, *et al.* Clinical practice guide for muscular injuries. Epidemiology, diagnosis, treatment and prevention. *Apunts Med Esport* 2009;64:179–203.
- 78 Renstrom P. Muscle injuries in sports. In: Ljungqvist A, Peltokallio P, Tikkanen H, eds. *Sports medicine in track and field athletics*. Kouvola, Finland: IAAF, 1985:17–28.
- 79 Pollock N, James S, Lee J, *et al.* British athletics muscle injury classification: a new grading system. *Br J Sports Med* 2014;48:1347–51.
- 80 Ekstrand J, Askling C, Magnusson H, *et al.* Return to play after thigh muscle injury in elite football players: implementation and validation of the Munich muscle injury classification. *Br J Sports Med* 2013;47:769–74.
- 81 Woodley SJ, Mercer SR. Hamstring muscles: architecture and innervation. *Cells Tissues Organs* 2005;179:125–41.
- 82 Guerrero M, Guiu-Comadevall M, Cadefau JA, *et al.* Fast and slow myosins as markers of muscle injury. *Br J Sports Med* 2008;42:581–4; discussion 584.
- 83 Cagnie B, Elliott J, O'Leary S, *et al.* Muscle functional MRI as an imaging tool to evaluate muscle activity. *J Orthop Sports Phys Ther* 2011;41:896–903.
- 84 Esposito A, Campana L, Palmisano A, *et al.* Magnetic resonance imaging at 7T reveals common events in age-related sarcopenia and in the homeostatic response to muscle sterile injury. *PLoS ONE* 2013;8:e59308.
- 85 Cermak NM, Noseworthy MD, Bourgeois JM, *et al.* Diffusion tensor MRI to assess skeletal muscle disruption following eccentric exercise. *Muscle Nerve* 2012;46:42–50.
- 86 Froeling M, Nederveen AJ, Nicolay K, *et al.* DTI of human skeletal muscle: the effects of diffusion encoding parameters, signal-to-noise ratio and T2 on tensor indices and fiber tracts. *NMR Biomed* 2013;26:1339–52.
- 87 Zarskaya T, Kumbhare D, Noseworthy MD. Diffusion tensor imaging in evaluation of human skeletal muscle injury. *J Magn Reson Imaging* 2006;24:402–8.
- 88 Nescolarde L, Yanguas J, Medina D, *et al.* Assessment and follow-up of muscle injuries in athletes by bioimpedance: preliminary results. *Conf Proc IEEE Eng Med Biol Soc* 2011;2011:1137–40.
- 89 Askling C, Schache A. *Brukner & Khan's clinical sports medicine*. McGraw-Hill Medical Australia 4th edn. 2011.