CASE REPORT

COLLES FRACTURE DUE TO WEIGHTLIFTING

P. H. ROWE, MA, MB, BChir

*Senior House Officer Orthopaedics, Brook General Hospital, Shooter's Hill, London, S.E. 18**Present Address:**Senior House Officer General Surgery, St. Helier Hospital, Carshalton*

ABSTRACT

The case report of a weightlifter who sustained a Colles fracture is described. Wrist injuries sustained in weightlifting are reviewed. They highlight the need for proper instruction and supervision in this sport.

A 21 year old porter, 73 kilograms in weight, had been weightlifting for 18 months under supervision. He presented in casualty with a painful wrist after lifting 79 kilograms (without supervision) at his club and then having lost control of the weight, stumbled backwards with sudden hyperextension of the wrists and landed on his elbow. A radiograph showed a comminuted fracture of the distal end of the radius, with dorsilateral angulation of the distal fragment and a fracture of the ulnar styloid process. The fracture was treated by closed reduction and immobilisation, healing was uneventful.

DISCUSSION

The object in weightlifting is to lift a heavy weight employing the strongest muscle groups at the point of greatest mechanical advantage, in a subtly efficient summation of forces. Injuries seen in weightlifters mainly involve the knee and back — the knee being particularly vulnerable at the beginning of the lift.

Wrist injuries are less well described. Reilly (1978) states that "deterioration may occur at the wrist" due to hypertension associated with repetitive lifting and holding weights overhead. Ryan and Saliccoli (1976) described 5 adolescent males ranging in age from 14-17 years, with fractures of the distal radial epiphysis; two patients had sustained bilateral fractures. All had been executing the military press at the time of the injury. All of the fractures were of a Type II epiphysis fracture of the distal radius as described by Salter (1963). Of the seven fractures in the five patients, three sustained associated fractures of the distal ulnar not through the

epiphysis of the ulnar. We have seen 2 adolescent weightlifters with fractures of the distal radial epiphysis also within the last year.

This patient presented with a Colles fracture, a previously undescribed injury in weightlifting. These injuries could be prevented by correct instruction in the technique of weightlifting and proper supervision. In

order to prevent these serious wrist injuries it is vital to have a person at each end of the bar to hold it if the lifter loses control.

ACKNOWLEDGEMENT

I would like to thank Mr. D. S. Porter for permission to publish his case.

REFERENCES

- Reilly, T, 1978. "Some observations on weighttraining". Brit.Journ.Sports Med. 12: 45-47.
- Ryan, J. R. and Saliciccoli, G. G., 1976. "Fractures of the distal radial epiphysis in adolescent weight lifters". Am.Journ.Sports Med. 4: 26-27.
- Salter, R. B. and Harris, W. R., 1963. "Injuries involving the Epiphyseal plate". J.Bone and Joint Surg. 45-A: 587-622.
-

B.A.S.M. TIES AND HEADSQUARES.

As the present stock of ties is now exhausted, another batch is already being manufactured. A wider blade, 4", with the Association's emblem as a multiple motif, will be available on the navy blue background as before, or on maroon. Another design with a single motif and the name of the Association on a stripe is also contemplated if there is a demand.- £3.00 + 20 pence postage & packing.

At the request of many lady members, headsquares with the Association's emblem on two corners are on order. - £2.50, + 20 pence postage and packing.

If there is a demand, woven blazer or track suit badges may be supplied, at approximately £4.00. Hand embroidered wire thread badges would be at least £8 or £9.

Ties and headsquares should be available well before Christmas, and orders should be sent to the Hon. Treasurer. Bank clearance charges of 55 pence should be added to cheques NOT drawn on U.K. banks in sterling, and an additional 25P if airmail is requested.