ovelat[®] Gel nucopolysaccharide, adrenocortical ext., salicylic acid

cooling, soothing & effective



- Direct to the pain
- Direct to the swelling

Movelat[®] Gel

- Tried Tested
- Trusted

The Direct Anti-inflammatory

Presentation: Movelat Gel contains mucopolysaccharide polysulphuric acid ester (heparinoid) 0,2g, adrenocortical extract 1,0g (equivalent to 20mg of corticosteroids), and salicylic acid 2,0g in a colourless gel base to 100g, Movelat Cream is also available containing the same active constituents in a

white vanishing cream base, Indications: Non-rheumatic disorders of the locomotor apparatus: sprains,

Indications: Non-rheumatic disorders of the locomotor apparatus: sprains, muscular strains and spasms, soft tissue injuries.
Rheumatic disorders: osteo-arthritis, fibrositis.

Dosage and administration: Movelat Gel: Two to six inches (5-15cm) to be applied gently to the affected area up to four times daily; may also be used as an ultrasound coupling medium,

Movelat Cream: Two to six inches (5-15cm) to be massaged into the affecte area up to four times daily; may also be used as a dressing, when it should be apolied thickly.

Contra-indications, warnings: Movelat Gel: Contains an alcohol and therefore should not be brought into direct contact with open wounds or mucous membranes.

Movelat Cream: Nil

Legal category: POM

sic NHS Prices: £3,90 per 100g tube of Gel or Cream

Movelat Gel Movelat Cream

0542/0006 0542/5001 Product Licence Holders and Distributors: Farillon Limited, Romford, Essex.



PANPHARMA LIMITED Hayes, Middx. UB4 0JN Tel: 01-561 8774

