INFECTIONS IN SPORT

Proceedings of the Sports Medicine Conference organised by the Edinburgh Post-Graduate Board for Medicine
4th March, 1988

INTRODUCTION

Dr. J. C. M. SHARP, FFCM, MRCP (Glas)

Consultant Epidemiologist, Communicable Diseases (Scotland) Unit,
Ruchill Hospital, Glasgow

The widening spectrum of sporting activities, their global extent and the increasing number of participants mean that almost every kind of human infection may be acquired directly or indirectly, in the pursuance of sport.

Up until 20 years or so ago, sport-related infections were largely confined to septic cuts and abrasions, possibly tetanus, the appropriately named "athletes foot" and other fungal infections, respiratory infections such as "flu" or "sore throat" or even on occasion pulmonary tuberculosis as happened to several players in a Scottish rugby club in the late 1940s.

While the potential or reality of most of these infections remain, some new ones have emerged in more recent years to cause additional concern. "Scrupox" in one form or another has always been around, but the fear of herpes has nowadays taken on a new dimension along with viral hepatitis and AIDS. Other viral infections identified in recent years include the Coxsackie group of enteroviruses, which is known to be associated with myalgic encephalomyelitis — the "post-viral fatigue syndrome".

With the range of sporting activities nowadays involving overseas travel on club tours or in competition, and especially so if to Africa or Asia, an increasing number of athletes and officials become exposed to additional hazards such as travellers' diarrhoea in its various guises (e.g. cholera or typhoid) or to malaria, certain types of which may result in death.

On Friday, 4th March, 1988, the morning session of the annual Sports Medicine Conference arranged by the Edinburgh Post-Graduate Board for Medicine, was devoted to "Infection in Sport". Five papers were presented by various speakers from general practice, bacteriology and infectious disease.

INFECTIONS ASSOCIATED WITH SPORT

Dr. R. W. A. GIRDWOOD, FRCPath

Consultant Microbiologist, Stobhill General Hospital, Glasgow

Sport associated infections can be divided conveniently into (1) infections acquired by direct person-to-person contact (2) infections acquired by the participant's involvement with the sport-related environment e.g. changing room, playing field and (3) infections acquired during foreign travel (J. H. Cossar — vide infra). In addition to the most frequently encountered topical infections some of the rarer systemic infections will also be mentioned.

Infections by Person-to-Person Contact

Infections acquired by this route include impetigo, erysipelas, herpes simplex and tinea barbae, all of which can be transmitted by the grinding physical contact found in such sports as wrestling and rugby (especially in the scrum)

— hence their frequently being known as "scrumpox". Stud abrasions and the wearing of facial stubble facilitate the inoculation of the infecting agent.

Impetigo is a superficial bacterial infection of the skin caused by *Staph. aureus* or *Strep. pyogenes*, and is characterised by small crusted spots, somewhat similar to pustular acne. Erysipelas is an acute spreading skin infection due to *Strep. pyogenes*, usually involving the face and is characterised by redness, pain and swelling. In both conditions some of the strains of *Strep. pyogenes* can cause kidney damage in the form of glomerulonephritis. A well documented example of rugby acquired skin infection resulting in glomerulonephritis was described by Ludlam and Cookson (1986) who use the term "scrum kidney".

Prevention of the spread of such infections is best achieved by a combination of a responsible attitude of the individual participant and clinical awareness of medical and paramedical personnel. Staphylococcal skin infections can be rendered less contagious by using topical antiseptics such as chlorhexidine and povidone iodine. Similarly newly acquired cuts and abrasions, in addition to prompt physical cleaning with soap and water should have topical antiseptics applied. Erysipelas requires treatment with systematic antibiotics.

Infections Acquired from the Changing Facility Environment

The most frequently occurring examples are the superficial dermatophyte infections of the skin, tinea pedis ("athlete's foot") and tinea cruris ("Jock rot"). The infecting fungi are most frequently *T. interdigitale* and *E. floccosum*.

Definitive diagnosis can be made by microscopy and culture of scrapings taken from the scaly edge of lesions. With these anthropophilic dermatophytes infection is from person to person usually via fomites such as towels, sports clothing and shower areas which have become contaminated with infected shed skin scales. Moist, macerated skin is more susceptible to infection and the warm humid environment found in communal changing rooms facilitates spread. It is imperative that there should be no sharing of towels, soaps or sports equipment. The use of prophylactic anti-fungal powders in undergarments and footwear is probably of value. Established infections require prolonged treatment with imidazole creams and sometimes systemic griseofulvin is necessary.

Other infections associated with sports complexes include pseudomonas folliculitis and legionella infections. Pseudomonas folliculitis is associated with whirlpool spas and jacuzzis. Infection occurs by a combination of maceration of the skin from excessive immersion and contamination of the water with *Pseudomonas aeruginosa*. Prevention is by regular maintenance of the facility and adequate chlorination of the water. In recent weeks in Scotland, infection by *Legionella micdadei* has been associated with a sports and leisure complex. Design and regular inspection and maintenance with particular reference to water storage and delivery systems are essential if this potential hazard is to be avoided.