Metabolic predictors of middle-distance swimming performance

J.P. Ribeiro, MD, ScD, E. Cadavid, MD, J. Baena, MD, E. Monsalvete, MD, A. Barna, MD and E.H. De Rose, MD

Exercise Research Laboratory, Federal University of Rio Grande do Sul, Porto Alegre, Brazil

To evaluate the capacity of different metabolic indices to predict performance in middle distance swimming, 15 competitive swimmers performed a submaximal and a maximal 400 metres freestyle swimming event. Expired gases were collected in Douglas bags immediately after the events for the determination of $\dot{V}O_2$ max. Arterialized blood samples were collected for the determination of maximal blood lactate concentration and the velocity corresponding to blood lactate concentration of 4 mM. The results demonstrated (means $\pm$ SD): maximal velocity of 1.44 ± 0.05 m.s⁻¹; velocity at 85 percent of $\dot{V}O_2$ max of 1.36 ± 0.04 m.s⁻¹; velocity at 4 mM of 1.32 ± 0.04 m.s⁻¹; $\dot{V}O_2$ max of 3.47 ± 0.5 l.min⁻¹; maximal blood lactate concentration of 11.8 ± 2.5 mM. Multiple regression analysis relating metabolic indices and maximal velocity demonstrated that only velocity at 85 percent of $\dot{V}O_2$ max ($r^2 = 0.81$) and velocity at 4 mM ($r^2 = 0.79$) were significant predictors. Thus, 79 percent of the variance in the performance of 400 m freestyle can be accounted for the velocity at 85 percent of $\dot{V}O_2$ max or the velocity at 4 mM. The success in this event seems to depend on the swimmer's capacity to achieve higher velocities with lower blood lactate levels and/or utilizing a lower percentage of their $\dot{V}O_2$ max.

Keywords: Maximal oxygen uptake, anaerobic threshold, blood lactate

It has been repeatedly shown that performance in long distance running and cycling is closely related to the individual's maximal oxygen uptake ($\dot{V}O_2$ max), to the percentage of $\dot{V}O_2$ max taxed during the event, and to ventilatory or lactate thresholds¹⁻⁸. However, these findings cannot be generalized to swimming due to the great variability in the swimming economy among different swimmers⁹. Costill *et al.*¹⁰ found a poor correlation between $\dot{V}O_2$ max and the performance in the 365.8 metres freestyle event, while Olbrecht *et al.* found a high correlation between the swimming velocity corresponding to the blood lactate concentration of 4 mM and the performance on a 30 minute swimming event¹¹. The 400 metres freestyle is an event in which both aerobic and anaerobic processes contribute to the energy release. However,

little is known about the value of different metabolic indices in the prediction of performance in this event¹².

This lack of data may be related to the fact that evaluation of the energy releasing processes in swimming is difficult to perform in most exercise physiology laboratories. However, new methods have been developed which allow the determination of $\dot{V}O_2$ and blood lactate concentration in response to different swimming velocities with the collection of expired air and blood samples in the recovery period^{13,14}. These methods present the advantage of estimating the contribution of anaerobic and aerobic processes at submaximal and maximal intensities while swimming, and relating these findings to training intensities^{10,15}. Thus, the present study was conducted to evaluate the capacity of metabolic indices derived from these new methods to predict the performance in the 400 metres swimming event. The results indicate that some of these factors may be important determinants of middle distance swimming performance.

Method

Fifteen competitive swimmers participated in the study. Their physical characteristics are described in Table 1. They were participating in intense physical training, they were all used to swimming the 400 metres freestyle swimming event, but not all of them were specialists in such an event. The athletes were informed about the risks involved in the participation in the experiments and signed an informed consent.

Table 1. Physical and performance characteristics of swimmers

	Mean	SD	Range
Age (years)	16	2	13-19
Weight (kg)	63	8	50-79
Height (cm)	171	4	159-185
Lean body mass (kg)	56	7	44-69
Water temperature (°C)	28.5	0.6	27-29
Velocity submax (m.s ⁻¹)	1.34	0.05	1.26-1.41
Velocity max (m.s ⁻¹)	1.44	0.05	1.35-1.54
Velocity 4 mM (m.s ⁻¹)	1.32	0.04	1.25-1.37
Velocity 85% $\dot{V}O_2$ max (m.s ⁻¹)	1.36	0.04	1.30-1.43

Address for correspondence: Jorge Pinto Ribeiro, Laboratorio de Pesquisa do Exercício, Universidade Federal do Rio Grande do Sul, Rua Felizardo 750, 90610 Porto Alegre RS, Brazil

© 1990 Butterworth-Heinemann Ltd
0306-4179/90/030196-05

At least 12 hours after the last training session, each swimmer performed two 400 metres freestyle swimming trials in an indoor swimming pool 25 metres long, according to the protocol described by Costill *et al.*¹⁰ and Olbrecht *et al.*¹¹. These trials were preceded by a warm-up period in which the athletes swam to get used to the swimming pool. The water temperature was maintained between 27 and 29 degrees centigrade, and the events were performed with a 20 minute interval period in which the swimmers remained seated out of the water.

In the first trial the athletes were asked to swim at a constant velocity in order to achieve 90 percent of their maximal velocity for the event. On the second trial the swimmers were asked to achieve their best time for the event. The subjects who could not achieve these goals came back on another day to swim at the appropriate velocities. During the recovery period of each trial, expired gas and blood samples were collected to evaluate the metabolic indices.

One stroke before the end of the events, the swimmers retained their breath, exhaling the air in a three-way respiratory mask as soon as the operator put it on their faces¹⁰. The face mask was connected by a short tube to a Douglas bag. The expired gas was collected over the first 20 seconds of the recovery period. Afterwards, air samples from the bags were collected in oiled syringes for analysis of O₂ and CO₂ concentration by the Sholander technique¹⁶, and the expired volume was measured in a dry gas meter. Conventional formulas¹⁷ were used to calculate the oxygen uptake for the submaximal ($\dot{V}O_{2\text{ submax}}$) and maximal ($\dot{V}O_{2\text{ max}}$) events. No correction was made for the backward extrapolation¹⁸ of these values because all subjects were studied in the same conditions. The results obtained from this method have been shown to be valid and reproducible^{10,13,18}.

$\dot{V}O_{2\text{ max}}$ was expressed in $\text{l}\cdot\text{min}^{-1}$, considering the body mass in $\text{ml}\cdot(\text{kg}\cdot\text{min})^{-1}$, and considering the lean body mass estimated by skinfold thickness¹⁹ in $\text{ml}\cdot(\text{kg LBM}\cdot\text{min})^{-1}$. $\dot{V}O_{2\text{ submax}}$ was expressed in $\text{l}\cdot\text{min}^{-1}$ and as a percentage of $\dot{V}O_{2\text{ max}}$. The results of $\dot{V}O_{2\text{ max}}$ and submax were plotted as a function of the swimming velocity in each of the events so that the velocity corresponding to 85 percent of $\dot{V}O_{2\text{ max}}$ could be interpolated or extrapolated for each individual. Figure 1 illustrates this procedure for two of the swimmers.

Twenty microlitre arterialized blood samples were collected from the ear lobe immediately before the events and at one, three and five minutes in the recovery period for later analysis in duplicate by a modified enzymatic method²⁰. The samples were deproteinized in 200 microlitres of perchloric acid and centrifuged for five minutes at 12000 rpm. Fifty microlitres of the supernatant were pipetted in 500 microlitres of a glycine hydrazide buffer solution, 50 microlitres of a NAD solution, and five microlitres of a lactate dehydrogenase solution. The assay was incubated for 60 minutes at 25 degrees centigrade, and the absorbance was read in a spectrophotometer at 364 nm. In our laboratory, the coefficient of variation for two samples is 4.5 percent for resting

Figure 1. Percentage of $\dot{V}O_{2\text{ max}}$ related to swimming velocity in two of the swimmers. For swimmer A the velocity corresponding to 85 percent of $\dot{V}O_{2\text{ max}}$ was extrapolated and for swimmer B it was interpolated. $\dot{V}O_{2\text{ max}}$ in swimmer A was $3.49 \text{ l}\cdot\text{min}^{-1}$, which corresponded to $60 \text{ ml}\cdot(\text{kg}\cdot\text{min})^{-1}$. $\dot{V}O_{2\text{ max}}$ in swimmer B was $3.09 \text{ l}\cdot\text{min}^{-1}$, which corresponded to $50.9 \text{ ml}\cdot(\text{kg}\cdot\text{min})^{-1}$.

and 5.1 percent for exercising conditions. The highest blood lactate concentration in the recovery period was used as the representative for each swimming velocity. These results were plotted as a function of swimming velocity and the velocity corresponding to the blood lactate concentration of 4 mM was determined by interpolation or extrapolation, as shown in Figure 2¹⁴. This intensity was arbitrarily chosen to represent the blood lactate response to submaximal swimming.

Figure 2. Blood lactate concentration related to swimming velocity for the same swimmers described in Figure 1. For both athletes the swimming velocity corresponding to a blood lactate concentration of 4 mM was extrapolated.

Descriptive data are presented as mean $\pm$ standard deviation. To determine the capacity of the metabolic indices to predict swimming performance in the 400 metres freestyle event, multiple linear regression analysis with the maximal swimming speed as the dependent variable was used²¹. Since six metabolic indices were used as independent variables, the Bonferroni correction was applied and the level of probability required for significance was 0.008 (0.05 divided by the number of variables)²².

Results

Table 1 contains the average performance of the swimmers. The best time for the event varied between 4min 19s, 86 hundredths, for the best swimmer and 4min 56s, 68 hundredths, for the swimmer with the poorest result. In the submaximal trial, the swimmers attained a speed which ranged from 89 to 95 percent of the velocity attained in the maximal trial. These findings resulted in metabolic findings which allowed the appropriate interpolation or extrapolation of the variables for all swimmers.

Table 2 contains the results for $\dot{V}O_2$ max and submax, as well as the blood lactate concentration before and after each trial. It is of note that the blood lactate concentration before the maximal event is higher than the one before the submaximal event. This was probably due to the fact that blood lactate levels had not returned back to baseline during the 20 minute interval period between the events.

Table 3 shows the simple correlation coefficients for each of the metabolic indices and the maximal swimming velocity in the 400 metres event. Only the velocity corresponding to 85 percent of $\dot{V}O_2$ max and

the velocity corresponding to the blood lactate concentration of 4mM correlated significantly to maximal swimming performance.

Figure 3 shows scattergrams for these associations. The maximal blood lactate concentration as well as the $\dot{V}O_2$ max expressed in absolute values and relative to body composition did not correlate significantly to maximal swimming performance. Multiple linear regression analysis showed that when the velocity corresponding to 85 percent of $\dot{V}O_2$ max and the velocity corresponding to the blood lactate concentration of 4mM were entered in the model, a multiple correlation coefficient of 0.91 was found ($r^2 = 0.83$, $p < 0.001$) with the maximal swimming velocity. When the velocity corresponding to the blood lactate concentration of 4 mM was added to the velocity corresponding to 85 percent of $\dot{V}O_2$ max as a

Table 2. Metabolic data in 15 swimmers

	Mean	SD	Range
$\dot{V}O_2$ submax (l.min ⁻¹)	2.83	0.50	1.91–3.67
$\dot{V}O_2$ submax (% $\dot{V}O_2$ max)	81	5	75–92
$\dot{V}O_2$ max (l.min ⁻¹)	3.47	0.53	2.56–4.52
$\dot{V}O_2$ max (ml.[kg.min] ⁻¹)	56	6	45–62
$\dot{V}O_2$ max (ml.[kgLBM.min] ⁻¹)	63	6	50–70
Lactate pre-submax (mM)	1.6	0.4	1–2.5
Lactate pre-max (mM)	2.3	0.8	1.2–4.0
Lactate submax (mM)	5.7	2.0	3.1–9.8
Lactate max (mM)	11.8	2.5	7.8–17.2

Table 3. Correlation between maximal swimming velocity in the 400 metres freestyle and metabolic indices

Index	r	r ²	p
Velocity at 85% $\dot{V}O_2$ max	0.90	0.81	< 0.001
Velocity at 4 mM	0.89	0.79	< 0.001
Lactate max	0.42	0.18	> 0.05
$\dot{V}O_2$ max (l.min ⁻¹)	0.30	0.09	> 0.05
$\dot{V}O_2$ max (ml.[kg.min] ⁻¹)	–0.08	0.006	> 0.05
$\dot{V}O_2$ max (ml.[kgLBM.min] ⁻¹)	0.07	0.005	> 0.05

Figure 3. a: Relation between the swimming velocity corresponding to 85 percent of $\dot{V}O_2$ max and maximal velocity is depicted; b: Relation between the velocity corresponding to 4mM and the maximal swimming velocity is presented.

second predictor, the predictability did not increase significantly. Thus, both the velocity corresponding to 85 percent of $\dot{V}O_2$ max and the velocity corresponding to 4 mM blood lactate concentration were found to be good isolated predictors of swimming performance in the 400 metres freestyle swimming event.

Discussion

In the 400 metres freestyle swimming event both aerobic and anaerobic energy releasing processes are activated. For this reason, it was our hypothesis that the maximal performance in such an event should be related to metabolic indices representative of both processes. It was a surprise to find out that $\dot{V}O_2$ max expressed in $l \cdot min^{-1}$ and relative to body composition did not correlate to maximal swimming velocity in an event which lasts four to five minutes. Although the protocol used does not assure that actual $\dot{V}O_2$ max was really attained, the maximal blood lactate concentrations are suggestive of a maximal effort. Moreover, previous studies have validated such protocol for the determination of $\dot{V}O_2$ max^{10,13}. Thus, we believe that the results are representative of the $\dot{V}O_2$ max during swimming for the athletes. Costill *et al.*¹⁰ have found a low, but significant, correlation coefficient ($r = 0.47$) between $\dot{V}O_2$ max and middle distance swimming performance. The fact that we found no significant association is probably related to the more homogeneous swimming performance of our athletes, who had much better results when compared to the swimmers studied by Costill *et al.*¹⁰.

The velocity corresponding to 85 percent of $\dot{V}O_2$ max was shown to be highly correlated to swimming performance in the 400 metres freestyle swimming event. Thus, performance in such an event does not depend directly on the swimmer's $\dot{V}O_2$ max, but on the capacity to achieve higher velocities less taxing on his aerobic reserve. An illustration of the importance of submaximal exercise capacity is shown in Figure 1, where two swimmers are compared. Swimmer A has a $\dot{V}O_2$ max higher than swimmer B. However, the latter shows better economy, as demonstrated by a much lower $\dot{V}O_2$ for a submaximal swimming velocity. To improve swimming performance, an athlete may increase his $\dot{V}O_2$ max or improve his economy. Our data indicate that, at least for the group studied, economy is more important than maximal aerobic power.

The capacity to tolerate higher blood lactate concentrations is considered an important factor in the performance of athletic events which last four to five minutes¹⁵. However, in our sample, there was no significant correlation between maximal blood lactate concentration and maximal velocity. On the other hand, we found a strong correlation ($r = 0.89$) between the velocity corresponding to four mM blood lactate concentration and maximal swimming velocity. This finding is in agreement with the results of other authors who correlated lactate and ventilatory thresholds with running and cycling performance¹⁻⁸.

Likewise, Olbrecht *et al.*¹¹ also found a strong correlation between the velocity corresponding to the blood lactate concentration of 4 mM and the performance in a 30 minute swimming event. However, these authors studied sport events which lasted more than ten minutes, while we studied a much shorter event. Thus, our results expand on this subject, demonstrating that the swimming velocity corresponding to a blood lactate concentration of 4 mM is a good predictor swimming performance in a relatively short lasting event.

As shown in Figure 2, an athlete may improve his swimming performance by reducing the blood lactate concentration at a given speed, or by tolerating higher maximal blood lactate concentrations. Several mechanisms may result in the reduction of submaximal blood lactate levels. The swimmer may increase his $\dot{V}O_2$ max after training, thus a given velocity would represent a lower relative intensity, resulting in lower blood lactate levels. Or the swimmer could, after training, reduce the blood lactate concentration at a given swimming velocity without changing $\dot{V}O_2$ max²³, by changes in the production and/or the metabolism of lactate at a certain percentage of $\dot{V}O_2$ max²⁴.

Finally, the swimmer could improve his economy, reducing the energetic cost for a certain swimming velocity. The fact that both the velocity corresponding to 85 percent of $\dot{V}O_2$ max and the velocity corresponding to 4 mM blood lactate concentration were found to be good predictors of maximal swimming performance suggests that, at least for the swimmers in this study, swimming economy, which is a factor which contributes to both indices, is the most important determinant of swimming performance in the 400 metres freestyle swimming event. However, it is possible that, for groups of athletes with better swimming performance, like the ones studied by Mader *et al.*¹⁴, the adaptations related to lactate metabolism might be more important.

Although the methods used in the present experiments had been previously validated¹¹⁻¹⁵, they have important limitations. The interpolation and extrapolation of the oxygen uptake curves are based on the assumption that, around the intensities studied, there is a linear relation between $\dot{V}O_2$ and swimming velocity, but $\dot{V}O_2$ increases curvilinearly with swimming velocity⁹. Moreover, the method also assumes that there is no plateau in $\dot{V}O_2$ close to $\dot{V}O_2$ max. Likewise, the blood lactate concentration increases exponentially to increments in swimming velocity¹⁴. However, all our swimmers were studied at intensities above 75 percent of their $\dot{V}O_2$ max, which resulted in blood lactate concentration higher than 3 mM, above the point where the most inclined deflections occur in the $\dot{V}O_2$ and blood lactate curves^{9,14}. Thus, an assumption was made that the response was linear at those intensities for interpolation or extrapolation. These limitations indicate that future studies should evaluate a larger number of submaximal swimming velocities.

Another important limitation, which has not been considered in previous studies on the relationship between metabolic indices and maximal perform-

ance, is the possibility of mathematical coupling between the independent and dependent variables²⁵. This occurs when part of a relationship between the variables can be explained by a direct or indirect mathematical relationship. In this study, as well as in previous ones^{3,7,8,11}, metabolic indices were expressed as the swimming velocity (or running velocity) at which the index was measured. Since the dependent variable is also expressed as the certain velocity, it is expected that the athlete who had the higher maximal velocity would also reach a certain metabolic index at a higher submaximal velocity. These comments, however, do not invalidate our findings or the conclusions of previous studies, but one should be aware that some of the relationships may be overestimated because of mathematical coupling²⁵.

Conclusion

The present study demonstrates that 79 percent of the variability of the swimming performance in the 400 metres swimming event can be accounted for in the swimming velocity corresponding to 85 percent of $\dot{V}O_2$ max or the velocity corresponding to a blood lactate concentration of 4 mM obtained by a relatively simple testing procedure. Both metabolic indices are good isolated predictors of middle distance swimming performance, indicating that the success in the 400 metres freestyle swimming event largely depends on the capacity of the athlete to reach higher swimming velocities with low blood lactate concentrations and/or taxing a lower percentage of $\dot{V}O_2$ max.

Acknowledgements

Drs. Ribeiro and De Rose are CNPq scientists.

References

- Barlow, K., Weltman, A., Schurrer, R. and Henritze, J. Prediction of maximal effort bicycle ergometer endurance performance *Int J Sports Med* 1985, **6**, 190–196
- Costill, D.L. Metabolic responses during distance running *J Appl Physiol* 1970, **28**, 251–255
- Farrel, P.A., Wilmore, J.H., Coyle, E.F., Billing, J.E. and Costill, D.L. Plasma lactate accumulation and distance running performance *Med Sci Sports Exercise* 1979, **11**, 338–344
- LaFontaine, T.P., Londeree, B.R. and Spath, W.K. The maximal steady state versus selected running events *Med Sci Sports Exercise* 1981, **13**, 190–191
- McLellan, T.M. and Skinner, J.S. Submaximal endurance performance related to the ventilation thresholds *Can J Appl Sports Sci* 1985, **10**, 81–87
- Reybrouck, T., Ghesquiere, J., Cattaert, A., Fagard, R. and Amery, A. Ventilatory thresholds during short- and long-term exercise *J Appl Physiol* 1983, **55**, 1694–1700
- Sjodin, B. and Jacobs, I. Onset of blood lactate accumulation and distance running performance *Int J Sports Med* 1981, **2**, 23–26
- Tanaka, K., Matsuura, Y., Matsuzaka, A., Hirakoba, K., Kumagai, S., Sun, S. and Asano, K. A longitudinal assessment of anaerobic threshold and distance running performance *Med Sci Sports Exercise* 1984, **16**, 278–282
- Pendergast, D.R., Di Prampero, P.E., Graig, A.B., Wilson, D.R. and Rennie, D.W. Quantitative analysis of the front crawl in men and women *J Appl Physiol* 1977, **43**, 475–479
- Costill, D.L., Kovaleski, D., Porter, D., Kirwan, J., Fielding, R. and King, D. Energy expenditure during front crawl swimming: predicting success in middle-distance events *Int J Sports Med* 1985, **6**, 266–270
- Olbrecht, J., Madsen, O., Mader, A., Liesen, H. and Hollmann, W. Relationship between swimming velocity and lactic concentration during continuous and intermittent training exercises *Int J Sports Med* 1985, **6**, 74–77
- Costill, D.L. Practical problems in exercise physiology research *Research Quarterly for Exercise and Sport* 1985, **56**, 378–384
- Leger, L.A., Seliger, V. and Brassard, L. Backward extrapolation of $\dot{V}O_2$ max values from the O_2 recovery curve *Med Sci Sports Exercise* 1980, **12**, 24–27
- Mader, A., Madsen, O. and Hollmann, W. Zur Beurteilung der lak- taziden Energiebereitstellung für Trainings- und Wettkampf- leistungen im Sport- schwimmen *Leistungssport* 1980, **10**, 263–268
- Troup, J. Setting up a season using scientific training. A guide on how to apply $\dot{V}O_2$ and lactate measures in establishing a training schedule *Swimming Technique* 1986, May–July: 8–16
- Scholander, P.P. Analyser for accurate estimation of respiratory gases in one half cubic centimeter samples *J Biol Chem* 1947, **167**, 235–250
- Consolazio, C.F., Johnson, R.E. and Pecora, L.J. 'Physiological measurements of metabolic functions in men' New York, McGraw, 1963
- Montpetit, R.R., Leger, L.A., Lavoie, J.M. and Cazola, G.A. $\dot{V}O_2$ peak during free swimming using the backward extrapolation of the O_2 recovery curve *Eur J Appl Physiol* 1981, **47**, 385–391
- Faulkner, J.A. Physiology of swimming and diving. In Falls, H.B. (ed.) 'Exercise physiology' New York, Academic Press, 1968, p 417
- Bergmeyer, H.U. 'Methods of enzymatic analysis' New York, Academic Press, 1974, pp 1464–1472
- Kleinbaum, D.G. and Kupper, L.L. 'Applied regression analysis and other multivariate methods' North Scituate, Duxbury, 1978, p 131
- Wallenstein, S., Zucker, C.L. and Fleiss, J.L. Some statistical methods useful in circulation research *Cir Res* 1980, **47**, 1–9
- Hurley, B.F., Hagberg, J.M., Allen, W.K., Seals, D.R., Young, J.C., Cuddihee, R.W. and Holloszy, J.O. The effect of training on blood lactate levels during submaximal exercise *J Appl Physiol* 1984, **56**, 1260–1264
- Brooks, G.A. Anaerobic threshold: review of the concept and directions for future research *Med Sci Sports Exercise* 1985, **17**, 22–31
- Archie, J.P. Mathematical coupling of data *Ann Surg* 1981, **193**, 296–303