Overtraining Syndrome

R. Budgett, MBBS, MA, Dip Sports Med, MRCP

The British Olympic Medical Centre

This review discusses the overtraining syndrome which is characterized by fatigue and underperformance precipitated by stress of training. Other stresses, depression and an increased susceptibility to infections may be important. Treatment requires rest and a stress management program over 3 months.

Keywords: Stress, underperformance, overtraining syndrome, staleness, chronic fatigue

Introduction

There have been numerous anecdotal reports of athletes underperforming for no apparent reason and taking weeks to recover form. The term 'overtraining syndrome' is often used to describe these athletes suffering from prolonged fatigue. This term implies causation, but there is no firm evidence for this, and athletes who complain of fatigue and underperformance may not all be suffering from a single pathophysiological entity.

Other terms such as 'burnout', 'staleness', and 'chronic fatigue syndrome' have been used to describe a similar picture. Some commentators feel the problem is post-viral in origin linking it with ME¹. The picture is complicated by a growing number of reports of immune dysfunction after prolonged exhaustive training, so that frequent or prolonged infections causing fatigue and malaise may be secondary to overtraining rather than the primary cause of the problem.

Definitions

Fatigue

This is a subjective condition synonymous with tiredness, lethargy and listlessness, with symptoms such as poor concentration and poor tolerance of activity. It contrasts with physiological fatigue which can be defined as the failure to sustain an expected or required muscular force. The mechanism depends on the duration of exercise, so that a sprinter will become fatigued in seconds in association with high

lactate levels, while a marathon runner will fatigue after nearly 2 hours due to glycogen depletion³. Fatigue in this review refers to the subjective condition.

Overtraining syndrome

1. This may be seen as prolonged fatigue and underperformance, following a period of heavy training or competition, lasting at least two weeks and confirmed on an appropriate ergometer, or by comparing training times or results in previous competition. Identifiable medical illness must have been excluded. This assumes that the stress of training is the major factor in the aetiology of prolonged fatigue in athletes.

2. Athletes who suffer recurrent infections during periods of maximum training or competition stress may suffer from overtraining syndrome.

This assumes that overtraining is the major stress causing immune dysfunction, producing increased vulnerability to infection.

Burnout/staleness

The definition is as for 1. in the overtraining syndrome. It is often used to describe swimmers⁴⁻⁵.

Chronic fatigue syndrome

This is the same definition as 1. in the overtraining syndrome. The term is used in American literature^{6,7}.

Post-viral infection/ME

This causes fatigue lasting many months following a viral, possibly enteroviral, infection⁸. Patients are unable to function in normal life let alone train. Few athletes fall into this category. There is no evidence that the proposed reduction in immune function in overtrained athletes predisposes to post-viral fatigue, so this cannot yet be defined as part of the overtraining syndrome.

Sympathetic and parasympathetic staleness

These terms have been used to sub-divide the overtraining syndrome, but are not generally used now due to the lack of evidence for a separate parasympathetic type which was supposedly characterized by hypoglycaemia during exercise and

Address for correspondence: Dr R. Budgett, The British Olympic Centre, Northwick Park Hospital, Watford road, Harrow, Middlesex HA1 3UJ, UK

© 1990 Butterworth-Heinemann Ltd
0306-3674/90/040231-06

Overtraining Syndrome: R. Budgett

exhaustion of the neuro-endocrine system. The sympathetic type was defined as essentially a response to prolonged stress and fits with overtraining definition 1.⁹⁻¹¹

Over-reaching

Purposeful overtraining may be part of a planned programme in order to stimulate adaptation. This is followed by a period of relative rest or tapering in order to allow full supercompensation, normally within 1-2 weeks¹².

Clinical picture

As suggested by the two definitions, there are two broad clinical presentations which may overlap.

Athletes may suffer from fatigue and a performance slump which does not improve despite resting over 2 days or reducing the training load for 2 weeks. They are often highly motivated, conscientious individuals who have responded to their lack of form by increasing the training load, leading to a vicious cycle of fatigue and underperformance¹².

Others have been entering a series of hard competitions, often with initial success, but then begin to become tired and produce steadily worse results¹³. These athletes are characterized by their reluctance to stop or reduce their level of training and competition due to their own ambition and determination sometimes combined with pressure from the coach, fellow athletes and parents.

Other elite competitors build up their training for two to three months, reaching a high level of performance, but then suffer from a viral infection which stops them in their tracks. This often occurs around the time of a major competition or series of races. After enforced rest for 1-2 weeks they restart training, but when they reach a certain level again break down and the cycle repeats itself^{14, 15}. As yet, there is only indirect evidence of immune dysfunction (Riddell: personal communication).

Reports of athletes presenting in this way are nearly all anecdotal, but the constant flow of reports from coaches and athletes show that they perceive there is a problem despite the fact that it is poorly defined¹². A large number of symptoms and signs have been described, but some of them relate to athletes who are only over-reaching, and others may represent underlying unidentified disease since they are inconsistently reported in different studies. There is no accepted gold standard for the diagnosis of the so called overtraining syndrome.

Symptoms

Athletes can only be labelled as overtrained if they are fatigued and underperforming. They may complain of fatigue at rest, with a loss of purpose, energy and competitive drive^{16, 17}. Feelings of helplessness, incompetence, being trapped in a routine, and emotional lability and loss of libido have been described^{10, 13}. Athletes often admit to increased

anxiety and depression (90%), irritability (70%) and sleep problems (90%) including difficulty getting to sleep, nightmares and waking in the night¹⁸. Loss of appetite, weight loss, heavy painful muscles and excessive sweating have been reported^{4, 11, 13}. Athletes have also been noted to complain of an increased susceptibility to infections^{2, 16} (Riddell: personal communication) and injuries¹⁹.

Signs

There are no physical signs consistently associated with the overtraining syndrome. The following have been reported in association: raised early morning heart rate, increased postural drop in blood pressure and rise in heart rate, and a slow return of pulse rate to normal after exercise^{20, 21}, decreased maximal lactate levels during exercise and decreased maximal power output¹⁰. There may be lymphadenopathy, diarrhoea, increased susceptibility to allergies and minor scratches may heal slowly¹³ (Riddell: personal communication). However, none of these signs are reliable indicators of the syndrome, since they can appear during periods of over-reaching without any drop in performance. They are reported inconsistently and some may represent underlying disease.

Precipitating factors

Physical stress

Stresses other than those of hard training and competition may lead to an inability to recover from a previously well tolerated training programme.

Glycogen depletion leads to rapid fatigue particularly after the anaerobic threshold is reached, and may predispose to more prolonged fatigue²². Thus, a poor diet with insufficient carbohydrates to meet training demands²³, as may occur when an athlete is attempting to lose weight, will increase the difficulty in coping with a particular programme.

Dehydration will also increase the perceived stress of exercise and is a potent cause of underperformance²⁴.

Infections, as well as being reported more frequently in hard training, will make it more difficult for athletes to tolerate the load.

Psychological stress

The profile of mood states questionnaire asking athletes to rate their current feelings or mood against a list of 65 words has been used to follow them in training. It shows reduced vigour and increased tension, depression, anger, fatigue and confusion (inverted iceberg profile) during periods of heavy training^{25, 26}. This is even worse in those athletes who suffer from prolonged fatigue and are labelled as stale or overtrained^{27, 28}. It has been suggested that academic, economic and social stressors may be important. They certainly feature prominently in anecdotal reports. However, there is no good evidence for their role due to the lack of studies on competitors who become overtrained.

Differential diagnosis

Serious disease

This is fortunately rare but it is important to exclude viral myocarditis or arrhythmias, bearing in mind that electrocardiograms are often abnormal, particularly at rest, in highly trained athletes²⁹.

Glycogen depletion

Prolonged inadequate carbohydrate intake may cause prolonged fatigue as well as precipitating a true overtraining syndrome.

Anaemia

It is important to distinguish between true anaemia which may be due to iron deficiency, particularly in female athletes³⁰, and the physiological reduction in haemoglobin due to increased plasma volume which has no effect on performance. Low iron stores as measured by serum ferritin levels were thought to be responsible for many cases of underperformance³¹, but it has been shown that levels as low as 12 ng/ml are not associated with underperformance. There is no advantage in raising this with iron supplements³².

Exercise induced asthma

This diagnosis is easily missed unless enquired for directly, and is a common cause of apparent recurrent colds and underperformance³³. A 15% or greater fall in peak expiratory flow rate or FEV1 after exercise is considered diagnostic³⁴.

Physical changes due to age

These are sometimes difficult to accept! The change in shape of female athletes, such as gymnasts or swimmers, at menarche is particularly dramatic if they have a few months off and put on weight. This will greatly affect performance but should not cause fatigue (P. Garret: personal communication).

Psychological

There is a psychological element to every athlete's underperformance, but in some this is of overriding importance so that the training stress is irrelevant. In these cases the athlete cannot be labelled as overtrained, and psychological help is far more important than rest from training.

Prevention

It is obviously important to avoid the precipitating factors discussed earlier, but early detection and a carefully designed training programme may also play a role in prevention.

Early detection

The careful monitoring of athletes who are overreaching has been suggested in order to pick up true overtraining or staleness in the early stages²¹. In

general, the longer the history of fatigue before training is stopped, the longer recovery takes¹². The drop in performance often become obvious too late to avoid a prolonged period of fatigue. Because of this, early morning heart rate²⁰ and mood state questionnaires²⁷ have been used. Others have suggested using serial measurements of haemoglobin, haematocrit or creatine kinase. Unfortunately, these tests are non-specific and reflect training load or may not even change at all^{10,16}. Thus, there is no objective early screen that can be reliably used to detect those athletes who will become overtrained during a period of overreaching.

Training

There is consensus that undertraining is better than overtraining, and this seems to be the surest way of avoiding prolonged fatigue²¹. However, maximal training is needed for an athlete to reach their full potential and various strategies have been suggested to allow this while minimizing the risk of the overtraining syndrome. Sudden increases in training load should be avoided, a steady increase of 5% per week being optimal for most athletes. Intensive exercise with short rests and frequent competition is dangerous. Monotonous training should also be avoided, particularly in swimmers¹¹. Notice should be taken of other stresses, both physical and psychological, since these may combine and exceed an individual's ability to adapt. Training should be reduced at these times⁹.

Management

Rest

Cases need to be assessed individually, and it is often difficult to persuade athletes that they need to rest. However, this seems to be the basis of treatment since an improvement in performance has been shown after 3–5 weeks rest, despite the fact that these athletes must become detrained in this time²⁸. The build up to full training may take up to three further months. It is thought important to avoid competition¹³, although individuals may produce personal best performances 'out of the blue' before apparent complete recovery⁶.

In the majority of athletes who are still able to work and function in normal life, a regime of relative rest has been suggested using gentle non-competitive exercise often in a different sport with no measure of performance available. Usually this low level of exercise is easily tolerated and the psychological and physical benefits will help speed recovery³⁴, while maintaining a degree of cardiovascular fitness. The emphasis should be on exercise as a therapeutic tool to speed rehabilitation, rather than as a means to obtain a training effect which is what the athletes are used to.

Regeneration

No controlled trials of treatment have been reported, but regeneration strategies have been widely used

particularly in the Eastern bloc³⁵. This involves a reduction in all stresses with rest, sleep, relaxation therapy and counselling as well as saunas, massage, hydrotherapy and physiotherapy⁵. Nutrition is looked at in detail and vitamins and other supplements given routinely although there is no proven benefit. Psychotherapy and counselling have been used in the USA to successfully treat underperforming athletes diagnosed as stale and depressed²⁵, but there are no reports on the use of antidepressants. However, there is one report of anabolic steroid use in an attempt to speed recovery³⁶, and no doubt others have tried this unproven and illegal tactic without reporting their results. Our own anecdotal observation suggests that recovery takes three to eight weeks. However, there is a danger of relapse at around 3 months. We advise reduced training and competition stress for up to 4 months.

Investigating the overtraining syndrome

There are only a few studies of truly overtrained elite competitors^{18, 25, 27, 28, 37} (Riddell: personal communication). Most investigators have concentrated on studying the response to severe exhaustive exercise in an attempt to find markers of and mechanisms to explain the overtraining syndrome. This ignores the previously discussed problem that overtrained competitors may be training no harder than their peers, and that it is something unique to that individual that causes prolonged breakdown with failure to adapt or benefit from training. Moreover, in those studies in which performance is monitored, there is generally a quick recovery or even an improvement at the end of the period of extreme over-reaching³⁸.

Routine tests as markers

Haemoglobin and haematocrit have been shown to fall during periods of heavy training³⁹, but this is probably physiological, and does not correlate with performance. Others have found no change¹⁶. ECGs are often abnormal in fit athletes and are no different during periods of heavy training or underperformance²⁹.

Creatine kinase has been used by many researchers, but unfortunately has been shown to rise in proportion to training load, presumably due to muscle damage, does not correlate with performance, and there is wide individual variability (50 fold)^{13, 20}. In a single study, thyroxine and TSH were reduced during a period of prolonged physical and mental stress, but levels did not correlate with performance and recovery took only two days.

Hormonal and endocrine changes

Cortisol is catabolic, helping to mobilize glycogen and free fatty acids, and rises after hard exercise as part of the stress hormone response^{10, 40, 41}. Psychological stress also raises levels, particularly before competition⁴². Testosterone has also been shown to rise in response to exercise, and there is evidence that levels are higher in well trained individuals^{17, 43, 44}.

Intense and prolonged training will raise cortisol levels even higher, but will reduce testosterone⁴⁵⁻⁴⁷. Thus, it has been suggested that hormonal changes are involved in or at least are markers of the overtraining syndrome¹⁷. It may be that at a critical testosterone/cortisol ratio the predominance of catabolism over anabolism means that an athlete is unable to recover from training, and continues to break down rather than adapting positively.

Harkonnen *et al.* looked at 15 long distance runners, and increased the training load of nine of them for just one week⁴⁸. The testosterone/cortisol ratio fell in eight of the nine who were classified as overtrained and in only one control. There was no record of chronic fatigue or underperformance. These athletes were therefore not truly overtrained, but merely exhibited a normal response to extreme training stress.

O'Connor *et al.* followed salivary cortisol levels in 14 female swimmers over 6 months²⁷. As well as the expected rise in baseline levels as training increased from 2,000 to 12,000 yards per day, they found significantly higher levels in those suffering from staleness and underperformance compared to those maintaining their performance. This correlated with depressed mood state²⁷.

Barron and Noakes in 1985 suggested that the overtraining syndrome was due to hypothalamic-pituitary dysfunction³⁷, as may be the case in female athletes who develop amenorrhoea. They found a significantly reduced hormonal response to insulin induced hypoglycaemia in four chronically fatigued and underperforming athletes compared to controls. However, basal cortisol levels were raised, making the interpretation of the results difficult, but further supporting the existence of high basal cortisol levels in overtrained athletes. Testosterone was not measured.

Mood state

Some commentators feel that the chronic fatigue seen in athletes is 'depression with a new face'. It is possible that the stress of heavy training and competition, combined with other stresses, can precipitate depression. This would certainly explain why some athletes are more susceptible to the so called overtraining syndrome than others. There is no doubt that the athletes are depressed with changes in global mood as measured by the Profile of Mood States (POMS) questionnaire^{27, 28}.

The symptoms of overtraining are very similar to those described in depression with psychomotor and physiological retardation, chronic fatigue, depressed appetite, weight loss, insomnia, muscle soreness, and tension and depression. However, the mood state changes in response to an increased training load are non-specific, with a decrease in vigour and increase in tension, depression, anger, vigour and confusion in all athletes, so it is difficult to use a mood state questionnaire to predict or diagnose the overtraining syndrome. Despite this, there is evidence that this change is even greater in underperforming athletes²⁷ and so may potentially be used to

support the diagnosis and monitor recovery since mood state improves with rest in parallel with performance²⁸.

Immune function

Anecdotal reports have linked susceptibility to infections to periods of heavy training, sometimes associated with prolonged fatigue and underperformance^{14,15} (Riddell: personal communication).

Many studies have investigated the effect of exercise on the immune system, and it is now accepted that there is a release of white blood cells causing a temporary leucocytosis after heavy exercise². However, there is no evidence that this of functional significance.

Some studies have shown a reduced T helper: T suppressor cell ratio after hard exercise^{49,50}, but this did not correlate with temporary suppression of lymphocyte function which has also been described⁵¹. Makinon showed reduced salivary IgA and IgM concentrations and reduced NK cell activity after one to two hours of heavy exercise⁵². However, these changes resolved soon after exercise, and are not found consistently. Nash showed that training enhanced immune function⁵³. Thus, there is still uncertainty about the effect of exercise and training on immune function which may depend on the pre-existing state of the athlete, and the type of exercise.

It is difficult to isolate the effect of psychological stress which has been shown to be capable of suppressing immunity² maybe by raising cortisol levels⁵⁴, although immunosuppression may occur in adrenalectomized animals⁵⁵.

Another explanation for the widely reported, but as yet unproven, susceptibility of athletes to infections during periods of chronic fatigue and underperformance, is through the reduced plasma glutamine levels which have been found in overtrained athletes¹⁸. This amino-acid is produced in muscle from the breakdown of other amino-acids⁵⁶, and is an essential fuel for all rapidly dividing cells⁵⁷. It donates nitrogen for purine synthesis, as well as being oxidized to provide energy. It also provides 35% of the energy for lymphocyte metabolism⁵⁸. Thus, low plasma levels may cause immunosuppression, leading to frequent infections and a worsening cycle of underperformance.

Virology

It has been suggested that the majority of cases of chronic fatigue in athletes are post-viral in origin¹. Infectious mononucleosis is a well known cause of chronic fatigue, and there have been cases of proven infection with enteroviruses and coxsackie B¹⁸, but the relevance of high titres to underperformance has been questioned. Subclinical infection or high titres did not correlate with objective or subjective underperformance in a group of 698 athletes¹⁵. In another study of 12 fatigued athletes, there were two positive monospots with atypical mononuclear cells (glandular fever), and two raised coxsackie B titres. However, no cause was found in the other eight⁵⁹.

If a definite viral cause is found, then a diagnosis of the overtraining syndrome cannot be made unless it is part of a pattern of recurrent infections at times of heavy training and competition stress. However, high viral titres have to be interpreted in their clinical context, and may be an incidental finding in fatigued competitors.

Conclusion

Current evidence on the pathophysiology of the overtraining syndrome is limited, but there is no doubt that the problem of unexplained underperformance does exist.

Investigators have suggested hormonal, psychological, immunological and post-viral aetiologies, and it is possible that all these play a part in different athletes. Increased catabolism, depression, infections, and the stress of training combined with other stresses, may all lead to fatigue and underperformance. The stress of training can cause depression, raised cortisol levels and reduced plasma glutamine levels which may all contribute to decreased immune function.

Management consists of rest and stress reduction through a regeneration programme including counselling and relaxation therapy. Recovery takes three to eight weeks.

References

- 1 Kevles, B. Flat tired out: How EBV, the mysterious chronic fatigue syndrome, has brought a group of women cyclists to a grinding halt *City Sports Magazine* 1988, 14 (9), 17-22
- 2 Keast, D. Exercise and the immune response *Sports Medicine* 1988, 5, 248-267
- 3 Brooks, G.A. and Fahey, T.D. Fatigue during muscular exercise *Exercise Physiology* Macmillan Publishers, London, 1984
- 4 Burke E. Blood testing to determine overtraining in swimmers *Swimming technique* 1981, 18 Nov; 29-33
- 5 Crampton, J. Regeneration vs burnout: prevention is better than cure! *Sports Coach* 1987, April/June, 7-11
- 6 Eichner, E. Chronic fatigue syndrome: How vulnerable are athletes? *Phys Sportsmed* 1989, 17 (6), 157-160
- 7 Eichner, E. Chronic fatigue syndrome: Searching for the cause and treatment *Phys Sportsmed* 1989, 17 (6), 142-152
- 8 David, A.S. Postviral fatigue syndrome: Time for a new approach *Br Med J (Clin Res)* 1988, 296, 696-699
- 9 Israel, S. Zur Problematic des Uebertrainings aus internistischer und leistungsphysiologischer Sicht *Medizin und Sport* 1976, 16 (1), 1-12
- 10 Kinderman, W. Das Uebertraining-Ausdruck einer vegetativen Fehlsteuerung *Deutsche Zeitschrift fur Sportmedizin* 1986, H8, 138-145
- 11 Kuipers, H. and Keizer, H.A. Overtraining in elite athletes *Sports Medicine* 1988, 6, 79-92
- 12 Brasher, C. The under-achievers who overtrain *Observer*: January 17th 1988
- 13 Noakes, T. *Lore of Running* Oxford University Press, Cape Town, 1986
- 14 Peters, E.M. Ultramarathon running and upper respiratory tract infections: An epidemiological survey *South African Med J* 1983, 64, 582-584
- 15 Roberts, J.A. Virus illness and sports performance *Sports Med* 1986, 3, 298-303
- 16 Costill, D.L. *Inside Running* Benchmark Press Inc. Indianapolis, 1986

- 17 Adlercreutz, H. Effect of training on plasma anabolic and catabolic steroid hormones and their response during physical exercise *Int J Sports Med* 1986, 7, 27–28 (supplement)
- 18 Budgett, R. et al. The overtraining syndrome/staleness (Abstract) International Olympic Committee World Congress on Sports Sciences, Colorado Springs, 1989
- 19 Barron, D. K. and Mellerowicz, H. Overtraining *Encyclopedia of Sports Sciences and Medicine* (Eds Larson and Leonard), Macmillan, New York, 1971
- 20 Dressendorfer, R.H. Increased morning heart rate in runners: a valid sign of overtraining? *Phys Sportsmed* 1985, 13 (8), 77–86
- 21 Ryan, A. Overtraining in athletes: A round table *The Physician and Sports Medicine* 1983, 11 (6), 93–110
- 22 Costill, D.L. Effects of repeated days of intensified training on muscle glycogen and swimming performance *Med Sci Sports Exerc* 1988, 20 (3), 249–54
- 23 Wheeler, K. Sports nutrition for the primary care physician: The importance of carbohydrate *Phys Sportsmed* 1989, 17 (5), 106–117
- 24 Costill, D.L. Nutrition and dietetics *Olympic Book of Sports Medicine* (Eds Dirix, A., Knuttgen, H. and Tittel, K.), Blackwell Scientific Publications, Oxford, 1988
- 25 Morgan, W.P. Psychological monitoring of overtraining and staleness *Brit J Sports Med* 1987, 21 (3), 107–114
- 26 Costill, D.L. and Morgan, W.P. Mood disturbance following increased training in swimmers *Med Sci Sports Exerc* 1988, 20 (4), 408–414
- 27 O'Connor, P.J. et al. Mood state and salivary cortisol levels following overtraining in female swimmers *Psychoneuroendocrinology* 1989, 14 (4), 303–310
- 28 Koutedakis, Y., Budgett, R. and Fullman, L. The role of physical rest for underperforming elite competitors *Br J Sports Med* (in press)
- 29 Hollman, W. *Sportsmedizin, Arbeits und Trainingsgrundlagen*, F.K. Schattauer Verlag, Stuttgart, 1980
- 30 Haymes, E.M. Nutrition for the female distance runner (ed) Drinkwater, B.L. *Female Endurance Athletes* Human Kinetics Publishers, Champaign, 1986
- 31 Eichner, E. The anaemias of athletes. *Phys Sports Med* 1986, 14 (9), 122–130
- 32 Newhouse, I., Clement, D., Taunton, J. and McKenzie, D. The effects of prelatent/latent iron deficiency on physical work capacity. *Med Sci Sports Exerc* 1989, 21 (3), 263–268
- 33 McCarthy, P. Wheezing or breezing through exercise-induced asthma *Phys Sportsmed* 1989, 17 (7), 125–130
- 34 Szabadi, E. Physical exercise and mental health *Br Med J* 1988, 296, 259–260
- 35 Paikov, V.B. Means of restoration in the training of speed skaters (condensed) *Soviet Sports Review* 1985, 20 (1), 9–12 (Translated from *Konkrobezhniv Sport* 1982, 1, 39–43)
- 36 Kereszty, A. Overtraining *Encyclopedia of Sports Sciences and Medicine* (ed) Larson and Leonard, 218–222, MacMillan, New York, 1971
- 37 Barron, G.L. and Noakes, T.D. Hypothalamic dysfunction in overtrained athletes *J Clin Endocrin Metab* 1985, 60 (4), 803–806
- 38 Costill, D.L. Physiological response to successive days of intense training in competitive swimmers *Med Sci Sports Exerc* 1988, 20 (3), 255–259
- 39 Rushall, B.S. Haematological responses to training elite swimmers *Canadian Journal of Applied Sports Sciences* 1980, 5, 164
- 40 Farrell, P.A. Review of ACTH in exercise *Exercise Endocrinology* Walter de Gruyter & Co., Berlin, 1985
- 41 Carli, G. Hormonal and metabolic effects following a football match *Int J Sports Med* 1986, 7, 36–38
- 42 Petralgia, F. Response of circulating adrenocorticotrophin, beta-endorphin, beta-lipotropin and cortisol to athletic competition *Acta Endocrinol (Copenh)* 1988, 118, 332–336
- 43 Fellmann N. Effect of endurance training on the androgen response to exercise in man *Int J Sports Med* 1985, 6, 215–219
- 44 Kinderman, W. Catecholamines, growth hormone, cortisol, insulin and sex hormones in anaerobic and aerobic exercise *Eur J Appl Physiol* 1982, 49, 389–399
- 45 Aakvaag, A. Hormonal changes in serum in young men during prolonged physical strain *Eur J Appl Physiol* 1978, 39, 283–291
- 46 Opstad, K. Altered physical response to short-term bicycle exercise in young men after prolonged physical strain, calorific deficiency, and sleep deprivation *Eur J Appl Physiol* 1980, 45, 51–62
- 47 Urhansen, A. Behaviour of testosterone, SHBG, and cortisol before and after a triathlon competition *Int J Sports Med* 1987, 8 (5), 305–308
- 48 Harkonnen, M. Biochemical indicators in diagnosis of overstrain condition in athletes *Sp Med and Ex Sci Proceedings of the Olympic Scientific Congress*, July, Eugene, Oregon USA, 1984
- 49 Gmunder, F.K. Effect of long term physical exercise on lymphocyte reactivity: similarity to spaceflight reactions *Aviation Space and Environmental Medicine* 1988, 59, 146–151
- 50 Burke, L.S. and Tan, S.A. The suppressive effect of stress from acute exhaustive exercise on T lymphocyte helper/suppressor cell ratio in athletes and non-athletes (Abstract) *Med Sci Sports Exerc* 1985, 17, 706
- 51 Cameron, A. and Keast, D. 'Haematological profiles and polyclonal lymphocyte function in continuous and intermittent exercise' MSc thesis, University of Western Australia, 1987
- 52 Makinnon, L. The effect of exercise on secretory and natural immunity *Adv Exp Med Biol* 1987, 216A, 869–876
- 53 Nash, H.L. Can exercise make us immune to disease? *Physician and Sports Med* 1986, 14, 250–253
- 54 Besedovsky H, del Ray, A and Sorkin, E. Immunological-neuroendocrine feedback circuits. In: Guillemin et al (eds). *Neural Modulation of Immunity*. Raven Press: New York, 1985
- 55 Stein, M. Bereavement stress and immunity. In *Neural Modulation of Immunity* (ed Guillemin et al.), Raven Press, New York, 1985
- 56 Felig, P. Amino acid metabolism in man *Ann Rev Biochem* 1975, 44, 993–995
- 57 Ardawi, S. and Newsholme, E.A. Glutamine metabolism in lymphocytes of the rat *Biochem J* 1983, 212, 835–842
- 58 Newsholme, E. *Biochemistry for the medical sciences* John Wiley, Chichester, UK, 1983
- 59 Roberts, J.A. Loss of form in young athletes due to viral infection *Br Med J* 1985, 290, 357–358