LETTERS TO THE EDITOR

Radionuclide imaging of gastric emptying

EDITOR, – I would like to comment on the paper by MacLaren *et al*¹ reporting the use of radionuclide imaging to determine gastric emptying during exercise. In their study they could not detect any difference in the rates of emptying of a 5% glucose solution and an iso-osmotic maltodextrin solution that appears to contain ~20% carbohydrate. There is a large body of evidence, accumulated over many years, that shows that carbohydrate concentration has a greater effect on the gastric emptying rate than osmolality within this concentration range.² The findings of this study therefore require examination.

It is known that single field scans, without correction for tissue attenuation, result in significant errors of measurement,³ and may well be the reason for the extremely variable emptying patterns shown in figs 2 and 3 of MacLaren's paper. The use of AUC values to compare emptying rates gives no indication of the shape of the emptying curves and reduces the sensitivity of the technique. I am curious to know what units are used in reporting the AUC values.

Without a measure of the sensitivity of this modification of the scintigraphic technique, or an indication of the power of the study, the value of this paper is debatable. I would suggest that the main conclusion of this study is not that the method was repeatable, but that it is not sensitive enough to detect differences in gastric emptying rates between 5% and 20% carbohydrate solutions.

JOHN LEIPER

Department of Environmental and Occupational Medicine, University Medical School, Foresterhill, Aberdeen AB9 2ZD

- 1 MacLaren D, Miles A, O'Neill I, Crichtley M, Grime S, Stockdale H. Use of radionuclide imaging to determine gastric emptying of carbohydrate solutions during exercise. *Br J Sports Med* 1996;30:20–3
- 2 Vist GE, Maughan RJ. The effect of osmolality and carbohydrate content on the rate of gastric emptying of liquids in man. *J Physiol (Lond)* 1995;482:523–31.
- 3 Collins PJ, Horowitz M, Cook DJ, Harding PE, Shearman DJC. Gastric emptying in normal subjects—a reproducible technique using a single scintillation camera and computer system. *Gut* 1983;24:1117–25.

The authors reply:

In response to Mr Leiper's letter, we offer the following explanations:

(1) We concur with the statement that there is a large body of evidence showing that the energy content of ingested fluid appears to be a primary determinant of gastric emptying, but that is not what we found in this study. The majority of studies on gastric emptying during exercise have focused on the intubation technique—this is the first study to examine gastric emptying directly using scintigraphic techniques while the subject is actually exercising.

(2) It is true that posterior to anterior movement of food from the fundus to the body of the stomach can, when imaging the subject using a single headed gamma camera

(we presume that is what Mr Leiper refers to as "single field scan"), lead to errors in the early assessment of gastric emptying. Dual headed cameras and geometric mean image analysis would help to resolve the problem of tissue attenuation. If Mr Leiper feels that without a dual headed camera and associated depth corrections, gastric emptying studies cannot correctly be performed, then he should be informed that the majority of nuclear medicine departments in the country that are involved in gastric emptying studies do not possess dual headed cameras. Yet these same departments produce clinically acceptable information which is of value to referring clinicians, and helps in the diagnosis and management of gastric disorders.

(3) We are aware of the potential errors in the quantification of gastric emptying studies using a single headed system, and it was for this reason that each subject acted as his own control. We made the assumption that any anatomical movement of the ingested material and any bowel overlap in the region of interest that may occur would be consistent for each individual. This is indeed what we found, for each individual the pattern of emptying on each of the four occasions was remarkably consistent and so repeatable. Any variations that occurred were due to interindividual differences not to intraindividual differences. Examination of figs 2 and 3 clearly reflect these findings. No subjects "dumped" the ingested material, but subjects could be classified as "slow emptiers" or "fast emptiers".

(4) The statement that the area under the curve (AUC) does not carry any information about the shape of the emptying patterns is true, but that is why we published figs 2 and 3. The AUC is an acceptable method of quantification in areas where conventional single exponential, double exponential, or power exponential curve fits are not acceptable. We admit that the sensitivity of AUC decreases as the emptying time increases. Since each subject's curves were normalised to their own maximum image count (which incidentally was the same frame on each of the subjects' four studies), and we never proposed to compare one subject with another, the question of units is irrelevant; however, the units we used were per cent retention time.

We do have to state quite vehemently that variations in gastric emptying found between individuals but not within individuals surely must reflect the fact that the scintigraphic technique we employed is sensitive enough or we would have obtained similar results throughout. The fact that our findings are not consistent with studies using the intubation technique needs elucidating.

D MACLAREN
A MILES
I O'NEIL
M CRITCHLEY
S GRIME
H STOCKDALE
Mountford Building, Byrom Street,
Liverpool L3 3AF

Systematic review of physical activity promotion strategies

EDITOR, – Your review article on systematic review of physical activity promotion strategies' reviews randomised controlled trials of physical activity promotion in apparently healthy, free living adults from the USA. From this it states that these findings do not support the promotion of physical activity in

the United Kingdom by general practitioner exercise prescription schemes.

It is established that the primary care physician is influential in promoting physical activity to patients.^{2–4} My own experiences confirm this. In the 10 exercise prescription schemes operating in East Sussex, initial attendance rates are over 90%. The importance of physical activity is firmly established in the scientific literature. Unfortunately, this is of little value to the general practitioner unless he has a practical means to help bring about a permanent change in lifestyle, incorporating regular and appropriate physical activity. Referring patients for exercise is a strategy to engage sedentary, at risk patients in regular controlled exercise in a safe environment. All general practitioners in the scheme undergo training, supported by a fully referenced manual. The centres are served by staff trained by the first university accredited course on GP referral schemes at the University of Brighton which reassures the GPs of the expected standard of care. The staff aim to educate patients concerning the benefits of activity and risks of continued inactivity, with the ultimate aim of empowering patients to control their own health and exercise more safely in the manner of their choosing. This may well not take place in the leisure centre—only approximately 20% continue their exercise within the centre on completion of the programme. Walking is one of the most commonly employed modes of exercise, contrary to the article's statement.

Raising the level of physical activity in the population is desirable and it would be foolish to surmise that one strategy alone will achieve this. Physician based counselling and referral for supervised activity is only a small part in that jig-saw. There appears to be a lack of understanding of the role of GP referral schemes. But throwing away one part of the jig-saw may ruin the whole picture. Regrettably Hillsdon and Thorogood¹ appear eager to discredit a system unique to the United Kingdom on the basis of American trials, with healthy subjects and only telephone follow up in many cases. The majority of home based programmes described had no postintervention follow up and it is therefore flawed to conclude that these were more likely to result in long term change in physical activity behaviour. While further research (and funding) in this area is required to substantiate long term behavioural change, it is important not to discourage those striving to motivate an at risk section of the population.

While appreciating that this is a review article, I have to confess that I was disappointed to see this paper accepted for publication in its current form. It bears so much resemblance to their previous review⁵ of controlled trials of physical activity that the introduction, discussion and "future research" merely display the value of the word processor, and it adds little to the published literature on this subject. Nine of the 12 papers were already discussed in the previous review.

A D J WEBBORN
Medical Advisor Exercise on Prescription,
East Sussex, Brighton and Hove
Health Authority

- 1 Hillsdon M, Thorogood M. A systematic review of physical activity promotion strategies. *Br J Sports Med* 1996;30:84–89
- 2 Lewis BS, Lynch WD. The effect of physician advice on exercise behaviour. *Prev Med* 1993;22:110–21.
- 3 Logsdon DN, Lazaro CM, Meier RV. The feasibility of behavioural risk reduction in primary care. *Am J Prev Med* 1989;5:249–56.

- 4 Long BJ *et al.* A multisite field test of the acceptability of physical activity counseling in primary care: project PACE. *Am J Prev Med* 1996;12: 73-81.
- 5 Hillsdon M, Thorogood M, Anstiss T, Morris J. Randomised controlled trials of physical activity promotion in free living populations: a review. *J Epidemiol Community Health* 1995;49:448-53.

BOOK REVIEWS

Anatomy and Human Movement.

Structure and Function. 2nd ed. Edited by N Palastanga, D Field, and R Soames. (Pp 894; soft cover £40.00.) London: Butterworth Heinemann, 1994. ISBN 0 7506 0970 2

The second edition of *Anatomy and human movement* is a very comprehensive functional anatomy textbook. It would be a very useful book for physiotherapists and for anyone involved in sports medicine.

The clinical significance of the joints, particularly the biomechanical aspects, is simply and clearly laid out, emphasising the stresses on the joints during the various phases of the gait cycle. The section on the

nervous system is also clearly laid out, but there is no section on the sensory motor tracks, which I believe would be a useful addition.

Most of the diagrams are very clear but in the illustrations that include the segment of the spinal cord, the posterior ganglion (which distinguishes the posterior root) has been left out. The embryology is basic and so is the section on the thorax and the abdomen. The evolution of the various grips in the hand is included and the muscles involved in the various grips are clearly demonstrated.

This book is a must for anyone involved or interested in locomotion, physiotherapy, and sports medicine.

MOIRA O'BRIEN
Dublin

Management Strategies in Athletic Training,

by R Ray. (Pp 256; £26.00.) Human Kinetics (Europe) Ltd, 1994. ISBN 0 87322 582 1

It is always sad when one is asked to review a book which is well presented, comprehensively researched, but inappropriate to the market for which it is being reviewed. *Management strategies in athletic training* is sadly one such publication.

In his preface the author himself identifies the main audience as athletics trainers at the

various stages of their professional development. Sadly in the United Kingdom we do not as yet have this specialisation and until such time as we do the book and information within it have no real role to play other than providing interest and background for anyone wanting further information on this profession. It would also give an insight to some of the cultural and ethical differences experienced by athletics trainers practising within the USA. Likewise the case studies would help to give interested parties an insight into the administrative and legal problems that are encountered by athletics trainers within the National Athletic Trainers Association.

I wish that I could be more positive. It is a well presented textbook. The author is an eminently qualified academic practitioner and a "recognised" leader in the field of athletic training administration. Unfortunately, however, I have to give an opinion based on the suitability of the publication in our environment and culture. Sadly therefore I must conclude by saying that other than for general interest and background information I do not think that it is relative or appropriate to sports medicine practitioners outwith the United States of America.

GRAHAM N SMITH
Glasgow

American College of Sports Medicine, Annual Congress, 29 May - 1 June 1996, Cincinnati, Ohio

With the help of a grant from the Sports Council (International Affairs Unit), I attended the 43rd Annual Meeting of the American College of Sports Medicine at the Cincinnati Convention Centre, Cincinnati, Ohio. This is the major event in the calendar of ACSM, and attracted in excess of four and a half thousand delegates.

While many of the speakers were Americans, contributions from South Africa and Australia ensured that some balance of approach was achieved. What a pity that no Briton was invited to give any of the keynote presentations—surely there would be a case for funding one of our eminent sports medicine or sports science authorities to present the current state of British achievement in their own specialist areas. There are such fundamental differences in the provision of health care here compared to the USA that I am sure this suggestion would arouse considerable interest.

The overall standard of presentation was excellent, although certain contributions left me feeling slightly cheated at the lack of new material produced.

One of the lasting impressions was the sheer enormity of the sports medicine network in the USA. Contributions from every conceivable specialty and subspecialty encompassed the full breadth of sport and exercise medicine. At a clinical level one could not help but be impressed by, and considerably envious of, the quality of facilities and personnel available to

assess and treat the athletic population. The importance of the team approach—involving the athletic trainer, physician, surgeon, nutritionist, sports scientist, and other allied professionals—was evident from many joint presentations of clinical topics.

The importance of proprioceptive training in the rehabilitation process came up time and again—and numerous examples were given which will be included in future treatment protocols for my patients.

On many occasions speakers emphasised that we must communicate with the coaches—we cannot treat athletes without an intimate knowledge of the demands and technical requirements of their sport. Similarly, the importance of educating our patients, not only in areas which may prevent injury or optimise performance, but in settling realistic goals in the rehabilitation process, were covered by several speakers.

Certain individual presentations left a lasting impression:

Dr John Lombardo's image of driving a Ferrari down a long straight highway and the driver's attitude to being caught speeding was both challenging and informative as an illustration of athletes' temptation to using performance enhancing substances. The dual role of the doctor as an educator and a healer, contrasting with the role of the doping officer as an enforcer and punisher, called us all to question our attitudes and motives when dealing with athletes. The distinction between recreational