



FOR ONE MONTH'S RELIEF

PRESCRIBING INFORMATION FOR FELDENE* (PIROXICAM) GEL 112g: UK Indications: FELDENE Gel is effective in the treatment of osteoarthritis of superficial joints and acute musculoskeletal disorders. Dosage: For external use only. Occlusive dressings should not be used. Apply one gram of FELDENE Gel and rub into the affected site three to four times daily leaving no residual material on the skin. Therapy should be reviewed after 4 weeks. Contra-indications Precautions: Contra-indicated in patients who have previously shown a hypersensitivity to FELDENE Gel or piroxicam in any of its forms, aspirin or other non-steroidal anti-inflammatory agents. If local irritation develops, discontinue FELDENE Gel. Keep away from the eyes and mucosal surfaces. Do not apply to sites affected by open skin lesions,

dermatoses or infection. Use in children: Not recommended. Use in pregnancy and lactation: The safety of FELDENE Gel during pregnancy and lactation has not been established. Side- Effects: Mild or moderate local reactions at the application site. Systemic reactions occur infrequently including nausea and dyspepsia; rare reports of abdominal pain and gastritis and isolated reports of bronchospasm and dyspnoea. Legal Category: POM. Package Quantities and Basic NHS Cost: FELDENE Gel 112g tube-

£7.84 (PL 0057/0284). Further information on request. **Pfizer Limited**, Sandwich, Kent. †FELDENE Gel is the first topical NSAID in the UK with a licensed indication for osteoarthritis.





A Division of Pfizer Ltd

Medi

Orthopaedic Supports



Genumed PLUS

Genumedi® PLUS "I" (inferior position silicone horseshoe shape buttress)

and Genumedi® PLUS "S"

(superior position horseshoe)

is a seamless knitted two-way stretch knee support.

Available in seven sizes for correct fitting.

INDICATIONS

- Osgood-Schlatter (Adolescent Knee)
- Chondromalacia of the patella
- Patella Tendinitis (Jumper's Knee)
- Patella-femoral pain syndrome

Genumed *

multi-dimensional stretch elastic knee support.

Available in seven sizes for correct fitting.

INDICATIONS

- Effusions caused by degenerative joint diseases
- Post-operative and post-traumatic inflammations
- Chondromalacia of the patella
- Irritations of the knee joint
- Patella-femoral pain syndrome
- Symptoms associated with ligament instability
- Adolescent knee

medi UK Itd.

Fields Yard, Plough Lane Hereford HR4 0EL Phone: (01432) 351682

Fax: (01432) 342383

If your patient requires a high quality, cost effective Orthopaedic joint support, then please consult your **medi** and brochure. Should you not have a brochure then phone 01432 351682.

We shall be pleased to supply you with one.