

ORIGINAL ARTICLE

Are unilateral and bilateral patellar tendinopathy distinguished by differences in anthropometry, body composition, or muscle strength in elite female basketball players?

J E Gaida, J L Cook, S L Bass, S Austen, Z S Kiss

Br J Sports Med 2004;38:581–585. doi: 10.1136/bjism.2003.006015

Background: Overuse injury to the patellar tendon (patellar tendinopathy) is a major reason for interrupted training and competition for elite athletes. In both sexes, the prevalence of unilateral and bilateral tendinopathy has been shown to differ. It has been proposed that bilateral pathology may have a different aetiology from unilateral pathology. Investigation of risk factors that may be unique to unilateral and bilateral patellar tendinopathy in female athletes may reveal insights into the aetiology of this condition.

Objectives: To examine whether anthropometry, body composition, or muscle strength distinguished elite female basketball players with unilateral or bilateral patellar tendinopathy.

Methods: Body composition, anthropometry, and muscle strength were compared in elite female basketball players with unilateral ($n = 8$), bilateral ($n = 7$), or no ($n = 24$) patellar tendinopathy. Body composition was analysed using a dual energy x ray absorptiometer. Anthropometric measures were assessed using standard techniques. Knee extensor strength was measured at 180°/s using an isokinetic dynamometer. z scores were calculated for the unilateral and bilateral groups (using the no tendinopathy group as controls). z scores were tested against zero.

Results: The tibia length to stature ratio was approximately 1.3 (1.3 SDs above zero in both the affected and non-affected legs in the unilateral group ($p < 0.05$). The waist to hip ratio was 0.66 (0.78) SD above zero in the unilateral group ($p < 0.05$). In the unilateral group, leg lean to total lean ratio was 0.42 (0.55) SD above zero ($p < 0.07$), the trunk lean to total lean ratio was 0.63 (0.68) SD below zero ($p < 0.05$), and leg fat relative to total fat was 0.47 (0.65) SD below zero ($p < 0.09$). In the unilateral group, the leg with pathology was 0.78 (1.03) SD weaker during eccentric contractions ($p < 0.07$).

Conclusions: Unilateral patellar tendinopathy has identifiable risk factors whereas bilateral patellar tendinopathy may not. This suggests that the aetiology of these conditions may be different. However, interpretation must respect the limitation of small subject numbers.

See end of article for authors' affiliations

Correspondence to:
Dr Cook, Musculoskeletal Research Centre, La Trobe University, 3086 Victoria, Australia; j.cook@latrobe.edu.au

Accepted 16 June 2003

Overuse injury to the patellar tendon (patellar tendinopathy) is common among athletes (particularly at elite levels) and is a major reason for interrupted training and competition.^{1,2} It is notoriously difficult to treat and may result in untimely retirement from competitive sport.³ Despite being first described in 1973,⁴ little is known about the aetiology of this condition. Strength and anthropometric measures have been associated with symptomatic patellar tendinopathy in male athletes, although the findings are equivocal.^{5–11} Similar data in female athletes are scarce.

The prevalence of unilateral and bilateral patellar tendinopathy has been shown to differ between the sexes; bilateral tendinopathy is equally prevalent, but unilateral tendinopathy is twice as common in males as females.¹² This suggests that bilateral pathology may have a different aetiology from unilateral pathology. This is supported by evidence that female athletes with unilateral patellar tendinopathy had significantly less posterior leg flexibility than subjects with bilateral patellar tendinopathy and subjects with normal tendons.⁵

The reason for the different prevalence and presentation of unilateral and bilateral tendinopathy may be individual and inherent characteristics. The genetic profile is associated with severe collagen disease,¹³ and, although yet to be shown, collagen changes in some presentations of tendon disease

may also have a genetic link. In the Achilles tendon, the ABO blood grouping has been associated with pathology; subjects with bilateral tendon rupture had the highest frequency of type O blood (71%) followed by unilateral tendon involvement (53%); both were higher than the general population (31%).¹⁴

Investigation of risk factors that may be unique to unilateral and bilateral patellar tendinopathy in women may reveal insights into the aetiology of tendon pathology in these two groups. Thus we asked the following question: are elite female basketball players with unilateral or bilateral patellar tendinopathy distinguished by differences in anthropometry, body composition, or muscle strength?

METHODS

Two Women's National Basketball League (WNBL) and six Australian Basketball Association (ABA) clubs were invited to participate in this research. Subjects were considered suitable for participation if they were currently engaged in full training and match responsibilities, had not had knee surgery in the preceding six months, and had never had surgery on the patellar tendon. Current symptoms of patellar tendinopathy did not exclude subjects if they met the above criteria. The faculty human ethics committees at La Trobe University, Deakin University, and the Radiation Safety Unit

approved all procedures. All subjects gave written informed consent.

Subjects recorded symptoms (VISA score¹⁵) and details on training, injuries, surgery, contraceptive use, and menstrual history. Physical performance, body composition, and anthropometric values were then tested. Subsequently an ultrasound examination of both patellar tendons was performed by an experienced musculoskeletal radiologist.

Physical and anthropometric testing

All physical and anthropometric testing was conducted in one session at the human movement laboratories at Deakin University by the same investigator (JEG) and a trained anthropometrist (SA). Standing and sitting heights were measured using a Holtain wall stadiometer and sitting height table (British Indicators Ltd, Burgess Hill, West Sussex, UK). Weight was measured using a Seca electronic scale (Seca Ltd, Birmingham, UK). Segmental limb lengths were measured using a Harpenden anthropometer (British Indicators Ltd); coefficient of variability was $0.1 \pm 0.1\%$.¹⁶

Physical performance testing measured sit and reach, vertical jump, and isokinetic quadriceps strength. Before a standard warm up and stretch, sit and reach was determined using standard procedures. Right leg, left leg, and bilateral vertical jump were then tested using a Vertec (Sports Imports Inc, Columbus, Ohio, USA). The protocol allowed no run up or arm swing during take off. A countermovement to 90° of knee flexion was allowed for all jumps. Lastly, isokinetic strength of the quadriceps was tested using a Cybex dynamometer (Lumex Inc, Ronkonkoma, New York, USA). Five familiarisation trials followed by three recorded trials were performed at 180°/s. Output was recorded as average work over the three trials.

Body composition

Body composition was analysed using a dual energy x ray absorptiometer (DPX-L; Lunar Radiation Corporation, Madison, Wisconsin, USA). Total and regional fat and lean body mass were determined using a total body scan.

Ultrasound examination

Each subject had an ultrasound examination of both patellar tendons. The examinations were performed by an experienced radiologist with additional musculoskeletal training. For the detection of hypoechoic lesions, both the intertester ($\kappa = 1$)¹⁷ and test-retest reliability (40/40)¹⁸ has been documented as excellent for a radiologist with musculoskeletal training.

Both transverse and longitudinal scans were conducted using a high resolution linear array with a 10 or 12 MHz transducer. Tendons were designated pathological if a hypoechoic lesion was visible in both the longitudinal and transverse scans. The ultrasound results for both tendons were used to assign subjects to one of three naturally occurring groups: no tendinopathy (control), unilateral tendinopathy, or bilateral tendinopathy.

Data analysis

Data for these three groups were then entered into SPSS (SPSS Inc, Chicago, Illinois, USA). Analysis was conducted using the median score of three trials for anthropometric data¹⁶ and mean of three trials for all other data. Data for the unilateral and bilateral tendinopathy groups were then expressed as z scores using the mean and standard deviation of the corresponding score from the control group. z scores for each leg were calculated from the comparable leg—that is, left or right leg—in the controls. A one sample t test was used to determine if any z scores deviated from zero. Results are reported to be significant at $p < 0.05$, and trends are reported

where $p < 0.10$, as this allows the reader to determine both the clinical and further research significance of these findings.¹⁹ For clarity, the unilateral and bilateral patellar tendinopathy groups will be referred to as the “unilateral” and “bilateral” groups respectively.

RESULTS

Thirty nine subjects were recruited, 16 from the WNBL and 23 from the ABA. The prevalence of tendinopathy (unilateral and bilateral) did not differ between the two levels of competition. Ultrasound examination showed a unilateral hypoechoic lesion in eight athletes, bilateral lesions in seven athletes, and no lesion in 24 athletes. Subjects with one or two hypoechoic lesions did not differ from controls in age, menstrual history, height, or weight (table 1). Subjects with one or two hypoechoic lesions, however, had trained a mean (SD) of 2.6 (1.4) more hours a week than controls in the preceding one to six months and had lower VISA scores (12.1 (5.7)) than controls (both $p < 0.05$) (table 2).

There were no detectable differences between the bilateral group and controls in any of the measured variables. In contrast, subjects in the unilateral group were distinguished from the control group by differences in anthropometry, body composition, and strength. The tibia length to stature ratio was approximately 1.3 (1.3) SDs above zero in both the affected and non-affected legs in the unilateral group ($p < 0.05$) (fig 1). Although no differences in waist or hip girths were detected between the groups, the waist to hip ratio was 0.66 (0.78) SD above zero in the unilateral group ($p < 0.05$) (fig 2). The unilateral group leg lean to total lean ratio was 0.42 (0.55) SD above zero ($p < 0.07$). In contrast, trunk lean to total lean ratio was 0.63 (0.68) SD below zero ($p < 0.05$) (fig 3). The reverse was evident for fat mass: the leg fat relative to total fat was 0.47 (0.65) SD below zero ($p < 0.09$) (fig 3). In the unilateral group the leg with pathology was 0.78 (1.03) SD weaker during eccentric contractions ($p < 0.07$).

DISCUSSION

In this study of elite female basketball players, several anthropometric and body composition variables distinguished athletes with one hypoechoic tendon from controls. No measured variable distinguished athletes with two hypoechoic tendons from controls.

Subjects with unilateral patellar tendinopathy had a higher waist to hip ratio than controls. This indicates a larger abdominal fat distribution relative to gluteofemoral fat deposits.¹⁶ Human fat distribution is controlled by a complex interaction of hormones and is particularly influenced by the female sex hormones (oestrogen and progesterone), but the mechanisms whereby hormones control this fat distribution are unclear.²⁰

Hormone concentrations may also influence susceptibility to tendinopathy. Oestrogen has been reported to directly affect fibroblast proliferation and collagen synthesis,^{21–24} and menopause is associated with an increase in Achilles tendon ruptures, a finding that is not paralleled in men of an equivalent age.²⁵ This study did not examine hormone concentrations and thus this hypothesis cannot be substantiated. However, these data together support the notion that hormone concentrations that may affect connective tissue are characterised by a redistribution of body fat to the truncal region. Furthermore, combining the decrease in trunk lean to total lean ratio along with trends for leg lean and fat observed within the data suggest that subjects with unilateral tendinopathy have unique body compositions. Together these data indicate an android distribution of both fat and lean mass in subjects with unilateral patellar tendinopathy.

Table 1 Age, menstrual history, height, weight, anthropometric data, body composition data, and strength data of female basketball players with unilateral or bilateral patellar tendinopathy and controls

	Unilateral	Bilateral	Control
Number	8	7	24
Age (years)	20 (2)	21 (3)	21 (3)
Age of menarche (years)	14 (1)	14 (2)	14 (1)
Number of cycles (last 12 months)	12 (0.5)	12 (0.5)	12 (1.5)
OCP (consecutive months)	13 (25)	17 (26)	29 (36)
Height (cm)	178 (10)	178 (9)	176 (7)
Weight (kg)	73 (13)	74 (13)	74 (9)
Anthropometry			
Sitting height (cm)	91 (5)	91 (2)	92 (3)
Waist girth (cm)	75 (7)	74 (6)	75 (6)
Hip girth (cm)	99 (7)	102 (9)	102 (5)
Waist:hip ratio	0.76 (0.03)	0.73 (0.02)	0.74 (0.03)
Right femur (cm)	45 (4)	47 (5)	45 (3)
Left femur (cm)	45 (4)	47 (5)	44 (3)
Right tibia (cm)	41 (3)	40 (4)	39 (2)
Left tibia (cm)	41 (3)	40 (4)	39 (2)
Sit and reach (cm)	6 (13)	9 (14)	4 (8)
Body composition			
Lean (kg)			
Legs	18.3 (1.8)	17.1 (2.3)	17.8 (1.8)
Trunk	23.0 (2.3)	22.6 (1.9)	23.2 (2.7)
Total	49.9 (4.6)	48.0 (4.9)	49.1 (5.1)
Fat (kg)			
Legs	8.5 (4.2)	10.1 (3.8)	9.4 (2.7)
Trunk	8.8 (4.4)	9.2 (3.2)	9.0 (2.8)
Total	19.6 (9.4)	21.6 (7.1)	20.6 (6.0)
Strength			
Concentric (J)			
Right	153 (30)	138 (24)	143 (25)
Left	150 (35)	140 (20)	144 (21)
Eccentric (J)			
Right	-213 (51)	-201 (69)	-234 (46)
Left	-224 (71)	-230 (56)	-233 (48)
Vertical jump (cm)			
Two legged	48 (6)	47 (4)	46 (4)
Right leg	38 (5)	34 (3)	35 (3)
Left leg	37 (5)	35 (4)	35 (4)

Values are mean (SD).
OCP, Oral contraceptive pill.

Patellar tendinopathy appears to result from multiple risk factors. Often the presence of pathology has attributed been to differences in limb lengths leading to poor movement mechanics or biomechanical factors.²⁶⁻²⁷ We found that the players with unilateral patellar tendinopathy had a longer tibia length relative to stature. This has not previously been reported, but there is evidence that an increase in both tibial and femoral lengths are associated with overuse knee injuries.²⁸

The simplest explanation for longer tibiae in subjects with unilateral patellar tendinopathy is that altered lower limb biomechanics increases the stress on the patellar tendon and

leads to tendinopathy. This appears unlikely, however, as only one tendon was affected while both tibiae were longer than the control group. Furthermore, all subjects with unilateral tendinopathy played right handed while equal numbers of left and right tendons were affected. Tibial length and unilateral patellar tendinopathy may be parallel phenomena both arising from an underlying characteristic in the individual. The relation appears complex, as the increase in

Table 2 Training history and VISA score for female basketball players with unilateral or bilateral patellar tendinopathy and controls

	Tendinopathy		
	Unilateral (n = 8)	Bilateral (n = 7)	Control (n = 24)
Training (hours a week)			
Past month	10 (4)	12 (6)	9 (2)
Past 1-6 months	12 (5)	12 (5)	9 (3)*
Past 6-12 months	10 (4)	10 (5)	9 (5)
VISA score	83 (18)	72 (23)	90 (10)*

Values are mean (SD).
*Significantly different from patellar tendinopathy (unilateral and bilateral), p<0.05.




Figure 1 Femur to stature ratio and tibia to stature ratio z scores for the affected and unaffected leg in subjects with unilateral and bilateral patellar tendinopathy (mean and 95% confidence interval (CI)). *Significantly different from zero, p<0.05.




Figure 2 Waist girth, hip girth, and waist to hip ratio z scores in subjects with unilateral and bilateral patellar tendinopathy (mean and 95% confidence interval (CI)). *Significantly different from zero, $p < 0.05$.

length is specific to the tibia in relation to stature. To affect the length of long bones and influence tendon collagen and/or matrix morphology suggests a fundamental characteristic, which may include individual genetic or hormonal profile. This finding, however, remains unclear.

The knee extensor mechanism of the affected leg in the unilateral tendinopathy group showed a trend toward decreased eccentric strength at 180°/s. Three observations infer that this finding results from inhibition of eccentric contractions rather than muscular atrophy. Firstly, these subjects have more leg musculature than controls. Secondly, knee extensor strength does not differ from controls for concentric contractions. Thirdly, elite basketball players do not tend to develop one sided weakness as the ability to lead with or jump from either leg is a highly valued attribute. Decreased eccentric strength may initially protect the tendon from abusive stress levels (highest during eccentric contractions) but a decrease in controlled muscle strength may ultimately be detrimental to tendons.²⁹

Limitations of the study

A number of the topics in the discussion led to the suggestion that female sex hormones may influence the risk of pathology in these athletes. This notion is speculative as sex steroids were not measured in this study and this warrants further investigation. Furthermore, the study design was cross sectional with relatively small subject numbers. Notwithstanding this, a number of significant results were reported. These data, however, need to be confirmed using large sample sizes.

CONCLUSION

These data suggest that unilateral and bilateral patellar tendinopathy have unique aetiology. Several variables in the group with unilateral patellar tendinopathy differed significantly from controls, whereas no variable distinguished subjects with bilateral patellar tendinopathy from controls. In fact, those variables that reached significance in the unilateral group tended toward the opposite direction in the bilateral group. Although further research is required to validate these findings in larger samples, across sports, and in male subjects, these data suggest that unilateral and bilateral patellar tendinopathy warrant separate investigation.

ACKNOWLEDGEMENTS

We thank the players and coaches of the Bullen Boomers (WNBL), Dandenong Rangers (WNBL and ABA), Eltham Wildcats (ABA),




Figure 3 Regional fat to total fat ratio and regional lean to total lean ratio z scores for the legs and trunk of subjects with unilateral and bilateral patellar tendinopathy (mean and 95% confidence interval (CI)). *Significantly different from zero, $p < 0.05$; †Significantly different from zero, $p < 0.10$.

Hume City Broncos (ABA), Kilsyth Lady Cobras (ABA), Knox Raiders (ABA), and Nunawading Spectres (ABA).

Authors' affiliations

J E Gaida, J L Cook, Musculoskeletal Research Centre, School of Physiotherapy, La Trobe University, Bundoora, 3086, Australia
S L Bass, S Austen, Physical Activity and Nutrition Research Unit, School of Health Sciences, Deakin University, Burwood, 3125, Australia
Z S Kiss, Medical Imaging Australia, Melbourne, Australia

REFERENCES

- 1 **Jarvinen M**. Epidemiology of tendon injuries in sports. *Clin Sports Med* 1992;11:493–504.
- 2 **Hickey GJ**, Fricker PA, McDonald WA. Injuries of young elite female basketball players over a six-year period. *Clin J Sport Med* 1997;7:252–6.
- 3 **Cook JL**, Khan KM, Purdam CR. Conservative treatment of patellar tendinopathy. *Phys Ther Sport* 2001;2:54–65.
- 4 **Blazina ME**, Kerlan RK, Jobe FW, et al. Jumper's knee. *Orthop Clin North Am* 1973;4:665–78.
- 5 **Cook JL**, Kiss ZS, Kahn KM, et al. Anthropometry, physical performance, and ultrasound patellar tendon abnormality in elite junior basketball players: a cross sectional study. *Br J Sports Med* 2004;38:206–9.
- 6 **Ferretti A**. Epidemiology of jumper's knee. *Sports Med* 1986;3:289–95.
- 7 **Hopper DM**, Hopper JL, Elliott BC. Do selected kinanthropometric and performance variables predict injuries in female netball players? *J Sports Sci* 1995;13:213–22.
- 8 **Lian O**, Engebretsen L, Ovrebo RV, et al. Characteristics of the leg extensors in male volleyball players with jumper's knee. *Am J Sports Med* 1996;24:380–5.
- 9 **Richards DP**, Ajemian SV, Wiley JP, et al. Relation between ankle joint dynamics and patellar tendinopathy in elite volleyball players. *Clin J Sport Med* 2002;12:266–72.
- 10 **Richards DP**, Ajemian SV, Wiley JP, et al. Knee joint dynamics predict patellar tendinitis in elite volleyball players. *Am J Sports Med* 1996;24:676–83.
- 11 **Witvrouw E**, Bellemans J, Lysens R, et al. Intrinsic risk factors for the development of patellar tendinitis in an athletic population. A two year prospective study. *Am J Sports Med* 2001;29:190–5.
- 12 **Cook JL**, Khan KM, Harcourt PR, et al. Patellar tendon ultrasonography in asymptomatic active athletes reveals hypoechoic regions: a study of 320 tendons. Victorian Institute of Sport Tendon Study Group. *Clin J Sport Med* 1998;8:73–7.
- 13 **Cole WG**. Genetics of connective tissue disease. *Med J Aust* 1993;158:678–80.
- 14 **Jozsa L**, Barzo M, Balint JB. Correlations between the ABO blood group system and tendon rupture. *Magy Traumatol Orthop Helyreallito Seb* 1990;33:101–4.
- 15 **Visentini PJ**, Khan KM, Cook JL, et al. The VISA score: an index of severity of symptoms in patients with jumper's knee (patellar tendinosis). Victorian Institute of Sport Tendon Study Group. *J Sci Med Sport* 1998;1:22–8.
- 16 **Norton K**, Olds T. *Anthropometrica*. Sydney: University of New South Wales Press, 1996.
- 17 **Black J**. Patellar tendinopathy: the intertester reliability of ultrasound imaging [honours thesis]. Bundoora: La Trobe University, 2002.
- 18 **Khan KM**, Cook JL, Kiss ZS, et al. Patellar tendon ultrasonography and jumper's knee in female basketball players: a longitudinal study. *Clin J Sport Med* 1997;7:199–206.
- 19 **Sterne JA**, Davey Smith G. Sifting the evidence: what's wrong with significance tests? *BMJ* 2001;322:226–31.

- 20 **Bjorntorp P.** The regulation of adipose tissue distribution in humans. *Int J Obes Relat Metab Disord* 1996;**20**:291–302.
- 21 **Liu SH, al-Shaikh R, Panossian V, et al.** Primary immunolocalization of estrogen and progesterone target cells in the human anterior cruciate ligament. *J Orthop Res* 1996;**14**:526–33.
- 22 **Liu SH, Al-Shaikh RA, Panossian V, et al.** Estrogen affects the cellular metabolism of the anterior cruciate ligament. A potential explanation for female athletic injury. *Am J Sports Med* 1997;**25**:704–9.
- 23 **Yu WD, Liu SH, Hatch JD, et al.** Effect of estrogen on cellular metabolism of the human anterior cruciate ligament. *Clin Orthop* 1999;**366**:229–38.
- 24 **Yu WD, Panossian V, Hatch JD, et al.** Combined effects of estrogen and progesterone on the anterior cruciate ligament. *Clin Orthop* 2001;**383**:268–81.
- 25 **Maffulli N, Waterston SW, Squair J, et al.** Changing incidence of Achilles tendon rupture in Scotland: a 15-year study. *Clin J Sport Med* 1999;**9**:157–60.
- 26 **Kujala UM, Osterman K, Kvist M, et al.** Factors predisposing to patellar chondropathy and patellar apicitis in athletes. *Int Orthop* 1986;**10**:195–200.
- 27 **Shambaugh JP, Klein A, Herbert JH.** Structural measures as predictors of injury basketball players. *Med Sci Sports Exerc* 1991;**23**:522–7.
- 28 **Dahlstrom S, Kujala UM.** Anthropometry and knee exertion injuries incurred in a physical training program. *J Sports Med Phys Fitness* 1990;**30**:190–3.
- 29 **Cook JL, Khan KM, Maffulli N, et al.** Overuse tendinosis, not tendinitis. Part 2. Applying the new approach to patellar tendinopathy. *Phys Sportsmed* 2000;**28**:31–2.

ELECTRONIC PAGES.....

BJSM Online case reports: <http://bjsm.bmjournals.com/>

The following electronic only articles are published in conjunction with this issue of *BJSM*.

The validity of capillary blood sampling in the determination of human growth hormone concentration during exercise in adult men

R J Godfrey, G Whyte, J McCarthy, A Nevill, T Head

Background: Studies measuring human growth hormone (hGH) in blood during exercise have mainly used venous sampling. The invasive nature of this procedure makes evaluation of hGH impossible in various exercise environments.

Objective: To determine whether capillary sampling could offer an alternative sampling method.

Methods: Capillary and venous blood samples were collected for determination of hGH at the end of each exercise stage during an incremental exercise test in 16 male club level competitive cyclists (mean (SD) age 30.8 (8.0) years, body mass 72.2 (7.1) kg, body fat 12.9 (3.5)%, peak oxygen consumption 4.18 (0.46) l·min⁻¹). Linear regression, from a plot of venous v capillary blood hGH concentration, showed a correlation coefficient of $r = 0.986$ ($p < 0.001$). When geometric means and log transformations were used, a coefficient of variation of 14.2% was demonstrated between venous and capillary flow for hGH concentration. The mean ratio limits of agreement were 0.62 (1.72)—that is, 95% of the ratios were contained between 0.36 and 1.07, with a mean of 0.62.

Conclusions: Capillary blood sampling is an acceptable alternative to venous sampling for determining hGH concentration during rest and exercise. Sample sites should not be used interchangeably: one site should be chosen and its use standardised.

(*Br J Sports Med* 2004;**38**:e27) <http://bjsm.bmjournals.com/cgi/content/full/38/5/e27>

Orbital emphysema following nose blowing as a sequel of a snowboard related head injury

Y Taguchi, Y Sakakibara, K Uchida, H Kishi

A case of orbital emphysema as a sequel of a snowboard related head injury is reported. It is believed that a fracture of the medial orbital wall was caused by the increased intraorbital pressure when the patient hit his forehead on the snowy ground, allowing air to enter the orbit when he blew his nose. Wearing goggles may prevent this type of sports related injury.

(*Br J Sports Med* 2004;**38**:e28) <http://bjsm.bmjournals.com/cgi/content/full/38/5/e28>

Self reported injury patterns among competitive curlers in the United States: a preliminary investigation into the epidemiology of curling injuries

J C Reeser, R L Berg

Objective: To investigate the injury patterns among competitive.

Methods: Participants at two curling championship events were asked to complete injury history questionnaires.

Results: 76 curlers (39%) participated; 79% of these reported curling related musculoskeletal pain, most commonly involving the knee (54%), back (33%), and shoulder (20%). Sweeping and delivering the stone were most likely to provoke symptoms. Time loss injuries were estimated to occur at a rate of 2 per 1000 athlete exposures.

Conclusions: Curling appears to be a relatively safe winter sport. Prospective studies are needed to confirm these preliminary findings and to further define the risk factors for curling related injuries.

(*Br J Sports Med* 2004;**38**:e29) <http://bjsm.bmjournals.com/cgi/content/full/38/5/e29>