

Warm up

## The power of placebo

P McCrory

Just when the silly season was starting to get on top of us, some welcome relief in the form of good solid science arrived on my desk. A study published in the *BMJ* looked at the therapeutic effect of wearing magnetic bracelets for pain control in osteoarthritis.<sup>1</sup>

I must admit my personal bias here. When I was the team doctor for one of our professional football clubs, we used to be inundated with salesmen pushing their magnetic wares. Magnetic bandages, pillows, shoe inserts, wrist bands, knee bands, and who knows what else. It seems the only limitation in their product range was their own imagination. Their strategy of leaving piles of their products around the football club was in the hope that they could then claim endorsement by some gullible celebrity athlete who would pick up a free sample. Not surprisingly, the footballers loved them. The combination of a free sample and some hocus science was an absolute winner. In fact, the more magnetic things they had attached to their bodies the better. It is like the old adage regarding footballers—if someone came along with a pill for

immortality, a footballer would take two of them.

You may well pooh-pooh this sales tactic but let me remind you that in 1999, the worldwide sales of magnetic products was estimated at \$5 billion USD (£2.6 billion, €3.8 billion).<sup>2</sup>

So you can understand my obvious satisfaction when I read the study of magnetic bracelets.<sup>1</sup> In a prospective RCT of 194 subjects with osteoarthritis of the hip and knee, three groups were studied for their pain levels wearing a standard strength bipolar magnetic bracelet, a weak magnetic bracelet, and a dummy non-magnetic bracelet. After 12 weeks of wearing the bracelets, all three groups reported less pain and the two groups with the magnetic bracelets reported less pain than the placebo group (note that it is impossible to blind the subjects to their use of a non-magnetic bracelet given that it doesn't stick to metal objects!) Despite this finding, there was no statistically significant difference in pain levels between the groups.

Previous studies in a variety of medical conditions have shown variable results from magnetic therapy with a

few studies reporting a benefit in pain reduction<sup>2-3</sup> whereas larger most studies have reported no beneficial effect from magnetic therapy.<sup>4-6</sup>

Interestingly the effect of the magnets on pain, although small, was similar to that seen with other traditional first line osteoarthritis treatments such as topical non-steroidal creams<sup>7</sup> and exercise therapy.<sup>8</sup> It is reassuring to know that in this era of evidence-based medicine that placebos still work.

*Br J Sports Med* 2005;**39**:125

### REFERENCES

- 1 Harlow T, Greaves C, White A, et al. Randomized controlled trial of magnetic bracelets for relieving pain in osteoarthritis of the hip and knee. *BMJ* 2004;**329**:1450-4.
- 2 Weintraub M. Magnetic bio-stimulation in painful diabetic peripheral neuropathy: a novel interventional randomized double blind double placebo cross over study. *Am J Pain Management* 1999;**9**:8-17.
- 3 Brown CS, Ling FW, Wan JY, et al. Efficacy of static magnetic field therapy on chronic pelvic pain: a double blind pilot study. *Am J Obstet Gynecol* 2002;**187**:1581-7.
- 4 Collocott E, Zimmerman J, White P, et al. Bipolar permanent magnets for the treatment of chronic low back pain. *JAMA* 2000;**283**:1322-5.
- 5 Caselli M, Clark N, Lazarus S, et al. Evaluation of magnetic foil and PPT insoles in the treatment of heel pain. *J Am Podiatr Med Assoc* 1997;**87**:11-6.
- 6 Winemiller M, Billow R, Lasowski E, et al. Effect of magnetic vs sham-magnetic insoles on plantar heel pain: a randomised controlled trial. *JAMA* 2003;**290**:45-56.
- 7 Bookman A, Williams K, Shainhouse J. Effect of a topical diclofenac solution for relieving symptoms of primary osteoarthritis of the knee: a randomized controlled trial. *Can Med Assoc J* 2004;**171**:333-8.
- 8 O'Reilly S, Muir K, Doherty M. Effectiveness of home exercise on pain and disability from osteoarthritis of the knee: a randomized controlled trial. *Ann Rheum Dis* 1999;**58**:15-9.

Warm up

## "Tomber dans les pommes" – can head injury cause brain damage?

P McCrory

Those readers looking for some amusing leisure time reading need look no further than the December issue of the *Canadian Medical Association Journal*. The enterprising authors—Antoine Cyr (age 5 years) and his brother Louise-Oliver (age 7 years) in conjunction with their father—analysed the adventures of Tintin in an attempt to discover why the Tintin grew no

taller from 1929 to 1975, never had a girlfriend, and never needed to shave.<sup>1</sup>

After an exhaustive review of the published literature, episodes of head injury were identified by the authors and the injury severity determined by the duration of loss of consciousness (as measured by the number of frames before Tintin returned to normal activ-

ity) and the number of stars revolving above Tintin's head following injury.

Rather surprisingly a total of 50 episodes of head injury were identified (43 were AAN grade 3 concussion injuries) in Tintin's career. The mean length of LOC was 7.5 frames. The number of concussive events per adventure ranged from two to six, and most were blunt object impacts. Despite the absence of formal neuroimaging or neuro-endocrine assessment, the authors postulated that the repeated head trauma resulted in hypogonadotropic hypogonadism and growth hormone deficiency, which in turn would explain his delayed statural growth, delayed onset of puberty, and lack of libido.

This is the first report of such a syndrome occurring following concussion. Pituitary and hypothalamic dysfunction is seen not uncommonly following severe head trauma or in the

setting of raised intracranial pressure after head injury but to date such a phenomenon has not been reported with concussive injury. As a result one would have to speculate that Tintin was in fact suffering from primary hypothalamic dysfunction and his clumsiness and repeated head trauma was incidental. The occurrence of repeated concussive injuries in athletes has been documented previously and thought to reflect the individuals risk taking behaviour rather than an intrinsic potential for injury<sup>2</sup>. In Tintin's case, annual neuropsychological screening may be advisable to detect and hopefully prevent any long term cognitive deterioration.

It has been suggested by some authors<sup>3,4</sup> that repeated concussive injuries may be a risk factor for "second impact syndrome"; however, this has been questioned due to the limited evidence for its existence.<sup>5</sup> In Tintin's case, fear of this putative complication

seems to have not been a management concern in his return to normal activity.

Some anecdotal return to play guidelines<sup>6,7</sup> would recommend immediate retirement or termination of the current adventure given that Tintin often experiences more than three concussive episodes in a short space of time. However, the Vienna guidelines<sup>8</sup> recommend no mandatory exclusion period but individualised assessment of recovery—a strategy that Tintin appears to have followed in preference. If these guidelines were strictly adhered to, however, then he would have to terminate the adventure following the initial injury and Snowy the dog or Captain Haddock may have to play a more central role in the story outcome.

I can only suggest that Tintin play close attention to the forthcoming Prague guidelines in view of his propensity of concussive injury in order to avoid long term problems.

*Br J Sports Med* 2005;**39**:125–126

## REFERENCES

- 1 **Cyr A**, Cyr L, Cyr C. Acquired growth hormone deficiency and hypogonadotropic hypogonadism in a subject with repeated head trauma. *Can Med Assoc J* 2004;**171**:1433–4.
- 2 **Gerberich SG**, Priest JD, Boen JR, Straub CP, Maxwell RE. Concussion incidences and severity in secondary school varsity football players. *Am J Public Health* 1983;**73**:1370–5.
- 3 **Cantu RC**. Second-impact syndrome. *Clin Sports Med* 1998;**17**(1):37–44.
- 4 **Kelly JP**, Nichols JS, Filley CM, *et al*. Concussion in sports. Guidelines for the prevention of catastrophic outcome. *JAMA* 1991;**266**:2867–9.
- 5 **McCroary PR**, Berkovic SF. Second impact syndrome. *Neurology* 1998;**50**:677–83.
- 6 **Cantu RC**. Guidelines for return to contact sports after cerebral concussion. *Phys Sportsmed* 1986;**14**:75–83.
- 7 **Kelly J**, Rosenberg J. Diagnosis and management of concussion in sports. *Neurology* 1997;**48**:575–80.
- 8 **Aubry M**, Cantu R, Dvorak J, *et al*. Summary and agreement statement of the first International Conference on Concussion in Sport, Vienna 2001. *Physician and Sports Medicine* 2002;**30**:57–62.