



Foot orthoses and gait: a systematic review and meta-analysis of literature pertaining to potential mechanisms

Kathryn Mills,^{1,2} Peter Blanch,² Andrew R Chapman,¹⁻³ Thomas G McPoil,⁴ Bill Vicenzino¹

¹School of Health and Rehabilitation Sciences, University of Queensland, Brisbane, Australia

²Australian Institute of Sport, Canberra, Australia

³McGill University, Montreal, Canada

⁴Northern Arizona University, Flagstaff, Arizona, USA

Correspondence to

Professor Bill Vicenzino, School of Health and Rehabilitation Sciences, Division of Physiotherapy, The University of Queensland, Brisbane QLD 4072, Australia; b.vicenzino@uq.edu.au

Accepted 4 November 2009

ABSTRACT

This article systematically reviews the available literature to improve our understanding of the physiological basis for orthoses under the kinematic, shock attenuation and neuromotor control paradigms. The propositions made under these three paradigms have not been systematically reviewed collectively, and as such, there is no single-point synthesis of this clinically relevant body of evidence and somewhat disparate findings. Our comprehensive search strategy yielded 22 papers. Under each paradigm, the role of orthoses with different design features including combinations of posting, moulding and density was analysed. Where possible, data have been pooled to provide an increased level of confidence in findings. The main findings in the kinematic paradigm were that posted non-moulded orthoses systematically reduced peak rearfoot eversion (2.12° (95% CI 0.72 to 3.53)) and tibial internal rotation (1.33° (0.12 to 2.53)) in non-injured cohorts. In the shock attenuation paradigm, it was found that non-posted moulded and posted moulded orthoses produced large reductions in loading rate and vertical impact force when compared with a control and to a posted non-moulded orthosis. The neuromotor control paradigm seems to be the least conclusive in its outcome. Based on our review, this paper concludes with rudimentary guidelines for the prescription of orthosis, that sports medicine practitioners may use in their clinical decision-making process. The need for further research focusing on the role of injury, particularly in neuromotor control modification and long-term adaptation to orthoses, was highlighted.

Inshoe foot orthoses are frequently used by clinicians¹⁻³ in the management of overuse injuries. The Australian Podiatry Council and American College of Foot and Ankle Orthopedics and Medicine³ define “an orthosis as an appliance to support, align, correct deformity or motion of parts of the body.”¹ The conventional kinematic paradigm, on which these definitions are founded, is based on the hypothesis that abnormal pronation of the subtalar joint contributes to lower limb injuries and that orthoses normalise pronation and subsequent coupled movements (eg, internal tibial rotation).⁴⁻⁶ However, this has been questioned.⁷⁻¹¹

In addition to the kinematic paradigm, two other major paradigms have been proposed^{8,9,12,13}; which are essentially the shock attenuation and neuromotor control paradigms. The former is based on the concept that the magnitude of force during impact

is a major contributor to overuse injuries¹⁴⁻¹⁶ and orthoses are proposed to reduce impact force by acting as a cushioning interface between the ground and foot. More recently, the neuromotor control paradigm has been proposed, whereby an orthosis may optimise performance and minimise muscle activity and fatigue by providing input through the sole of the foot.^{8,9,17}

A source of confusion for both the researcher and clinician is the array of materials with various properties (type, density or hardness/firmness) that are either custom moulded or prefabricated into various shapes, which can be further customised by the addition of posting or wedging so as to tilt the device from the horizontal. A systematic review of the literature is timely to provide a critically evaluative synthesis of the physiological basis for orthosis therapy during gait under the kinematic, shock attenuation and neuromotor control paradigms. There is no other single source of evidence of the data synthesised from these paradigms, although there are isolated systematic reviews without meta-analyses available.^{8,12,18-21} This will assist clinicians in their prescription and fitting of orthoses and highlight areas for future research.

METHODOLOGY

Search strategy

We undertook a comprehensive, sensitive literature search strategy of Sportsdiscus, Medline, Cinahl, PubMed, Cochrane and Pedro databases from 1971 to September 2008 (fig. 1). Keywords used in the search strategy focused on the three identified paradigms: “ortho*, insole, shoe, foot, electromyography (EMG), muscle activity, biomechanics, kinetic, kinematic, shock attenuation, shock absorption, overuse injur*, leg, lower limb,” with no language restriction. Reference lists of reviews in similar topics and papers that met the inclusion criteria were hand searched (K.M.). Titles and (where necessary) abstracts retrieved by initial search were screened (K.M.), with only clinical trials meeting initial criteria considered for further review.

Inclusion and exclusion criteria

Included studies focused on the mechanism of action, rather than efficacy. Excluded were cohorts with neurological (eg, cerebral palsy), systemic (eg, diabetes and rheumatoid arthritis) and degenerative (eg, osteoarthritis) conditions, because these may complicate the analysis of gait.




Figure 1 Search strategy; *, three studies categorised in both kinematic paradigm and shock attenuation; one study categorised in both kinematic and neuromotor control paradigm.

For each paradigm, papers examining tasks other than gait were excluded because the kinematics, kinetics, muscle activity and shock attenuation in activities such as landing, step-ups, single-leg squats and balance assessments are too dissimilar to gait and often too heterogeneous for pooling.^{22–23} Papers studying three-dimensional kinematics were only included because with two-dimensional analysis, movement in the frontal plane is strongly affected by the alignment of the foot in the transverse plane.^{4,24–26}

Quality assessment

Since there is no validated quality assessment tool suitable for the repeated-measures, laboratory-based study designs included in our review, we adapted the Quality Index,²⁷ which is purported to be superior because it encompasses a profile of scores for rating: reporting, internal validity, power and external validity.²⁷ However, we only used relevant items, such as the reporting items, external validity items 11–13, internal validity (bias) items 14–16 and 18–20 and internal validity (confounding) items 26 and 27. A score for participant characteristics (item 3) was only recorded if studies described participants' injury type, physical activity levels and foot posture. Randomisation (item 23), as described in the Quality Index, was not applicable to all studies (eg, within- participants), so we quarantined it from the overall score and modified it, awarding studies a point if the order of intervention(s) and control were randomised. The maximum score obtainable was 28 from the index.

Data synthesis

Quantitative data synthesis was conducted using Cochrane Review Manager (V.5) with data extracted directly from the papers, and when not available, we attempted to contact the

authors. Mean difference between orthosis and comparator conditions and its 95% confidence interval (CI) was calculated. CI containing “0” represents a null effect. Estimates of the treatment effect are also provided in the form of an effect size (ES; difference in mean scores divided by pooled SD),²⁸ and classified as trivial (0–0.2), small (0.2–0.6), moderate (0.6–1.2) and large (>1.2),²⁸ thereby allowing a common metric across all measures.

RESULTS

Search results

Twenty-two papers that studied 30 different designs of orthoses on 34 kinematic and 18 kinetic variables were included (table 1). Analysis only included kinematic or kinetic variables that were investigated by more than three papers.

We categorised orthoses into three categories: (1) posted non-moulded, which refers to orthoses that were not contoured to the participant's foot (eg, flat) but with added posting; (2) non-posted moulded, which refers to orthoses that were custom made or contoured to the individual but with no posting; and (3) posted moulded that had both custom-contouring and additional posting. Density was not included in this categorisation, nor orthoses that had an irregular surface (several inbuilt raised areas), but were reported separately.

Studies were further categorised with regard to injury status of the cohorts: (1) no history of injury; (2) history, where participants were injured before orthosis prescription but were improving or asymptomatic during the study; and (3) currently injured. In addition, gait was divided into walking and jogging. Data pooling was conducted when investigations were the same across orthosis design, injury status and gait (ie, walking or jogging). Comparisons were also made either between an orthosis and a control (shoe, running sandal, flat ethylene-vinyl acetate (EVA) insert) or between two orthoses with different designs (posting; moulding; density; location of posting, eg, anterior, lateral, posterior, medial; height of posting, eg, inverted, 4°, etc).

Quality

Quality index scores ranged from 17 to 24, of a possible 28 (mean 20.4; table 2). Studies scored similarly in their reporting styles. Sixteen of the 22 papers presented sufficient data to extract and calculate point estimates of effect (eg, mean and SD/error or ES). Only eight studies attempted to describe adverse or longer-term effects of wearing orthoses. For internal and external validity, all studies performed poorly in three items: (1) generalisability of the sample (item 12), (2) blinding participants (item 14), and (3) blinding assessors (item 15). Thirteen papers randomised the order of intervention(s) and control.

Kinematic effects on the foot and shank

Fifteen papers presented kinematic and kinetic outcomes, studying 29 different orthosis designs, with kinematic variables of rearfoot eversion, tibial internal rotation, and rearfoot eversion velocity (tables 3–5) and kinetic variables of maximum ankle inversion moment and maximum knee external rotation moment (table 6).

Orthoses versus control

Rearfoot eversion

We found 38 comparisons of an orthosis versus a control. Of these, 28 involved orthoses of various posting and moulding designs. A further four comparisons investigated a semicustom orthosis described as a “mould-of best-fit,”²⁹ five comparisons involved different density materials and one comparison

Table 1 Studies included in the review

Author	Activity	N-total (sex)	Injury history*	Previous orthosis experience (mo)	Paradigm	Intervention	Review category	Control	Comments
Branthwaite <i>et al</i> ³²	Walking	9 M	1	0	Kinematic	(a) Biplanar: tilts calcaneus in frontal and sagittal planes; (b) Cobra: feature and arch profile and heel cup	Posted non-moulded, Posted non-moulded	Running sandal	Insufficient data to calculate CI
Butler <i>et al</i> ⁴¹	Jogging	15	1	0	Shock attenuation	(a) Suborthlene: rigid; (b) EVA foam: soft	Posted non-moulded, Posted non-moulded	Shoe	Heel counter removed
Eng and Pierrynowski ³⁸	Walking and jogging	10 F	3	See comments	Kinematic	Flat Spenco insole postedmedially in rearfoot and forefoot with rubber wedges	Posted non-moulded	Shoe	Participants had previously been prescribed orthoses by the same physical therapist and physician, time period not specified
MacLean <i>et al</i> ³⁶	Jogging	12 F	2	Not stated	Kinematic, shock attenuation	Semirigid thermoplastic functional foot orthosis intrinsically posted to calcaneal vertical and inverted an addition 5°. Featured added EVA rearfoot stabiliser and EVA top cover	Posted moulded (inverted)	Shoe	Measures taken initially then post 6-weeks of wearing orthosis
MacLean <i>et al</i> ³⁷	Jogging	15 F	1	0	Kinematic	Cast to calcaneal vertical and intrinsically posted an additional 5°. Extrinsic rearfoot post and EVA cushioning top cover	Posted moulded	Shoe	Authors state effect sizes (ES) but do not provide enough data to calculate mean difference and 95% CI
McCulloch <i>et al</i> ⁵	Walking at 2 speeds	10 (5 M, 5 F)	3	See comments	Kinematic	Rigid and semirigid posted to correct "biomechanical dysfunction"	Posted moulded	Shoe	Reported as "maximum pronation." However, methodology describes rear-foot eversion
Miller <i>et al</i> ⁴²	Walking	25 (13 M, 12 F)	1	Not stated	Shock attenuation	Custom plastic polymer orthoses with medial firm heel post	Posted moulded	Shoe	Not enough data to calculate mean differences and 95% CI
Mündermann <i>et al</i> ³⁰	Jogging	21 (9 M, 12 F)	1	0	Kinematic, shock attenuation	(a) EVA bottom layer with extrinsic 6 mm rearfoot and forefoot post; (b) neutral polypropylene shell; (c) polypropylene shell with extrinsic EVA 6 mm	Posted non-moulded, non-posted moulded, posted moulded	Flat EVA insert in running sandal	Ran for 2 weeks in running sandal before data collection
Mündermann <i>et al</i> ⁴⁶	Jogging	20 (9 M, 12 F)	1	Not stated	Neuromotor control	(a) Top layer consisted of 3-mm Spenco. Bottom layer consisted of full-length 6-mm EVA wedge; (b) custom polypropylene shell with no extrinsic posting; (c) custom polypropylene shell with 6-mm extrinsic EVA post added to medial rearfoot and forefoot areas	Posted non-moulded, non-posted moulded, posted moulded	Flat EVA insert in running sandal	Ran for 2 weeks in running sandal before data collection
Murley and Bird ⁴⁷	Walking	15 (7 M, 10 F)	1	Not stated	Neuromotor control	Custom-made rigid with 15° post	Posted moulded	Shoe	Orthosis worn for 4 weeks before data collection

Continued

Table 1 Continued

Author	Activity	N-total (sex)	Injury history*	Previous orthosis experience (mo)	Paradigm	Intervention	Review category	Control	Comments
Nawoczenski and Ludewig ⁴⁴	Jogging	12 (6 M, 6 F)	3	0	Neuromotor control	Custom semirigid polypropylene with extrinsic rearfoot and intrinsic forefoot post	Posted moulded	Running sandal	Orthosis worn for 3–4 weeks before data collection. Authors do not provide sufficient data to calculate ES.
Nester <i>et al</i> ⁴⁰	108 steps/min	15 (8 M, 7 F)	1	Not stated	Kinematic, shock attenuation	High-density EVA with medium density 10° post placed medially laterally	Posted non-moulded	Shoe	Data presented as graphs only
Nigg <i>et al</i> ³¹	Jogging	15 M	1	0	Kinematic	Prefabricated rigid EVA with 4.5-mm medial or lateral posts: (a) full-length medial; (b) full-length lateral; (c) 0.5 length medial; (d) 0.5 length lateral	Posted non-moulded	Manufacturer's insert in shoe	Insufficient data to calculate ES
Nigg <i>et al</i> ³⁴	Jogging	12 M	1	0	Kinematic	Bilayer orthoses (moulding not specified) made from polyurethane uppers and polyethylene or EVA lower layer: (a) Medium upper (Shore C65), soft EVA lower (shore C75); (b) soft upper (Shore C50), soft polyethylene lower (Shore C78); (c) soft upper (Shore C50), hard lower (Shore C84); (d) hard upper (Shore C70), soft polyethylene lower (Shore C78); (e) hard upper (Shore C70), hard lower (Shore C84)	Density	Shoe	Heel counter removed. Not enough data provided to calculate mean difference or CI
Nigg <i>et al</i> ⁴³	Jogging	16 M	1	Not stated	Shock attenuation	Commercially available viscoelastics orthoses with Shore values: (a) 26 (+stabiliser); (b) 28 (+stabiliser); (c) 29 (+stabiliser); (d) 34 (+stabiliser)	Density	Shoe with manufacturer's insert ± horse shoe-shaped rearfoot stabiliser	When compared with shoe with rearfoot stabiliser, orthosis condition also had rearfoot stabiliser.
Stackhouse <i>et al</i> ³⁹	Jogging	15	1	0	Kinematic	Semirigid functional orthotic devises fabricated from suborthelene with neoprene covers with 6° varus posting	Posted moulded	Shoe	2 weeks adjustment period before data collection. Heel counter removed. Insufficient data to calculate CI and ES
Stacoff <i>et al</i> ³³	Walking	8 M	3	0	Kinematic, neuromotor control	(a) Bilayer orthosis (hard lower, soft upper) posted supporting the calcaneus at the sustentaculum tali; (b) hard density, supporting the foot medially and laterally; (c) designed with inbuilt 4 raises (3 mm, 2 mm) at midfoot, metatarsal head and toes	Posted non-moulded, posted moulded, irregular surface	Running sandal	Comparisons made between orthoses and first control trials (start of testing period)

Continued

Table 1 Continued

Author	Activity	N-total (sex)	Injury history*	Previous orthosis experience (mo)	Paradigm	Intervention	Review category	Control	Comments
Tomaro and Burdett ⁴⁵	Walking	10 (3 M, 7 F)	2	>6	Neuromotor control	Sporthotic devices, modified to individuals' subtalar neutral	Posted moulded	Shoe	
Williams <i>et al</i> ⁹⁵	Jogging	11 (5 M, 6 F)	2	Chronic	Kinematic	Custom-made rigid. Internally posted to the forefoot deformity and externally posted to: (a) 4°; (b) 15° to 30°	Posted moulded, posted moulded (inverted)	Shoe	
Zifchock and Davis ²⁹	Walking	37 (17 M, 20 F)	1	0	Kinematic	Semigraphite with vinyl covers: (a) custom: made from plaster casts of individuals' foot; (b) semicustom: mould-of-best-fit chosen from range available	Posted moulded, posted moulded	Shoe	Divided into high and low arch based on arch height index

*Injury history: 1, no history of injury; 2, history of injury; 3, current injury. EVA, ethylene-vinyl acetate; M, male; F, female.

involved an orthosis with irregular surface. Nineteen comparisons measured peak rearfoot eversion and 19 measured rearfoot eversion excursion (table 3). Results of four papers detailing 11 comparisons did not provide enough data to calculate point estimates of effect.

The effect of orthoses on peak rearfoot eversion seems dependent upon design. For posted non-moulded orthoses, pooled data from two studies of participants with no history of injury revealed a 2.12° (95% CI 0.72 to 3.53) reduction in peak rearfoot eversion during jogging (fig. 2).^{11, 30} This finding is in line with the overall tendency for there to be a reduction in peak rearfoot eversion in four non-pooled comparisons during jogging^{11 30 31} and two of the three that studied walking.³²

For posted moulded orthoses, pooled data from two studies^{5 33} of currently injured participants revealed a 1.95° (0.1 to 3.79) reduction in peak rearfoot eversion. This finding is in line with two of the three comparisons involving currently injured cohorts which reported moderate (non-significant, eg, CI contained 0) effects in favour of the orthoses.^{5 33}

Of non-pooled data, only one comparison reported a statistically significant effect. A posted non-moulded orthosis produced a 2.3° (0.78 to 3.82) reduction of moderate effect (ES 0.92).³⁰ One comparison evaluated the irregular surfaced orthosis (ES 0.28) and another made five comparisons of different material density, where authors did not provide enough information for point estimates of effect, showing small non-systematic effects (table 3).^{33 34}

With regard to rearfoot eversion excursion, two comparisons examined the effect of inverted orthoses in cohorts with a history of injury. One study evaluated an inverted orthosis that was posted between 15° and 30° (often called a Blake orthosis),³⁵ the other inverted the posting an additional 5° from neutral.³⁶ In both studies, the individuals had been wearing their inverted orthoses for a minimum of 6 weeks. Pooled data revealed that the inverted orthoses had no effect (0.21° (-1.81 to 2.23)) on rearfoot eversion excursion. This finding contrasts that of MacLean *et al*,³⁷ who examined a pragmatic prescription of their custom-moulded device (±posting) and qualitatively reported a significant but small (ES 0.32) reduction in rearfoot eversion excursion between 15% and 50% of stance. The lack of any substantial effect is in line with all other comparisons regardless of orthosis'

design or injury status. Eng and Pierrynowski³⁸ reported near-significant moderate reductions in the midstance phase of walking (ES 0.82) and contact phase of jogging (ES 0.73) in 10 adolescents with diagnosed patellofemoral pain syndrome.

Tibial internal rotation

There were 24 comparisons between orthoses and a control involving tibial internal rotation, four made during walking. Eighteen provided enough information to calculate point estimates of effect and confidence intervals (table 4). The majority of comparisons were based on different posting and moulding designs, one used an orthosis with irregular surface and five comparisons were made between a control and orthoses of different densities.

Three sets of pooling were possible. Pooling from two comparisons^{11 30} involving participants with no history of injury wearing posted non-moulded orthoses showed a decrease of 1.33° (0.13 to 2.53) in tibial internal rotation when jogging (fig. 2). Likewise, pooled data from two comparisons^{33 38} involving currently injured participants, also wearing posted non-moulded orthoses, found a reduction of 1.66° (0.2 to 3.13) in tibial internal rotation after touchdown during walking gait (fig. 2). Pooling of two comparisons involved cohorts with a history of injury who had worn their inverted posted moulded orthoses for a minimum of 6 weeks.^{35 36} Wearing the inverted orthoses did not change tibial rotation during jogging (2.22° (-0.83 to 5.26); fig. 2).

From data that were not pooled, Stacoff *et al*¹¹ found a posted non-moulded orthosis, consisting of a medial post placed under the calcaneus added to a prefabricated orthosis, decreased tibial internal rotation by 1.59° (0.21 to 2.97) when jogging. Eng and Pierrynowski³⁸ also observed a significant reduction during the touch-down phase of walking gait (1.9° (0.35 to 3.45) using a similar posted non-moulded device. These findings are of large (ES 1.43) and moderated effect (ES 1.07), respectively.

Five qualitative comparisons between orthoses of different densities and a control found that each of the 12 participants responded differently to different orthoses with a tendency towards a reduction in tibial rotation.³⁴

Table 2 Quality index score

Author	Item number																				Total			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	18	19	20	26	27			
Branthwaite <i>et al</i> ³²	1	1	1	1	1	1	0	0	1	0	1	0	1	0	0	1	1	1	1	1	1	5	19	
Butler <i>et al</i> ⁴¹	1	1	1	1	1	1	1	1	0	1	1	0	1	0	0	1	1	1	1	1	1	1	5	21
Eng and Pierrynowski ³⁸	1	1	1	1	2	1	1	1	1	1	1	1	1	0	0	1	1	1	1	1	1	1	5	24
MacLean <i>et al</i> ³⁶	1	1	1	1	1	0	0	0	1	0	1	0	1	0	0	1	1	1	1	1	1	1	5	18
MacLean <i>et al</i> ³⁷	1	1	1	1	2	1	1	0	1	1	1	1	1	0	0	1	1	0	1	1	1	1	5	22
McCulloch <i>et al</i> ⁵	1	1	1	0	1	1	1	0	1	0	1	0	1	0	0	1	1	1	1	1	1	1	5	19
Miller <i>et al</i> ⁴²	1	1	1	1	0	0	0	0	1	1	1	0	1	0	1	1	1	1	1	1	1	1	5	19
Mündermann <i>et al</i> ³⁰	1	1	1	1	2	1	1	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	22
Mündermann <i>et al</i> ⁴⁶	1	1	1	1	2	0	0	1	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	21
Murley and Bird ⁴⁷	1	1	1	1	1	0	0	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	19
Nawoczenski <i>et al</i> ⁶	1	1	1	1	2	1	1	1	1	1	1	1	1	0	0	1	1	1	1	1	1	1	5	24
Nawoczenski and Ludewig ⁴⁴	1	1	1	1	2	0	1	1	1	1	1	1	1	0	0	1	1	1	1	1	1	1	5	23
Nester <i>et al</i> ⁴⁰	1	0	0	1	1	1	1	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	19
Nigg <i>et al</i> ⁴³	1	1	1	0	0	1	1	1	1	0	1	0	1	0	0	1	1	1	1	1	1	1	5	19
Nigg <i>et al</i> ³⁴	1	1	1	0	0	1	0	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	18
Nigg <i>et al</i> ³¹	1	1	1	1	0	1	1	0	1	0	1	0	1	0	0	1	1	1	1	1	1	1	5	19
Stackhouse <i>et al</i> ³⁹	1	1	1	1	1	1	0	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	20
Stacoff <i>et al</i> ¹¹	1	1	1	1	0	1	0	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	3	17
Stacoff <i>et al</i> ³³	1	1	1	1	2	1	1	1	1	0	1	1	1	0	0	1	1	1	1	1	1	1	4	22
Tomaro and Burdett ⁴⁵	1	1	1	0	0	1	1	0	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	19
Williams <i>et al</i> ³⁵	1	1	1	1	2	1	1	1	1	1	1	0	1	0	0	1	1	1	1	1	1	1	5	23
Zifchock and Davis ²⁹	1	1	1	1	2	0	0	0	1	0	1	0	1	1	1	1	1	1	1	1	1	1	5	21

Rearfoot eversion velocity

We found 19 comparisons over eight studies that examined the effects of orthoses on rearfoot eversion velocity (table 5).

Two comparisons^{11 30} that examined the effect of posted non-moulded orthoses on healthy cohorts during jogging were pooled and showed that orthoses had no effect on rearfoot eversion velocity ($-14.16^\circ/s$ (-50.34 to 22.03); fig. 3). This finding is in line with data that were not pooled across all orthoses' designs,^{5 29 30 33 39} with the exception of two comparisons. MacLean *et al*³⁷ reported that a posted moulded orthosis cast to calcaneal vertical produced a moderate (ES 0.95) reduction in velocity during the first 15% of stance phase. A later study reported a tendency for an inverted orthosis to increase velocity ($38.66^\circ/s$ (-22.44 to 99.76), ES 0.51) on initial use by individuals with a history of injury.³⁶

Maximum ankle inversion moment

We found 16 comparisons between orthoses and controls measuring maximum ankle inversion moment; in 10 of which, we could generate point estimates of effect and confidence intervals (table 6). No pooling was possible, and results were conflicting between studies where point estimates of effect could and could not be calculated.

Of the 10 comparisons where point estimates of effect could be calculated, only two significant effects were found. In a healthy cohort, a posted non-moulded orthosis significantly reduced maximum ankle inversion moment with moderate effect (6 Nm (1.28 to 10.72), ES 0.77) when jogging.³⁰ However, no effect was found using an orthosis of similar design in a currently injured cohort.³³ An inverted Blake orthosis³⁵ had a moderate reducing effect on maximum ankle inversion moment by 0.14 Nm/kg/m (-0.25 to -0.03 (ES 1.06)).

Of the six comparisons where the point estimate of effects could not be calculated, MacLean *et al*³⁷ and Stackhouse *et al*³⁹ reported that their posted moulded orthoses produced a significant reduction in ankle inversion moment between 5% and 75% of stance phase and, on average, 24% throughout stance,

respectively. Similarly, Nigg *et al*³¹ reported that orthoses with either a full length or half length medial post both significantly reduced inversion moment.

Maximum knee external rotation moment

Twelve comparisons were found, five of which provided enough data to calculate effect and confidence intervals, but pooling was not possible (table 6).^{30 36}

Where point estimates of effect and confidence intervals were calculated, orthoses had no effect on maximum knee external rotation moment regardless of design.^{30 36} Six of the remaining comparisons supported these findings. In one comparison between an orthosis with a full-length 4.5-mm medial post and control, the authors reported increased maximum knee external rotation moment (27.6%) but did not provide enough data to enable the calculation of confidence intervals.³¹

Between-orthoses comparisons

Orthoses of differing designs (eg, Cobra, Blake inverted, biplanar, moulded, posted, irregular surface) have been compared. Data pooling was not possible because of the large variability in design features of the compared orthoses.

Rearfoot eversion

We found 11 comparisons investigating the effects of differing orthoses on peak eversion and four on rearfoot eversion excursion. The majority (seven) of comparisons were between orthoses of different posting and moulding designs. Other comparisons included an irregularly surfaced orthosis, two posted moulded orthoses (one custom-made and the other "mould-of-best-fit") and different posting placement and amount.

With peak rearfoot eversion, a posted moulded orthosis was more effective than a posted non-moulded orthosis (3.3° (1.53 to 5.07), ES 1.83) and the irregular surface orthosis (2.4° (0.44 to 4.36), ES 1.2) in a currently injured cohort.³³ In participants with no history of injury, the posted non-moulded orthosis was more effective (3.2° (1.28 to 5.12), ES 1.01).³⁰

Table 3 Rearfoot eversion results where point estimates of effect and confidence intervals were able to be calculated

Authors	Outcome	Intervention	Comparator	Mean difference ° (95% CI)	Effect size (ES)
Branthwaite <i>et al</i> ³²	Peak eversion	Biplanar	Control	-3.1	0.62
		Cobra	Control	-2.1	0.46
		Biplanar	Cobra	No difference	
Eng and Pierrynowski ³⁸	Eversion excursion	Posted non-moulded	Control	Walking: contact -1.8 (-5.13 to 1.53)	0.47
				Midstance -1.8 (-3.73 to 0.13)	0.82
				Propulsion -0.4 (-2.42 to 1.62)	0.17
				Jogging: contact -2.5 (-5.49 to 0.49)	0.73
				Midstance -0.8 (-2.42 to 0.82)	0.43
				Propulsion -1.7 (-4.15 to 0.75)	0.61
MacLean <i>et al</i> ³⁶	Eversion excursion	Posted moulded (Inverted)	Control	Initial: 1.56 (-2.05 to 5.17)	0.35
				After 6 weeks of wear: 0.88 (-2.97 to 4.73)	0.18
MacLean <i>et al</i> ³⁷	Peak eversion Eversion excursion	Posted moulded	Control	Peak eversion: reduction Decreased calcaneal eversion angle	0.32 0.25
McCulloch <i>et al</i> ⁵	Peak eversion	Posted moulded	Control	Walking 3.2 km/h: -4.0 (-8.7 to 0.7)	0.75
				Walking 4.8 km/h: -2.7 (-7.29 to 1.89)	0.52
Mündermann <i>et al</i> ³⁰	Peak eversion	Posted non-moulded	Control	-2.3 (-3.82 to -0.78)	0.92
		Non-posted moulded	Control	0.6 (-0.85 to 2.05)	0.25
		Posted moulded	Control	0.9 (-0.93 to 2.73)	0.3
		Posted non-moulded	Non-posted moulded	-2.9 (-4.47 to -1.33)	1.11
		Posted non-moulded	Posted moulded	-3.2 (-5.12 to -1.28)	1.01
		Non-posted moulded	Posted moulded	-0.3 (-2.17 to 1.57)	0.1
Nigg <i>et al</i> ³¹	Peak eversion	Full-length medial post	Control	-1.5 SD ± 1.3	
	Eversion	Full-length medial post	Control	-2 SD ± 1.5	
	Excursion	Full-length lateral post	Control	2.1 SD ± 1.7	
Stacoff <i>et al</i> ³³	Peak eversion	Posted non-moulded	Control	1.5 (-0.43 to 3.43)	0.76
		Posted moulded	Control	-1.8 (-3.81 to 0.21)	0.88
		Irregular surface	Control	0.6 (-1.51 to 2.71)	0.28
		Posted moulded	Posted non-moulded	-3.3 (-5.07 to -1.53)	1.83
		Posted moulded	Irregular surface	-2.4 (-4.36 to -0.44)	1.2
		Irregular surface	Posted non-moulded	-0.9 (-2.77 to 0.97)	0.47
Stacoff <i>et al</i> ¹¹	Peak eversion	Posted non-moulded (Anterior)	Control	-0.83 (-4.34 to 2.68)	0.29
	Eversion	Posted non-moulded (Posterior)	Control	-1.03 (-4.79 to 2.73)	0.34
	Excursion	Posted non-moulded (Posterior)	Posted non-moulded (Anterior)	-0.20 (-4.01 to 3.61)	0.07
		Posted non-moulded (Anterior)	Control	-0.26 (-3.64 to 3.12)	0.1
		Posted non-moulded (Posterior)	Control	-0.34 (-3.63 to 2.95)	0.13
		Posted non-moulded (Anterior)	Posted non-moulded (Posterior)	-0.08 (-3.44 to 3.28)	0.03
Williams <i>et al</i> ³⁵	Peak eversion	4° posted moulded	Control	1.59 (-1.66 to 4.84)	0.41
	Eversion excursion	Inverted posted moulded	Control	1.2 (-1.87 to 4.27)	0.3
		Inverted posted moulded	4° posted moulded	-0.39 (-3.4 to 2.62)	0.11
		4° posted moulded	Control	-0.82 (-3.42 to 1.78)	0.29
		Inverted posted moulded	Control	-0.05 (-2.42 to 2.32)	0.02
		4° posted moulded	Inverted posted moulded	-0.77 (-3.37 to 1.83)	0.25
Zifchock and Davis ²⁵	Peak eversion	Posted moulded (semicustom)	Control	Low arch: 0.06 (-1.45 to 1.58)	0.03
	Eversion excursion	Posted moulded (custom)	Control	High arch: -0.03 (-2.02 to 1.95)	0.01
		Posted moulded (semicustom)	Posted moulded (custom)	Low arch: 1.13 (-1.46 to 3.37)	0.29
		Posted moulded (semicustom)	Control	High arch: 0.28 (-1.88 to 2.45)	0.08
		Posted moulded (custom)	Control	Low arch: -1.07 (-3.06 to 1.46)	0.28
		Posted moulded (custom)	Posted moulded (semicustom)	High arch: -0.32 (-2.09 to 1.46)	0.11
		Posted moulded (semicustom)	Posted moulded (custom)	Low arch: -0.92 (-2.46 to 0.61)	0.39
				High arch: -0.84 (-2.86 to 1.18)	0.26
				Low arch: -1.13 (-2.70 to 0.44)	0.47
				High arch: -0.76 (-2.76 to 1.23)	0.24
		Low arch: -0.2 (-1.4 to 1.0)	0.11		
		High arch: -0.08 (-1.92 to 1.76)	0.03		

No orthosis design, degree or placement of posting seemed to be more effective in influencing excursion.

Tibial internal rotation

Of the eight between-orthosis comparisons (table 4), only one reported a moderate non-significant effect (ES 1.07). A reduction of 1.05° (-0.16 to 2.26) in tibial internal rotation occurred when a non-moulded orthosis with a medial post placed under the calcaneus was compared with a post placed under the medial arch.¹¹ It would seem that no design feature is more effective at reducing tibial internal rotation.

Rearfoot eversion velocity

We found seven between-orthosis comparisons for rearfoot eversion velocity. Six involved different posting and moulding features^{20 30 32} and the other compared a posterior post with an anterior post (table 5).¹¹

Jogging in a posted non-moulded orthosis was found to have a moderate effect over posted moulded orthoses (91.5°/s (7.98 to 175.02), ES 0.66) and non-posted moulded orthoses (ES 0.6) when participants had no history of injury.³⁰ No other significant results were found.

Table 4 Tibial internal rotation comparisons of studies where point estimates of effect and confidence intervals were able to be calculated

Authors	Intervention	Comparator	Mean difference (95% CI)	Effect size (ES)
Eng and Pierrynowski ³⁸	Posted non-moulded	Control	Walking: contact -1.9 (-3.45 to -0.35), midstance 0 (-2.29 to 2.29), propulsion -0.4 (-2.46 to 1.66) Jogging: contact -0.4 (-2.42 to 1.62), midstance 0.0 (-0.81 to 0.81), propulsion -0.6 (-2.62 to 1.42)	1.07, 0, 0.17 0.15, 0, 0.26
MacLean <i>et al</i> ²⁰⁰⁸ ³⁶	Posted moulded (Inverted)	Control	Initial -1.38 (-4.99 to 2.33) After 6 weeks of wear -0.14 (-4.48 to 4.2)	0.31 0.03
Mündermann <i>et al</i> ³⁰	Posted non-moulded	Control	-0.5 (-2.95 to 1.95)	0.12
	Non-posted moulded	Control	-0.6 (-2.99 to 1.79)	0.15
	Posted moulded	Control	-0.5 (-2.98 to 1.98)	0.12
	Non-posted moulded	Posted non-moulded	-0.1 (-2.52 to 2.32)	0.02
	Posted non-moulded	Posted moulded	0 (-2.51 to 2.51)	0.0
	Non-posted moulded	Posted moulded	-0.1 (-2.55 to 2.35)	0.02
Nawoczenski <i>et al</i> ⁶	Posted moulded (low rearfoot), posted moulded (high rearfoot)	Control, control	Internal rotation with respect to rearfoot: 2.0 (-4.31 to 0.31), -2.3 (-6.26 to 1.66)	0.76, 0.37 0.31, 0.3
	Posted moulded (low rearfoot), posted moulded (high rearfoot)	Control, control	Total internal/external rotation: -1.2 (-4.64 to 2.24), -1.6 (-6.34 to 3.14)	
Stacoff <i>et al</i> ³³	Posted non-moulded	Control	0.3 (-4.16 to 4.76)	0.07
	Posted moulded	Control	-0.2 (-4.71 to 4.31)	0.04
	Irregular surface	Control	0.4 (-4.06 to 4.86)	0.09
	Posted moulded	Posted non-moulded	-0.5 (-5.96 to 3.96)	0.11
	Posted non-moulded	Irregular surface	-0.1 (-4.51 to 4.31)	0.02
	Posted moulded	Irregular surface	-0.6 (-5.06 to 3.86)	0.13
Stacoff <i>et al</i> ¹¹	Posted non-moulded (anterior)	Control	-0.54 (-1.37 to 0.29)	0.81
	Posted non-moulded (posterior)	Control	-1.59 (-2.97 to -0.21)	1.43
	Posted non-moulded (posterior)	Posted non-moulded (anterior)	-1.05 (-2.26 to 0.16)	1.07
Williams <i>et al</i> ³⁵	4° posted moulded	Control	3.33 (-1.94 to 8.6)	0.53
	Inverted posted moulded	Control	4.5 (-0.23 to 8.77)	0.88
	4° posted moulded	Inverted oosted moulded	-1.17 (-5.55 to 3.21)	0.22

Table 5 Rearfoot eversion velocity comparisons for studies where point estimates of effect and confidence intervals were able to be calculated

Authors	Intervention	Comparator	Mean difference (95% CI)	Effect size (ES)
Branthwaite <i>et al</i> ³²	Biplanar	Control	-4 (-20.37 to 12.37)	0.23
	Cobra	Control	5 (-7.55 to 17.55)	0.35
	Biplanar	Cobra	-9 (-26.4 to 8.4)	0.48
MacLean <i>et al</i> ³⁶	Posted moulded (Inverted)	Control	Initial: 38.66 (-22.44 to 99.76) After 6 weeks of wear: 37.58 (-43.9 to 119.06)	0.51 0.37
McCulloch <i>et al</i> ⁵	Posted moulded	Control	Rate of pronation first 10% stance walking 3.2 km/h: 4.5 (-13.68 to 22.68), walking 4.8 km/h: 1.5 (-21.17 to 24.17) Rate of pronation second 10% stance walking 3.2 km/h: -6.2 (-23.94 to 11.54), walking 4.8 km/h: 1.1 (-17.22 to 19.42)	0.22, 0.06 0.31, 0.05
Mündermann <i>et al</i> ³⁰	Posted non-moulded	Control	-71.8 (-159.78 to 16.18)	0.49
	Non-moulded posted	Control	12.1 (-78.74 to 102.94)	0.08
	Posted moulded	Control	19.7 (-70.01 to 109.41)	0.13
	Posted non-moulded	Non-posted moulded	-83.9 (-168.63 to 0.83)	0.6
	Posted non-moulded	Posted moulded	-91.5 (-175.02 to -7.98)	0.66
	Non-posted moulded	Posted moulded	-7.6 (-94.13 to 78.93)	0.05
Stackhouse <i>et al</i> ³⁹	Posted moulded	Control	-7.85 (not significant)	
Stacoff <i>et al</i> ¹¹	Posted non-moulded (anterior)	Control	5.0 (-35.61 to 45.61)	0.15
	Posted non-moulded (posterior)	Control	-2.42 (-42.12 to 37.28)	0.08
	Posted non-moulded (posterior)	Posted non-moulded (Anterior)	-7.42 (-45.51 to 30.67)	0.24
Zifchock and Davis 2008 ²⁵	Posted moulded (semicustom)	Control	Low arch: -10.87 (-41.37 to 19.63)	0.23
	Posted moulded (custom)	Control	High arch: -14.32 (-50.92 to 22.28)	0.25
	Posted moulded (custom)	Posted moulded (semicustom)	Low arch: -14.14 (-44.87 to 16.58)	0.3
	Posted moulded (custom)	Posted moulded (semicustom)	High arch: -20.51 (-56.08 to 15.06), -3.27 (-33.87 to 27.32), -6.19 (-34.14 to 21.76)	0.37, 0.07, 0.14


Maximum ankle inversion moment

Of the seven between-orthoses comparisons for maximum ankle inversion moment, there were six between orthoses of different moulding and posting designs and another between a

moulded-inverted orthosis to a moulded orthosis with a 4° post (table 6). The latter showed a small tendency for the inverted posted moulded orthosis to reduce maximum ankle inversion by 0.07 Nm/kg/m (-0.04 to 0.18) over the comparator (ES 0.58).³⁵

Table 6 Kinematic outcomes for comparisons where point estimates of effect and confidence intervals were able to be calculated

Authors	Outcome	Intervention	Comparator	Mean difference (95% CI)	Effect size (ES)
MacLean <i>et al</i> ³⁶	Maximum ankle inversion moment	Posted moulded (Inverted)	Control	Initial: -0.05 (-0.22 to 0.12),	0.23, 0.22
		Posted moulded (Inverted)	Control	After 6 weeks of wear: -0.07 (-0.59 to 0.45)	0.3, 0.05
Mündermann <i>et al</i> ³⁰	Maximum ankle inversion moment	Posted non-moulded	Control	-6.0 Nm (-10.72 to -1.28)	0.77
		Non-posted moulded	Control	-2.7 Nm (-7.84 to 2.44)	0.32
	Maximum knee external rotation moment	Posted moulded	Control	-4.8 Nm (-9.91 to 0.31)	0.57
		Posted non-moulded	Non-posted moulded	-3.3 Nm (-8.22 to 1.62)	0.41
	Posted non-moulded	Moulded-posted	-1.2 Nm (-6.08 to 3.68)	0.15	
		Non-posted moulded	Non-posted moulded	-2.1 Nm (-7.39 to 3.19)	0.24
	Posted non-moulded	Control	1.5 Nm (-1.24 to 4.24)	0.33	
		Control	1.4 Nm (-1.02 to 3.82)	0.35	
	Posted moulded	Control	1.8 Nm (-0.68 to 4.28)	0.44	
		Posted non-moulded	Posted non-moulded	-0.1 Nm (-2.84 to 2.64)	0.02
	Posted non-moulded	Posted moulded	-0.3 Nm (-3.09 to 2.49)	0.06	
		Non-posted moulded	Posted moulded	-0.4 Nm (-2.88 to 2.08)	0.1
Stacoff <i>et al</i> ³³	Maximum ankle inversion moment	Posted non-moulded	Control	-0.50 Nm (-11.29 to 10.29)	0.05
		Posted moulded	Control	1.4 Nm (-9.89 to 12.69)	0.12
		Irregular surface	Control	0.6 Nm (-10.46 to 11.66)	0.05
		Posted non-moulded	Posted moulded	-1.9 Nm (-12.06 to 8.26)	0.18
		Posted non-moulded	Irregular surface	-1.1 Nm (-11.0 to 8.8)	0.11
		Posted moulded	Posted moulded	-0.8 Nm (-11.24 to 9.64)	0.07
Williams <i>et al</i> ³⁵	Maximum ankle inversion moment	4° posted moulded	Control	-0.06 Nm/kg/m (-0.20 to 0.06)	0.41
		Posted moulded (Inverted)	Control	-0.14 Nm/kg/m (-0.25 to -0.03)	1.06
		Posted moulded (converted)	4° posted moulded	-0.07 Nm/kg/m (-0.18 to 0.04)	0.58

**Figure 2** Forest plot of data pooling for rearfoot eversion and tibial internal rotation. Filled diamonds represent pooled data.


Maximum knee external rotation moment

We found four between-orthoses comparisons in the literature regarding maximum knee external rotation moment (table 6), all equivocal.^{30 40}

Shock attenuation

The search strategy yielded 42 comparisons, from six studies on shock attenuation.^{30 36 40-43} Orthoses were compared on the basis of density (22), with or without rearfoot stabilisation, or

Rearfoot eversion velocity

**Figure 3** Forest plot of data pooling for rearfoot eversion velocity. Filled diamonds represent pooled data.

variations of posting and moulding (20). The outcome measures consisted of tibial acceleration (3), loading rate (19), vertical impact force (16) and vertical ground reaction force (4) (table 7).

Tibial acceleration

Three comparisons involving two densities and a control showed no differential effects on tibial acceleration (ES 0.01-0.16).⁴¹

Loading rate

Comparisons were divided into those investigating density (11),^{32 40 43} and those investigating design (7).^{30 36} The former did not differentiate on loading rate,⁴³ regardless of the presence of rearfoot stabilisation. When orthoses differed in design, non-posted moulded and posted moulded both had significant moderate effects over the control (ES 0.69 and 0.95, respectively).³⁰ These results were found in participants without injury. A reduction in loading rate brought about by a posted moulded orthosis was also reported in participants with a history of injury (-19.56 BW/s (-37.24 to -1.88)), but only after 6 weeks.³⁶

Vertical impact force

We found three studies of vertical impact force with 16 comparisons of orthoses differing in density and design.^{30 36 43}

Table 7 Shock attenuation paradigm comparisons when point estimates of effect and confidence intervals were able to be calculated

Authors	Outcome	Intervention	Comparator	Mean difference (95% CI)	Effect size (ES)
Butler <i>et al</i> ⁴¹	Tibial acceleration (g) Loading rate (BW/s)	Posted moulded (rigid)	Control	-0.2 (-2.35 to 1.95)	0.07
		Posted moulded (Soft)	Control	0.3 (-1.92 to 2.52)	0.1
		Posted moulded (rigid)	Posted moulded (soft)	-0.5 (-2.72 to 1.72)	0.16
		Posted moulded (rigid)	Control	-6 (-16.19 to 4.19)	0.47
		Posted moulded (Soft)	Control	-3.4 (-12.55 to 5.75)	0.27
		Posted moulded (rigid)	Posted moulded (soft)	-2.6 (-14.37 to 9.17)	0.16
MacLean <i>et al</i> ³⁶	Impact peak (BW) Maximum loading rate (BW/s)	Posted moulded	Control	Initial: -0.11 (-0.28 to 0.06), 6 weeks: 0.1 (-1.17 to 0.97)	0.52, 0.42
		Posted moulded	Control	Initial: -12.04 (-33.51 to 9.43), 6 weeks: -19.56 (-37.24 to -1.88)	0.43, 0.89
Mündermann <i>et al</i> ³⁰	Vertical impact peak (N) Maximum loading rate (N/s)	Posted non-moulded	Control	20.3 (-137.45 to 178.05)	0.04
		Non-posted moulded	Control	-98.7 (-249.39 to 51.99)	0.4
		Posted moulded	Control	-146.8 (-249.9 to 1.30)	0.6
		Non-posted moulded	Posted non-moulded	-119 (-272.92 to 34.92)	0.47
		Posted moulded	Posted non-moulded	-167.1 (-318.48 to -15.72)	0.67
		Posted moulded	Non-posted moulded	-48.1 (-192.11 to 95.91)	0.2
		Posted non-moulded	Control	1 (-5.96 to 7.96)	0.09
		Non-posted moulded	Control	-7.7 (-14.41 to -0.99)	0.69
		Posted moulded	Control	-10.5 (-17.15 to -3.85)	0.95
		Non-posted moulded	Posted non-moulded	-8.7 (-15.66 to -1.74)	0.76
		Posted moulded	Posted non-moulded	-11.5 (-18.4 to -4.6)	1.01
		Posted moulded	Non-posted moulded	-2.8 (-9.45 to 3.85)	0.25
Nigg <i>et al</i> ⁴³	Vertical impact force (N) Loading rate (kN/s)	Shore 26	Control	-118 (-179.88 to 115.88)	0.47
		Shore 28	Control	-92 (-255.62 to 71.62)	0.39
		Shore 29	Control	-88 (-273.76 to 97.76)	0.33
		Shore 34	Control	-63 (-235.31 to 109.31)	0.25
		Shore 26 + stabiliser	Control + stabiliser	-52 (-193.06 to 89.06)	0.26
		Shore 28 + stabiliser	Control + stabiliser	-32 (-179.88 to 115.88)	0.15
		Shore 29 + stabiliser	Control + stabiliser	-40 (-179.99 to 99.99)	0.2
		Shore 34 + stabiliser	Control + stabiliser	-52 (-202.6 to 98.6)	0.24
		Shore 26	Control	-2.92 (17.39 to 11.55)	0.14
		Shore 28	Control	-0.09 (-14.18 to 14)	0.0
		Shore 29	Control	-6.27 (-20.88 to 8.34)	0.29
		Shore 34	Control	-2.48 (-15.81 to 10.85)	0.13
		Shore 26 + stabiliser	Control + stabiliser	10.39 (-4.65 to 25.43)	0.48
		Shore 28 + stabiliser	Control + stabiliser	12.97 (-2.59 to 28.53)	0.58
		Shore 29 + stabiliser	Control + stabiliser	5.22 (-9.7 to 20.14)	0.24
		Shore 34 + stabiliser	Control + stabiliser	4.07 (-11.48 to 19.62)	0.18

Varying the material densities, irrespective of rearfoot stabilisation, had no effect on impact force. Large interparticipant variability was again found.⁴³

In terms of design, a significant moderate attenuating effect was found in favour of a posted moulded orthosis compared with a posted non-moulded orthosis (167.1 N (15.72 to 318.48), ES 0.67) in uninjured participants.³⁰ No difference existed between a posted moulded orthosis compared with control in a cohort with history of injury.³⁶

Vertical ground reaction force

Miller *et al*⁴² reported that their posted moulded orthosis produced less vertical ground reaction force at 10% and 20% of the total stance phase. This effect was not apparent when a medially posted (10°) high-density EVA orthosis and a laterally posted (10°) high-density EVA orthosis was compared with control.⁴⁰ Insufficient information was presented to calculate point estimates of effect.

Neuromotor control

Eight comparisons were found fitting the inclusion criteria. Orthoses were of different designs, including an irregularly surfaced orthosis, and were compared with a control condition. Of these comparisons, two studies (two comparisons) provided enough information to calculate point estimates of effect and their confidence intervals (table 8).^{44 45} The main outcome measure was the amplitude of EMG signal of several

muscles of the shank (tibialis anterior (TA), peroneus longus (PL), medial gastrocnemius (MG), lateral gastrocnemius, soleus (Sol), tibialis posterior (TP)) and the thigh (vastus lateralis, vastus medialis, rectus femoris, biceps femoris (BF)). Tomaro and Burdett⁴⁵ also measured the duration of TA EMG signal.

Shank muscles

Five comparisons were made for each TA, PL and MG. Jogging in a posted moulded orthosis produced significant increases in TA and PL amplitudes and a significant decrease in the amplitude of MG for uninjured participants.⁴⁶ For these participants, posted non-moulded and non-posted moulded orthoses were reported to increase both PL and MG in different phases of gait and EMG bandwidths.⁴⁶ Walking in a moulded orthosis posted to 15° produced an increase of 19% maximum voluntary contraction (MVC) in PL for participants with no injury.⁴⁷ For participants with current injury, a posted moulded orthosis also significantly increased TA amplitude (37% of MVC (5.44 to 68.56), ES 0.67).⁴⁴ Walking studies found no effect on MG. Tomaro and Burdett⁴⁵ studied individuals with a history of injury who had worn, for a minimum of 6 months, posted moulded orthoses. They reported no difference between the orthoses and control in TA and PL amplitude as well as no change in TA duration (2.6% (-2.89 to 8.09), ES 0.41).

Comparisons were also made for Sol and TP. No change in Sol amplitude was reported when participants with no history of injury walked in moulded orthoses posted to 15° compared

Table 8 Neuromotor control paradigm comparisons where point estimates of effect and confidence intervals were able to be calculated

Authors	Muscles	Intervention	Comparator	Mean difference (95% CI)	Effect size (ES)
Nawoczinski and Ludewig ⁴⁴	TA, MG, VM, VL, BF (% MVC)	Posted moulded	Control	TA: 37.5 (5.44 to 68.56), MG: -7.6 (-23.87 to 8.67), VM: -2.2 (-18.66 to 14.26), VL: -4.3 (-26.84 to 18.24), BF: -11.1 (-20.31 to -1.89)	0.67, 0.26, 0.08, 0.12, 0.68
Tomaro and Burdett ⁴⁵	TA, PL, LG	Posted moulded	Control	TA duration: 2.6% (-2.89% to 8.09%) amplitude (μV), TA: 4 (-117.28 to 125.28), PL: -7 (-93.4 to 79.4), LG: 10 (-45.22 to 65.22)	0.41, 0.03, 0.07, 0.16

LG, lateral gastrocnemius; MVC, maximum voluntary contraction; VL, vastus lateralis; VM, vastus medialis.

with walking in a control.⁴⁷ Similarly, no systematic results were found in TP amplitude in currently injured individuals walking in posted non-moulded, posted moulded and irregular surface orthoses.³³

Thigh muscles

Four comparisons were made for the quadriceps and BF. When participants with no history of injury jogged in posted moulded orthoses, all four muscles significantly increased in amplitude during various phases of gait and across different EMG bandwidths. Similar increases were also present in BF when participants wore either posted non-moulded and non-posted moulded orthoses.⁴⁶ These findings are in discordance with a study of currently injured individuals wearing posted moulded orthoses. BF amplitude significantly and moderately (ES 0.68) reduced by 11.1% of MVC (1.89 to 20.31) throughout stance, and there was no change in the activity of vastus medialis or vastus lateralis.⁴⁴

DISCUSSION

Research has primarily focused on the kinematic paradigm and least on neuromotor control. To the extent that synthesis was possible, data pooling revealed that posted orthoses that were not moulded reduced peak rearfoot eversion and tibial internal rotation in non-injured, whereas moulded orthoses with or without posting produced large reductions in loading rate and vertical impact force.

Data pooling of the kinematic paradigm showed a relatively small effect ($\sim 2^\circ$) in reducing rearfoot eversion and tibial internal rotation using skin markers to record motion, which may well be an overestimate of actual bone motion.⁴⁸ It is currently unknown whether this small motion reduction is clinically beneficial, although Nawoczinski *et al*⁶ posited that such small changes may be clinically relevant in injured runners because of the potential for cumulative effects from the high volume of repetitive/cyclical motion. More importantly, individual studies showed large confidence intervals, which indicates that practitioners should tailor their approach to each individual's clinical presentation and apply sound clinical reasoning skills when considering this pooled data.

An interesting and unexpected outcome of shock attenuation data was that altering material density had no systematic effect on tibial acceleration, loading rate or vertical impact force,^{41 43} whereas orthosis' moulding reduced loading rate^{30 36} and may favourably affect vertical impact and ground reaction forces.^{30 36 42}

Only two neuromotor control studies reported data sufficient to derive point estimates of effect, but pooling was not possible (cohorts too dissimilar). Orthoses seem to increase TA and PL activity, variably influence MG activation levels depending on speed of gait and differentially change MG and thigh muscles contingent on injury status.^{44 46 47} Further study is urgently needed in this paradigm.

The methodological quality assessment of the studies in this review identified the main issue as being the non-specific categorisation of injury type. Studies investigating cohorts with past and current injuries included a range of lower limb injuries, thereby making it difficult to apply pooled results to a specific injury in practice.

We only reviewed studies using three-dimensional motion analysis (excluding two-dimensional) because we believe that doing so ensured the most accurate representation of motion. A limitation of this approach is that we may have overlooked meaningful data, though it would seem that this is not the case.^{25 26 49} We also only pooled data from studies using similar orthoses. It must be acknowledged that not all orthoses are exactly the same, so that our pooled point estimates of effect may be underestimates. However, this should be counterbalanced by the increased precision gained by pooling data from a number of studies. Using similar orthoses for pooling strengthens the findings for specific commonly used features of orthoses, such as posting and moulding.

SUMMARY

The major conclusion of this review is that there is a large amount of variability with regard to how patients respond to orthoses. Meta-analysis showed that an orthosis that is posted without any customisation (or individual moulding) produces greater motion control at the rearfoot and tibia than a control. Orthoses that have been individually contoured are more effective at attenuating loading rate and vertical impact force than if only posted. The majority of the evidence base has been derived from individuals with no history of injury but are likely still useful as rudimentary clinical guides for sports medicine practitioners. Future research needs to focus on neuromotor control effects, especially in those with injury.

Acknowledgements K. Mills and Dr AR Chapman are supported by the Australian Research Council. Financial support for this research was received from the Australian Research Council (Australian Research Council Linkage Project Grant LP0668233).

Funding Australian Research Council, Linkage Project Grant LP0668233, first Floor, 8, Brindabella Circuit, Brindabella Business Park, Canberra Airport Act 2609, Australia.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Australian Podiatry Council.** Clinical guidelines for orthotic therapy provided by podiatrists. 1998. <http://www.apodc.com.au> (accessed March 2009).
2. **Landorf K,** Keenan AM, Rushworth RL. Foot orthosis prescription habits of Australian and New Zealand podiatric physicians. *J Am Podiatr Med Assoc* 2001;**91**:174–83.
3. **The American College of Foot and Ankle Orthopedics and Medicine.** Prescription custom foot orthoses practice guidelines. 2004. <http://www.acfoam.org>. (accessed March 2009).

4. **Johanson MA**, Donatelli R, Wooden MJ, *et al*. Effects of three different posting methods on controlling abnormal subtalar pronation. *Phys Ther* 1994;**74**:149–58; discussion 158–61.
5. **McCulloch MU**, Brunt D, Vander Linden D. The effect of foot orthotics and gait velocity on lower limb kinematics and temporal events of stance. *J Orthop Sports Phys Ther* 1993;**17**:2–10.
6. **Nawoczenski DA**, Cook TM, Saltzman CL. The effect of foot orthotics on three-dimensional kinematics of the leg and rearfoot during running. *J Orthop Sports Phys Ther* 1995;**21**:317–27.
7. **McPoil TG**, Hunt GC. Evaluation and management of foot and ankle disorders: present problems and future directions. *J Orthop Sports Phys Ther* 1995;**21**:381–8.
8. **Nawoczenski DA**, Janisse DJ. Foot orthoses in rehabilitation—what's new. *Clin Sports Med* 2004;**23**:157–67.
9. **Nigg BM**, Nurse MA, Stefanyshyn DJ. Shoe inserts and orthotics for sport and physical activities. *Med Sci Sports Exerc* 1999;**31**:S421–8.
10. **Nigg BM**, Wakeling JM. Impact forces and muscle tuning: a new paradigm. *Exerc Sport Sci Rev* 2001;**29**:37–41.
11. **Stacoff A**, Reinschmidt C, Nigg BM, *et al*. Effects of foot orthoses on skeletal motion during running. *Clin Biomech (Bristol, Avon)* 2000;**15**:54–64.
12. **Razeghi M**, Batt ME. Biomechanical analysis of the effect of orthotic shoe inserts: a review of the literature. *Sports Med* 2000;**29**:425–38.
13. **Stefanyshyn DJ**, Hettling BA. Running injuries and orthotics. *Int Sport Med J* 2006;**7**:13.
14. **Bobbert MF**, Yeadon MR, Nigg BM. Mechanical analysis of the landing phase in heel-toe running. *J Biomech* 1992;**25**:223–34.
15. **Hohmann E**, Wörtler K, Imhoff AB. MR imaging of the hip and knee before and after marathon running. *Am J Sports Med* 2004;**32**:55–9.
16. **Nigg BM**. Impact forces in running. *Curr Opin Orthop* 1997;**8**:43–7.
17. **Nigg BM**. The role of impact forces and foot pronation: a new paradigm. *Clin J Sport Med* 2001;**11**:2–9.
18. **Collins N**, Bisset L, McPoil T, *et al*. Foot orthoses in lower limb overuse conditions: a systematic review and meta-analysis. *Foot Ankle Int* 2007;**28**:396–412.
19. **Hatton AL**, Dixon J, Rome K, *et al*. Effect of foot orthoses on lower limb muscle activation: a critical review. *Phys Ther Rev* 2008;**13**:280–93.
20. **McMillan A**, Payne C. Effect of foot orthoses on lower extremity kinetics during running: a systematic literature review. *J Foot Ankle Res* 2008;**1**:13.
21. **Murley GS**, Landorf KB, Menz HB, *et al*. Effect of foot posture, foot orthoses and footwear on lower limb muscle activity during walking and running: a systematic review. *Gait Posture* 2009;**29**:172–87.
22. **Kovács I**, Tihanyi J, Devita P, *et al*. Foot placement modifies kinematics and kinetics during drop jumping. *Med Sci Sports Exerc* 1999;**31**:708–16.
23. **Tillman MD**, Hass CJ, Chow JW, *et al*. Lower extremity coupling parameters during locomotion and landings. *J Appl Biomech* 2005;**21**:359–70.
24. **Eng JJ**, Winter DA. Kinetic analysis of the lower limbs during walking: what information can be gained from a three-dimensional model? *J Biomech* 1995;**28**:753–8.
25. **McClay I**, Manal K. The influence of foot abduction on differences between two-dimensional and three-dimensional rearfoot motion. *Foot Ankle Int* 1998;**19**:26–31.
26. **Areblad M**, Nigg BM, Ekstrand J, *et al*. Three-dimensional measurement of rearfoot motion during running. *J Biomech* 1990;**23**:933–40.
27. **Downs SH**, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;**52**:377–84.
28. **Hopkins W**. A new view of statistics. 2007. <http://www.sportsci.org/resource/stats/index.html>. (accessed March 2009).
29. **Zifchock RA**, Davis I. A comparison of semi-custom and custom foot orthotic devices in high- and low-arched individuals during walking. *Clin Biomech (Bristol, Avon)* 2008;**23**:1287–93.
30. **Mündermann A**, Nigg BM, Humble RN, *et al*. Foot orthotics affect lower extremity kinematics and kinetics during running. *Clin Biomech (Bristol, Avon)* 2003;**18**:254–62.
31. **Nigg BM**, Stergiou P, Cole G, *et al*. Effect of shoe inserts on kinematics, center of pressure, and leg joint moments during running. *Med Sci Sports Exerc* 2003;**35**:314–19.
32. **Branthwaite HR**, Payton CJ, Chockalingam N. The effect of simple insoles on three-dimensional foot motion during normal walking. *Clin Biomech (Bristol, Avon)* 2004;**19**:972–7.
33. **Stacoff A**, Kramers-de Quervain I, Dettwyler M, *et al*. Biomechanical effects of foot orthoses during walking. *The Foot* 2007;**17**:143–53.
34. **Nigg BM**, Khan A, Fisher V, *et al*. Effect of shoe insert construction on foot and leg movement. *Med Sci Sports Exerc* 1998;**30**:550–5.
35. **Williams DS** 3rd, McClay Davis I, Baitch SP. Effect of inverted orthoses on lower-extremity mechanics in runners. *Med Sci Sports Exerc* 2003;**35**:2060–8.
36. **MacLean CL**, Davis IS, Hamill J. Short- and long-term influences of a custom foot orthotic intervention on lower extremity dynamics. *Clin J Sport Med* 2008;**18**:338–43.
37. **MacLean C**, Davis IM, Hamill J. Influence of a custom foot orthotic intervention on lower extremity dynamics in healthy runners. *Clin Biomech (Bristol, Avon)* 2006;**21**:623–30.
38. **Eng JJ**, Pierrynowski MR. The effect of soft foot orthotics on three-dimensional lower-limb kinematics during walking and running. *Phys Ther* 1994;**74**:836–44.
39. **Stackhouse CL**, Davis IM, Hamill J. Orthotic intervention in forefoot and rearfoot strike running patterns. *Clin Biomech (Bristol, Avon)* 2004;**19**:64–70.
40. **Nester CJ**, van der Linden ML, Bowker P. Effect of foot orthoses on the kinematics and kinetics of normal walking gait. *Gait Posture* 2003;**17**:180–7.
41. **Butler RJ**, Davis IM, Laughton CM, *et al*. Dual-function foot orthosis: effect on shock and control of rearfoot motion. *Foot Ankle Int* 2003;**24**:410–14.
42. **Miller CD**, Laskowski ER, Suman VJ. Effect of corrective rearfoot orthotic devices on ground reaction forces during ambulation. *Mayo Clin Proc* 1996;**71**:757–62.
43. **Nigg BM**, Herzog W, Read LJ. Effect of viscoelastic shoe insoles on vertical impact forces in heel-toe running. *Am J Sports Med* 1988;**16**:70–6.
44. **Nawoczenski DA**, Ludewig PM. Electromyographic effects of foot orthotics on selected lower extremity muscles during running. *Arch Phys Med Rehabil* 1999;**80**:540–4.
45. **Tomaro J**, Burdett RG. The effects of foot orthotics on the EMG activity of selected leg muscles during gait. *J Orthop Sports Phys Ther* 1993;**18**:532–6.
46. **Mündermann A**, Wakeling JM, Nigg BM, *et al*. Foot orthoses affect frequency components of muscle activity in the lower extremity. *Gait Posture* 2006;**23**:295–302.
47. **Murley GS**, Bird AR. The effect of three levels of foot orthotic wedging on the surface electromyographic activity of selected lower limb muscles during gait. *Clin Biomech (Bristol, Avon)* 2006;**21**:1074–80.
48. **Reinschmidt C**, Van Den Bogert, AJ, Lundberg A, *et al*. Tibiofemoral and tibio-calcaneal motion during walking: external vs. skeletal markers. *Gait Posture* 1997;**6**:98–109.
49. **Cornwall MW**, McPoil TG. Comparison of 2-dimensional and 3-dimensional rearfoot motion during walking. *Clin Biomech (Bristol, Avon)* 1995;**10**:36–40.