Injury risk of playing football in Futsal World Cups

Astrid Junge,¹ Jiri Dvorak^{1,2}

¹FIFA Medical Assessment and Research Center (F-MARC) and Schulthess Klinik, Zurich, Switzerland
²Fédération Internationale de Football Association (FIFA), Zurich, Switzerland

Correspondence to

Dr Astrid Junge, Schulthess Klinik, F-MARC, Lengghalde 2, CH-8008 Zurich, Switzerland; medical@fifa.org

Accepted 4 September 2010

Published Online First

19 October 2010

ABSTRACT

Background While injuries in football (soccer) have been the subject of numerous publications, little information is available about the injury risk in futsal.

Objective Analysis of the incidence and characteristics of injury in futsal players during top-level international tournaments.

Design Prospective survey.

Methods Player injuries during three consecutive Futsal World Cups were analysed using an established injury report system. The physicians of all participating teams reported all injuries after each match on a standardised injury report form. The average response rate was 93%.

Results A total of 165 injuries were reported from 127 matches, which is equivalent to an incidence of 195.6 injuries per 1000 player hours (95% CI 165.8 to 225.6) or 130.4 injuries per 1000 player matches (95% CI 110.5 to 150.3). The majority of injuries were caused by contact with another player, and 36% of the injuries occurred during non-contact activities. Most injuries affected the lower extremity (70%), followed by head and neck (13%), upper extremity (10%) and trunk (7%). The most frequent diagnoses were contusion of the lower leg (11%), ankle sprain (10%) and groin strain (8%). On average, one time-loss injury in every two matches was reported.

Conclusion While the location and diagnosis of injuries were similar in the two types of football, fewer injuries in futsal than in football were caused by contact with another player and by foul play. More information on injury mechanisms in futsal is needed to develop specific injury preventive interventions.

INTRODUCTION

Futsal is the five-a-side version of football played on a smaller pitch and usually indoors. It is played by over a million registered players worldwide (http://www.fifa.com/mm/document/fifafacts/bcoffsurv/bigcount.summaryreport_7022.pdf) and is a growing sport in many countries. It started in South America in 1930 and the first World Futsal Championship was held 1982 in Brazil. Since 1989, a Futsal World Cup is organised by the Fédération Internationale de Football Association (FIFA). The important differences in the laws of the game between football and futsal are listed in table 1.

A few scientific studies have been published on match demands and fitness parameters of futsal players.^{1–5} From the analyses of heart rates and time-motion during competitive matches, the authors concluded that futsal is a multiple-sprints sport with more high-intensity phases than football.⁴ While injuries in football have been the subject of numerous

publications,^{6–7} only two studies on injuries in futsal were found in the literature.^{8–9} In a recent national survey on sports injuries in The Netherlands,⁸ futsal was among the top 10 sports that contributed most to the absolute number of sports injuries and had the highest injury incidence per 10 000 h of sports participation (55.2; 95% CI 42.7 to 71.3). The injury rate was about 2.7 times higher than in football (20.3; 95% CI 18.4 to 22.4), but it seems that the populations participating in futsal and in football differed in age and gender.

Futsal is different from indoor football (indoor soccer), which is more common in the USA and Canada. The most important differences are that a team includes six players, and the ball can be played directly off the wall that surrounds the playing field. Few studies on injuries in indoor soccer have been published.^{10–14} The comparison of injury rates in indoor and outdoor football had inconsistent results.^{10–14}

The aim of this study was to analyse the incidence and characteristics of injury during Futsal World Cups and compare the results with the equivalent outdoor tournament (FIFA World Cup™).

METHODS

The injury surveillance system applied is well established in international football¹⁵ and other team sports^{16–17} competitions. Since 1998 it has been implemented routinely in more than 40 international football tournaments.^{15–20} The injury definition and data-collection procedure are in accordance with the respective consensus statement.²¹ An injury was defined as any physical complaint during a match which received medical attention from the team physician, regardless of the consequences with respect to absence from match or training.²¹ The physicians of all participating teams reported all injuries after each match on a standardised injury report form. Details of the methodology are published elsewhere.^{15–17–18}

Calculation of injury rates and statistical analysis

Injury rates were expressed per match, per 1000 player hours and per 1000 player matches. Player hours were calculated by multiplying the number of returned injury report forms by five players by 40/60 h. Extra time and reduced numbers of players on the field were not taken into account because only few matches went into extra time, and the exclusion of a player is a rare exception.¹⁵ Injuries per 1000 player hours were calculated: number of injuries multiplied by 1000 divided by total player hours. Player

This paper is freely available online under the BMJ Journals unlocked scheme, see <http://bjsm.bmj.com/info/unlocked.dtl>

matches were calculated by multiplying number of returned injury report forms by five players. Injuries per 1000 player matches were calculated: number of injuries multiplied by 1000 divided by total player matches. For incidence rates, 95% CIs were calculated as the incidence ± 1.96 times the incidence divided by the square root of the number of injuries. Statistical methods applied were frequencies, cross-tabulations and χ^2 test. Significance was accepted at the 5% level.

Tournaments

Three consecutive Futsal World Cups (2000, 2004, 2008) were included in the present study. These tournaments comprised 136 matches, equivalent to 906.7 player hours (table 2). A total of 253 injury report forms, covering 843 player hours, were completed and returned. The mean response rate was 93%.

Table 1 Differences in the laws of the game between football and futsal

	Football	Futsal
Pitch	90–120×45–90 m	38–42×18–25 m
Ball	410–450 g, circumference 68–70 cm	400–440 g, circumference 62–64 cm
Players per team	11	5
Substitutions	3	Unlimited 'flying' (12 players on a team)
	No substitution for player sent off	Player sent off can be substituted for after 2 min or other has scored
Duration	45 min halves	20 min halves
	Running clock	Stopped clock
Time-outs	No time-outs	1 time-out per half and team
Restart the match	No absolute time limit	4 s rule on restarts
	Throw-in	Kick-in
Offside rule	Yes	No
Accumulated fouls	No	Yes

RESULTS

A total of 165 injuries were reported from the three tournaments, which is equivalent to an incidence of 1.30 injuries per match (95% CI 1.10 to 1.50), 195.6 injuries per 1000 player hours (95% CI 165.8 to 225.6) or 130.4 injuries per 1000 player matches (95% CI 110.5 to 150.3). The injury rate was similar in both halves, and in the first and second 10 min periods of both halves.

Most injuries were caused by contact with another player, and 36% of the injuries occurred during non-contact activities. Less than half (46%) of the contact injuries (43 of 93), or 26% of all injuries, were caused by foul play based on the judgement of the team physician, but about two-thirds of these (27 of 43) were followed by a sanction of the referee.

The injuries affected the lower extremity (n=115; 69.7%) most frequently, followed by head and neck (n=21; 12.7%), upper extremity (n=17; 10.3%) and trunk (n=12; 7.3%). The body parts predominantly injured were knee (n=26; 15.8%), thigh (n=23; 13.9%), ankle and lower leg (each n=20; 12.1%). Most injuries were diagnosed as contusions (n=73; 44.2%), sprains or ligament ruptures (n=32; 19.4%) and strains or muscle fibre ruptures (n=29; 17.6%). Seven injuries were diagnosed as concussion, three as ligament rupture in the knee, two as fracture (finger and foot) and one as a lesion of the meniscus (table 3). The most frequent diagnoses were contusion of the lower leg (n=18; 10.9%), ankle sprain (n=16; 9.7%) and groin strain (n=13; 7.9%).

Information in relation to time loss in sport was available for 138 (84%) injuries. At least 67 injuries (48.6%) were expected to prevent the player from participating in match or training. The incidence of time-loss injuries was 79.5 per 1000 player hours (95% CI 59.7 to 98.5), 53.0 per 1000 player matches (95% CI 40.3 to 65.7) or approximately one in every two matches. The most prevalent diagnoses of time-loss injuries were groin and thigh strains and ankle sprains (each n=8; 11.9%).

Table 2 Incidence and characteristics of injuries in Futsal World Cups 2000–2008

Tournament	Guatemala 2000 ¹⁵	Chinese Taipei 2004	Brazil 2008
No of matches	40	40	56
Injury report forms returned	66 (82.5%)	80 (100%)	107 (95.5%)
Player hours documented	220	266.7	356.7
No of injuries	42	63	60
Per 1000 player hours (95% CI)	190.9 (133.1 to 248.7)	236.0 (177.8 to 294.2)	168.5 (125.8 to 211.2)
Per 1000 player matches (95% CI)	127 (88.6 to 165.4)	158 (119.0 to 197.0)	111 (83.0 to 149.0)
Circumstances	N (%)	N (%)	N (%)
Non-contact injuries	10 of 41 (24%)	19 of 58 (33%)	28 of 58 (48%)
Contact injuries	31 of 41 (76%)	39 of 58 (67%)	30 of 58 (52%)
Contact-injury caused by foul	14 of 27 (52%)	25 of 39 (64%)	4 of 27 (15%)
Foul sanctioned by the referee	6 of 13 (46%)	18 of 25 (72%)	3 of 4 (75%)
Estimated severity of injury	N (%)	N (%)	N (%)
0 days	21 (53.3%)	29 (64%)	21 (41%)
1–3 days	12 (31.6%)	8 (18%)	23 (43%)
4–7 days	1 (2.6%)	6 (13%)	0 (0%)
>1 week <1 month	3 (7.9%)	2 (3%)	8 (15%)
>1 month	1 (2.6%)	–	1 (2%)
Missing	4	18	7
Time-loss injuries*	17	18	32
Per 1000 player hours	≥77.2	≥67.5	≥89.9
(95% CI)	(40.5 to 113.9)	(22.3 to 98.7)	(58.8 to 121.0)
Per 1000 player matches	≥52	≥45	≥60
(95% CI)	(27.3 to 76.7)	(24.2 to 65.8)	(39.2 to 80.8)

*Information on time loss from sport is missing in 27 injuries.

Table 3 Diagnosis of injuries in FIFA Futsal World Cups 2000–2008

Location and type	No of injuries		
	Total*	Without absence	With absence
Head/neck	21 (12.7%)*	16 (13.9%)	4 (6.0%)
Concussion	7*	4	2
Sprain	4	4	0
Contusion	8	8	0
Laceration	2	0	2
Trunk	12 (7.3%)	6 (8.3%)	6 (9.0%)
Strain	2	0	2
Contusion	6	5	1
Laceration	2	1	1
Others	2	0	2
Upper extremity	17 (10.3%)*	13 (18.1%)	3 (4.5%)
Fracture	1	1	0
Sprain	3	2	1
Contusion	10*	9	–
Others	3	1	2
Hip/groin	17 (10.3%)*	8 (11.1%)	8 (11.9%)
Strain (all groin)	13	5	8
Others (all hip)	4*	3	–
Thigh	23 (13.9%)*	3 (4.2%)	15 (22.4%)
Strain	9	1	8
Contusion	10*	2	3
Others	4	0	4
Knee	26 (15.8%)*	6 (8.3%)	14 (20.9%)
Ligament rupture	3*	–	2
Lesion of meniscus	1	–	1
Sprain	9*	1	6
Contusion	10*	4	3
Others	3	1	2
Lower leg	20 (12.1%)*	11 (15.3%)	5 (7.5%)
Contusion	18*	9	5
Strains	2	2	0
Ankle†	20 (12.1%)*	6 (8.3%)	9 (13.4%)
Sprain	16*	5	8
Contusion	3*	1	1
Foot/toe	9 (5.5%)*	3 (4.2%)	3 (4.5%)
Fracture	1	0	1
Strain	1	0	1
Contusion	7*	3	1

*Information on time loss from sport is missing in at least one injury (total: 27 injuries).

†Information on type is missing in one case.

DISCUSSION

This present study is the first survey of the incidence and characteristics of injuries in top-level futsal players. The methodology applied is well established, and its strengths and weaknesses have been discussed previously.^{15 18}

During the three recent FIFA Futsal World Cups, the team physicians reported 165 injuries, equivalent to 195.6 injuries per 1000 player hours (95% CI 165.8 to 225.6). This injury rate was 2.6 times higher than the average of the FIFA World Cups 1998,¹⁵ 2002¹⁹ and 2006²⁰ (74.2; 95% CI 67.5 to 80.9). A higher exposure-time related injury rate in futsal compared with football was also reported from a national survey on sports injuries in The Netherlands.⁸ However, football is played with a running clock, while in futsal the clock is stopped when the ball is out of play (see table 1). Junge *et al*¹⁷ have previously discussed the methodological dilemma when comparing injury incidences of different sports and concluded that it is more adequate to compare the injury rate in relation to 'athlete's

exposures' or for team sports: player matches.^{17 22} The injury incidence per 1000 player matches was similar in the Futsal and the Football World Cups (111.2; 95% CI 101.1 to 121.3). Almost half (48.6%) of the injuries incurred in the Futsal World Cups were expected to prevent the player from participating in match or training but two-thirds (68%) in the equivalent football competitions. Thus, the incidence of time-loss injuries per 1000 player matches was significantly lower in the Futsal than in the Football World Cups (72.4; 95% CI 62.3 to 82.5).

Similar to football injuries,^{6 7} most injuries in the present study were incurred by contact with another player and affected the lower extremity. However, a more detailed comparison of injuries incurred in the Futsal and the FIFA World Cups showed that a higher percentage of futsal than football injuries were incurred without contact with another player (37% vs 27%). In addition, in futsal, fewer contact injuries were caused by foul play than in football (46% vs 56%), probably because futsal referees punished injury-causing fouls more frequently than football referees (64% vs 55%).

The majority of injuries affected the lower extremity, and the most frequent diagnoses were contusions of the lower leg and ankle sprain, as reported previously for elite male football players.^{6 7} In addition, groin strains were among the three most common time-loss injuries which is different from top-level football competitions.^{15 19 20 23} It can be assumed that multiple sprints with frequent changes in direction contribute to the higher proportion of non-contact injuries, such as groin strains.

Seven concussions were reported from the three Futsal World Cups compared with six in the three football world cups. If these figures are related to the number of player matches, the incidence of concussions was 3.6 times higher in futsal than in football (5.15 vs 1.42 per 1000 player matches, NS). In the present study, all concussions were caused by contact with another player, and half of the injury situations were classified as foul play by the team physicians compared with 26% of all injuries. In international football, video analysis of the injury situation has shown that the (unfair) use of the arm to tackle the opposing player during vertical jumps was most likely to cause a head or neck injury, but none of the incidents resulting in a concussion was deemed to be a foul by the match referee.²⁴ Based on the results of that study, the referees of the 2006 FIFA World Cup™ were encouraged to severely sanction injurious fouls such as intentional elbows to the head. In the 2006 FIFA World Cup™, only one concussion occurred, compared with four in 2002, and the total number of head injuries dropped from 25 to 13.²⁰

Future studies should investigate injury mechanisms in futsal using video analysis, which has provided valuable information about injury mechanisms in professional male football players.^{25 26} Such analysis may provide additional information to develop preventive interventions to reduce the incidence of injury in futsal competitions.

The present study on injuries during top-level competitions and a Dutch national survey⁸ showed that injuries are frequent in futsal. In light of the growing popularity of futsal, future studies should evaluate the incidence, characteristics and causation of injuries on all levels of play during the season as a basis for the development of injury-prevention programmes.

CONCLUSION

The exposure-time-related incidence of injury in world cups was higher in futsal than in football, but the risk of a time-loss

What is already known on this topic

- Futsal is a popular sport in many countries, and a Futsal World Cup takes place every 4 years.
- There is only one study on the incidence of futsal injury, and this stated that futsal was among the top 10 sports that contributed most to the absolute number of sports injuries in The Netherlands.
- In this national survey on sports injuries in The Netherlands, futsal had the highest injury incidence per 10 000 h of sports participation.

What this study adds

- Based on exposure time, the incidence of injury in world cups is higher in futsal than in football, but the risk of a time-loss injury per player match was lower in futsal than in football.
- The majority of injuries were incurred due to contact with another player as reported previously for match injuries of football players, but in contrast to the corresponding tournaments of male football players, fewer futsal injuries were caused by foul play.
- The location and type of injuries were similar in football and futsal, but groin strains were more frequent in the Futsal than in the Football World Cups.
- More information on the incidence, characteristics and causation of futsal injuries on all levels of play during the season are required to develop specific injury-prevention programmes.

injury in relation to player matches was lower. While the location and type of injuries were similar between the two ways of playing football, the injury mechanism varied with fewer futsal than football injuries caused by contact with another player and by foul play. However, concussions are frequent in futsal and (too) often caused by unfair tackles. More information on injury mechanisms in futsal is needed to develop specific injury-preventive interventions.

Acknowledgements The authors highly appreciate the cooperation of all team physicians who volunteered their time to collect the data for this project. The authors thank R Charles and A König, for their help with the data collection, and D O'Riordan, for his assistance in preparing the manuscript.

Funding The authors gratefully acknowledge Fédération Internationale de Football Association for the funding of this study.

Competing interests None.

Ethics approval Ethics approval was provided by the University of Zurich, Switzerland.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. **Barbieri FA**, Gobbi LT, Santiago PR, *et al*. Performance comparisons of the kicking of stationary and rolling balls in a futsal context. *Sports Biomech* 2010;**9**:1–15.
2. **Alvarez JC**, D'Ottavio S, Vera JG, *et al*. Aerobic fitness in futsal players of different competitive level. *J Strength Cond Res* 2009;**23**:2163–6.
3. **Tessitore A**, Meeusen R, Pagano R, *et al*. Effectiveness of active versus passive recovery strategies after futsal games. *J Strength Cond Res* 2008;**22**:1402–12.
4. **Castagna C**, D'Ottavio S, Granda Vera J, *et al*. Match demands of professional Futsal: a case study. *J Sci Med Sport* 2009;**12**:490–4.
5. **Barbero-Alvarez JC**, Soto VM, Barbero-Alvarez V, *et al*. Match analysis and heart rate of futsal players during competition. *J Sports Sci* 2008;**26**:63–73.
6. **Junge A**, Dvorak J. Soccer injuries: a review on incidence and prevention. *Sports Med* 2004;**34**:929–38.
7. **Giza E**, Micheli LJ. Soccer injuries. *Med Sport Sci* 2005;**49**:140–69.
8. **Schmikli SL**, Backx FJ, Kemler HJ, *et al*. National survey on sports injuries in The Netherlands: target populations for sports injury prevention programs. *Clin J Sport Med* 2009;**19**:101–6.
9. **Cain LE**, Nicholson LL, Adams RD, *et al*. Foot morphology and foot/ankle injury in indoor football. *J Sci Med Sport* 2007;**10**:311–19.
10. **Hoff GL**, Martin TA. Outdoor and indoor soccer: injuries among youth players. *Am J Sports Med* 1986;**14**:231–3.
11. **Putukian M**, Knowles WK, Swere S, *et al*. Injuries in indoor soccer. The Lake Placid Dawn to Dark Soccer Tournament. *Am J Sports Med* 1996;**24**:317–22.
12. **Lindenfeld TN**, Schmitt DJ, Hendy MP, *et al*. Incidence of injury in indoor soccer. *Am J Sports Med* 1994;**22**:364–71.
13. **Emery CA**, Meeuwisse WH. The effectiveness of a neuromuscular prevention strategy to reduce injuries in youth soccer: a cluster-randomised controlled trial. *Br J Sports Med* 2010;**44**:555–62.
14. **Emery CA**, Meeuwisse WH. Risk factors for injury in indoor compared with outdoor adolescent soccer. *Am J Sports Med* 2006;**34**:1636–42.
15. **Junge A**, Dvorak J, Graf-Baumann T, *et al*. Football injuries during FIFA tournaments and the Olympic Games, 1998–2001: development and implementation of an injury-reporting system. *Am J Sports Med* 2004;**32**(1 Suppl):80S–9S.
16. **Langevoort G**, Myklebust G, Dvorak J, *et al*. Handball injuries during major international tournaments. *Scand J Med Sci Sports* 2007;**17**:400–7.
17. **Junge A**, Langevoort G, Pipe A, *et al*. Injuries in team sport tournaments during the 2004 Olympic Games. *Am J Sports Med* 2006;**34**:565–76.
18. **Junge A**, Dvorak J. Injuries in female football players in top-level international tournaments. *Br J Sports Med* 2007;**41**(Suppl 1):i3–7.
19. **Junge A**, Dvorak J, Graf-Baumann T. Football injuries during the World Cup 2002. *Am J Sports Med* 2004;**32**(1 Suppl):23S–7S.
20. **Dvorak J**, Junge A, Grimm K, *et al*. Medical report from the 2006 FIFA World Cup™ Germany. *Br J Sports Med* 2007;**41**:578–81; discussion 581.
21. **Fuller CW**, Ekstrand J, Junge A, *et al*. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med* 2006;**40**:193–201.
22. **Junge A**, Engebretsen L, Alonso JM, *et al*. Injury surveillance in multi-sport events: the International Olympic Committee approach. *Br J Sports Med* 2008;**42**:413–21.
23. **Ekstrand J**, Timpka T, Häggglund M. Risk of injury in elite football played on artificial turf versus natural grass: a prospective two-cohort study. *Br J Sports Med* 2006;**40**:975–80.
24. **Fuller CW**, Junge A, Dvorak J. A six year prospective study of the incidence and causes of head and neck injuries in international football. *Br J Sports Med* 2005;**39**(Suppl 1):i3–9.
25. **Fuller CW**, Smith GL, Junge A, *et al*. The influence of tackle parameters on the propensity for injury in international football. *Am J Sports Med* 2004;**32**:43S–53S.
26. **Andersen TE**, Larsen Ø, Tenga A, *et al*. Football incident analysis: a new video based method to describe injury mechanisms in professional football. *Br J Sports Med* 2003;**37**:226–32.