



Editor's choice
Scan to access more
free content

What is the evidence for chronic concussion-related changes in retired athletes: behavioural, pathological and clinical outcomes?

Paul McCrory,¹ Willem H Meeuwisse,^{2,3} Jeffrey S Kutcher,⁴ Barry D Jordan,⁵ Andrew Gardner⁶

► Additional data are published online only. To view this file please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2013-092248>)

For numbered affiliations see end of article.

Correspondence to

Dr Paul McCrory, The Florey Institute of Neuroscience and Mental Health, Heidelberg, VIC 3084 Australia; paulmccrory@icloud.com

Received 25 January 2013

Accepted 26 January 2013

ABSTRACT

Objective The purpose of this paper was to review the current state of evidence for chronic traumatic encephalopathy (CTE) in retired athletes and to consider the potential differential diagnoses that require consideration when retired athletes present with cognitive and psychiatric problems.

Data sources MEDLINE, CINAHL, EMBASE, Mosby's Index, PsycEXTRA, PsycINFO and Scopus. Key words included CTE, dementia pugilistica, punch drunk syndrome, traumatic encephalopathy, CTE, repetitive head injury, sports concussion, multiple concussions, chronic concussions, subconcussive blow and sports-related traumatic brain injury.

Results At present, there are no published epidemiological, cross-sectional or prospective studies relating to modern CTE. Owing to the nature of the published studies, being case reports or pathological case series, it is not possible to determine the causality or risk factors with any certainty. As such, the speculation that repeated concussion or subconcussive impacts cause CTE remains unproven. The extent to which age-related changes, psychiatric or mental health illness, alcohol/drug use or coexisting dementing illnesses contribute to this process is largely unaccounted for in the published literature.

Conclusions At present, the interpretation of causation in the modern CTE case studies should proceed cautiously. The causal assumptions require further prospective or longitudinal studies on the topic.

INTRODUCTION

Considerable attention surrounds the potential for long-term problems in athletes with high exposure to head impacts during a career in collision sport.¹ There is evidence^{2–4} supporting an association between long-term cognitive, neurobehavioural, psychiatric problems and participation in sport.⁵ Given that sports-related concussion is a common injury and that concussive or subconcussive blows to the head or body are an inevitable consequence of sports participation, if a causal relationship between these impacts and later-life neuropsychiatric disease exists, then potentially an enormous number of retired athletes would be at risk.⁶ Based on the published case studies, however, one estimate is that fewer than 4% of retired US professional football players may be at risk for this condition⁷ rather than all exposed athletes, raising the issue that this may not be a part of impact exposure but rather due to other as yet unidentified factors.

A recent report by McKee *et al*⁸ has suggested that chronic traumatic encephalopathy (CTE) may represent a unique tauopathy with characteristic pathological stages; however, the published methodology does not allow a causal relationship to be determined between concussion or subconcussive impacts being a risk factor for CTE.

The purpose of this paper was to review the current state of evidence for CTE in retired athletes and to consider the potential differential diagnoses that require consideration when retired athletes present with cognitive and psychiatric problems.

METHODS

Articles were retrieved via online database searching, hand-searching reference lists and cited reference searches. The online databases of MEDLINE, CINAHL, EMBASE, Mosby's Index, PsycEXTRA, PsycINFO and Scopus were searched. Key words, MeSH terms and combinations of these were used to systematically search the databases. Key words included CTE, dementia pugilistica, punch drunk syndrome, traumatic encephalopathy, CTE, repetitive head injury, sports concussion, multiple concussions, chronic concussions, subconcussive blow and sports-related traumatic brain injury (TBI).

CLINICAL SYNDROMES OF LONG-TERM PROBLEMS FOLLOWING CONCUSSION

The clinical characterisation of the presentations of athletes with chronic postconcussive symptoms is poorly defined and may reflect intrinsic differences (eg, genetic) between individuals rather than the oversimplified understanding that these syndromes are due to concussive or subconcussive trauma alone. In some cases, the persistent 'postconcussive' symptoms may, in large part, be due to unrecognised depression or anxiety,⁹ which may be labelled as 'overtraining' or even athlete 'burnout'.¹⁰

If we assume concussion is the acute clinical syndrome with 90–95% of cases recovering in less than 10 days, then there is a small group where *prolonged postconcussive symptoms* exist, that is, where the symptoms persist more than 10 days after a single episode of acute concussion (5–10% of cases) but full recovery eventually ensues usually within a matter of weeks.¹¹ The clinical and neuropsychological features are those of a resolving acute injury and perhaps this entity should be considered as part of the acute syndrome.

In addition to the acute syndrome, a number of distinct but as yet poorly defined clinical subsets exist that may shed light on the process of recovery

To cite: McCrory P, Meeuwisse WH, Kutcher JS, *et al*. *Br J Sports Med* 2013;**47**:327–330.

from concussive injury. These subsets are based on clinical experience and are not supported by hard scientific evidence at this time. Nevertheless, it may be useful to consider the full spectrum of presentations of athletes with persistent or permanent symptoms and the terminology of these states.

These include:

- ▶ Prolonged postconcussive symptoms as discussed above.
- ▶ Persistent postconcussion symptoms after one or more concussions where recovery is slow and may take months to years.¹² The DSM-IV criteria for Post Concussion Syndrome is invoked only if the symptoms last beyond 3 months. Approximately 80% of such patients recover fully with time.¹³ Cognitive testing reveals mainly attentional deficits, and structural neuroimaging is normal.
- ▶ Permanent PCS—as per above, but these individuals do not recover fully, and this has been estimated at between 10% and 20% of all cases of persistent concussion symptoms. Functional MR and/or electrophysiological change may be present, but normal structural neuroimaging is typical.¹⁴
- ▶ Proposed CTE where chronic cognitive and/or neurobehavioural dysfunction exists and a pathological diagnosis is subsequently confirmed.^{3 6} It is worth noting that there are cases where the clinical phenotype is the same as pathologically confirmed CTE cases; however, no neuro-pathological change is demonstrated at autopsy.⁸

As well as these broad categories, there are other athletes who *de novo* manifest mental health issues (depression, anxiety, suicide) or neurobehavioural problems in the absence of persistent or prolonged concussive symptoms dating from the time of an injury. Whether these represent a variant of the subsets above or simply reflect the high incidence of such disorders in society remains to be elucidated. The risk factors for such complications following concussion remain unclear, although they may be related to repeated concussions, at least in retrospective surveys.^{2 15} Prospective studies of head-injured individuals, with neuropathological and clinical verification, are needed to improve understanding of head trauma as a risk factor for sequelae.¹⁶

As well as clinical symptoms, there is some limited objective evidence of persistent neurophysiological, cognitive¹⁶ and radiological¹⁷ deficits up to 30 years following concussion.

PHYSIOLOGY OF CONCUSSION

Understanding the pathophysiology of concussion would be expected to lead to an improvement in the assessment of deficit and recovery following injury, and to facilitate the accurate classification of severity. For example, little is known about the anatomical localisation of common clinical features such as headache, loss of consciousness (LOC), difficulty with concentration, sleep disturbance and fatigue. Moreover, it is currently unknown whether mild TBI reflects a single clinical entity with a linear spectrum of injury or even distinct injury subtypes. It is hoped that this understanding may give some insight into whether *in vivo* diagnosis of CTE is possible.

In animal models following acute injury, the release of neurotransmitters and ionic fluxes occurs (known as the 'neurometabolic cascade'),¹⁸ which in turn leads to changes in cell membrane function. Animal studies suggest that, during the glucose metabolic depression phase (1–10 days postinjury), the brain is more vulnerable to repeat injury.¹⁸ Changes in the intracellular fluid status or the presence of axonal swelling may be detected using imaging techniques such as advanced diffusion weighted imaging (DWI), which also allow mapping of white

matter fibre tracts in the central nervous system.¹⁹ Preliminary studies have demonstrated DWI changes in the acute setting following mild TBI in a small cohort of adolescent patients with normal CT scans as well as in a cohort of asymptomatic professional boxers.²⁰ Functional neuroimaging techniques, for example, functional MRI (fMRI) have demonstrated changes in brain function following sport-related TBI.²¹ The main limitation of this technique is that it only reveals regions of the brain that are active in the specific cognitive task being studied. Another imaging modality, Functional Connectivity, is able to detect realtime resting state networks and may provide an alternative to fMRI for identifying brain subregions and networks that are affected in mild TBI.²² Other techniques such as MR spectroscopy (MRS) allow the detection of metabolic disturbances following mild TBI through the measurement of intracellular metabolites. Preliminary studies in a small cohort of collegiate athletes suggest a role of mitochondrial dysfunction in the postinjury metabolic depression.²³ Other studies using MRS have also demonstrated that the *N*-acetylaspartate/creatine ratio (which reflects neuronal cell damage) is related to injury severity and outcome even when white matter appears normal on MRI.²⁴

DEMENTIA RATES IN THE GENERAL POPULATION

Given that cognitive impairment is one of the key features of the proposed CTE, it is important to understand the risk of this in similar age groups in the general population. The reported incidence and prevalence rates of dementia vary according to the population studied, and prevalence approximately doubles every 5 years from the age of 65 years. The incidence of dementia (all causes) in the 30-year-old to 64-year-old group is 54/100 000, and for the 45-year-old to 64-year-old group, it is 98.1 : 100 000.²⁵

Whether mild, or repetitive mild TBI (mTBI) increases an individual's risk for developing Alzheimer's disease (AD) has been a long-standing topic of contention. While mixed results have been reported regarding the association between moderate and severe TBI and AD, the association between mTBI and AD appears to be less strong. For example, in a systematic review, Bazarian *et al*²⁶ concluded that there was limited support for the notion that mTBI (with LOC) results in an increased risk for later life AD. Further, the authors also concluded that there was insufficient evidence to determine whether an association between mTBI (without LOC) and AD exists.

DEPRESSION IN THE GENERAL POPULATION

Depression and cognitive impairment are both common conditions in older age, and they frequently occur together. A US-based epidemiology study examining major depression reported that the incidence of this disorder in men aged 25–34, 35–44 and 45–54 years was 12.3%, 11.0% and 8.6%, respectively.²⁷ This is much higher among individuals with dementia, where it has been reported that 25–50% of all patients with dementia will develop depression at some point over the course of their illness.^{28 29}

DEPRESSION, COGNITIVE IMPAIRMENT AND NEURODEGENERATIVE DISEASES IN FORMER ATHLETES

In a study of retired National Football League (NFL) players, responses to questions regarding clinical depression revealed that 269 (11.1%; 95% CI 9.9% to 12.3%) of 2434 respondents reported a previous diagnosis of clinical depression. In comparison to retired NFL players with no history of concussion, retired players with a history of one or two previous

concussions were 1.5 times more likely to be diagnosed with depression, while those with a history of three or more previous concussions were found to be three times more likely to be diagnosed with depression.¹⁵

The results of a screening survey of 513 retired NFL players (average age=61 years) reported that 35% produced scores suggesting a possibly mild cognitive impairment.² In another recent study of former NFL players, the death rate from neurodegenerative diseases was three times greater than that of the general population, although the number of cases in this study was small. More specifically, the rates of diagnosis of AD and amyotrophic lateral sclerosis (ALS) were found to be four times higher in former NFL players than the general population.³⁰ Of the 334 former athletes in this cohort, seven (2.1%) had ALS listed on their death certificates. This issue of ALS is in keeping with the TDP43 tauopathy seen in CTE cases.³¹

BRAIN PATHOLOGY IN NORMAL AGEING

The Honolulu-Asia Aging Study provides a unique longitudinal model of ageing and disease and offers considerable insight into what may be considered *normal ageing* from a pathological standpoint.³² From 1991 to 1993, 3 yearly follow-up clinical and formal neuropsychological examinations have been conducted in 3508 men who were free of dementia and a total of 593 standardised brain autopsies have been conducted.³²

Postmortem brain examination demonstrated diverse pathology, even in individuals clinically diagnosed as 'pure AD,' with fewer than 50% demonstrating the typical pathological features of AD. Furthermore, neuropathological abnormalities were also observed in approximately 40% of neuropsychologically normal patients.³²⁻³⁴ Similarly, disparity between clinical presentation (ie, living diagnosis) and postmortem neuropathology has also been demonstrated in other ageing study samples.³⁵⁻³⁶ It is now clear from these ageing studies that postmortem findings may not represent pathology and may not equate to clinical symptomatology or a syndrome³⁷ and may be seen in cognitively normal older adults. Whether the 'gold standard' of neuropathology in neurological disease is true has been recently questioned.³⁸

THE 'CLASSIC' SYNDROME OF CTE IN PROFESSIONAL BOXERS

In 1928, Dr Harrison Martland first described the 'punch drunk' or CTE state in retired boxers.³⁹ The incidence of classic CTE has proven difficult to establish, due, in the main, to a lack of prospective studies. Roberts,⁴⁰ who randomly sampled 250 retired boxers from a cohort of 16 781 UK boxers registered between 1929 and 1955, reported that in 37 boxers (17%) clinically demonstrable lesions of the nervous system were present. Notably, the oldest boxers in this cohort fought in the late 1800s, in an era where bare-knuckle championships were still conducted, frequent fights occurred even when boxers were concussed and there was little medical supervision or weight matching of boxers; however, only 11 of the 37 cases were elaborated on in this study. A close analysis of the clinical details raises suspicion regarding the certainty of the diagnosis of CTE in most cases.⁴¹

In the published cases, cognitive deterioration was typically detected 10–20 years subsequent to cessation of exposure to repetitive head trauma (ie, postretirement).⁶ Interestingly, in all cases where details were provided, the physical signs but not the cognitive deficits progressed postretirement. There are two distinct clinical syndromes that have been demonstrated in this data set; the first (which occurs in approximately 70% of cases) includes dysarthria, pyramidal problems and cognitive deficits.

As the disease manifests clinically, these cognitive abnormalities include difficulties in memory, information processing speed, insight and orientation. The second clinical syndrome (in approximately 30% of cases) includes dysarthria and pyramidal problems, but with intact cognitive abilities.⁴⁰⁻⁴²⁻⁴⁴ Movement disorders were reported to be present in approximately two in every five reported cases.³

One of the difficulties of ascribing the clinical syndrome solely to boxing is the presence of comorbidities plus risk factors for other conditions that may also result in cognitive deterioration. A case report of a champion boxer with cognitive decline highlights these issues of multifactorial causation of cognitive problems.⁴⁵

The neuropathological features of classic CTE have been described in detail⁴⁶ and typically result in a cavum septum pellucidum with septal fenestration; cerebellar scarring involving Purkinje cell loss and thinning of the granular layer; degeneration of the substantia nigra and locus caeruleus; and diffuse neurofibrillary tangles (NFT) involving the medial temporal region, uncus, amygdala, hippocampus, parahippocampal gyrus and fusiform gyrus along with the more lateral temporal, insular and frontal cortices. The extent of neuropathology appears to be positively correlated with the level of exposure.⁴⁶ Roberts *et al*⁴⁴ examined 14/15 brains originally described by Corsellis *et al*,⁴⁶ as well as six additional boxers' brains using immunocytochemistry, and 19/21 cases also demonstrated widespread diffuse amyloid deposits.

Recently, a propagation model of neurodegeneration has been proposed, suggesting that tau positive NFT phosphorylation may progressively spread from one neuron to adjacent neurons in the absence of ongoing triggering factors.⁴⁷⁻⁵⁰ The implication of this finding is that the neuropathological findings may differ or progress in different stages of the condition. This finding may help resolve the differences between reports of specific sites of tau positive NFT deposition in CTE and other studies that suggest widespread changes.

THE 'MODERN' SYNDROME OF CTE IN FOOTBALLERS

Recent publications³⁻⁵¹ on CTE in retired athletes have introduced a number of conceptual changes in the clinical features⁵²⁻⁵³ and outcomes, and also the neuropathology, as compared with the classic entity described by Roberts⁴⁰ and Corsellis *et al*⁴⁶ in their boxing subjects.

PUBLISHED MODERN CTE CASES

A number of cases of CTE in retired athletes have been published. The index case was reported by Omalu and colleagues in 2005, with additional cases reported subsequently.⁴⁻⁵¹⁻⁵⁴⁻⁵⁶ Recently, the Boston University group published their experience of CTE, with 80 athlete brain donors (22 of whom were also military veterans) with a history of repetitive brain injury, and they found that 80% of these cases demonstrated the characteristic pathology for CTE.⁸

Signs and symptoms

The modern CTE description suggests that symptoms such as gait disorders, speech slowing and extrapyramidal signs may be present; however, neuropsychiatric and behavioural symptoms tend to predominate early.⁵⁷ The most common symptoms reported are mood disorder (mainly depression), paranoia, agitation, social withdrawal, poor judgement and aggression. Cognitive impairment tends to emerge as the major neuropsychiatric feature in the latter stages⁵⁸⁻⁵⁹ and typically includes impairment across the domains of orientation, memory,

language, attention, information processing speed and executive functioning.⁶⁰ These cognitive symptoms have been proposed to progress in a somewhat predictable manner,³ whereas the classic CTE entity reported little progression of cognitive deficits.⁴⁶

Neuropathology

The reported neuropathological characteristics of both entities (classic and modern CTE) appear to be more closely related and share a number of common features such as fenestrated septum pellucidum, cerebral atrophy, tau+NFT inclusion (although found in greater amounts in modern CTE), β -amyloid deposition (found in less amounts in modern CTE), reduced pigmentation of the substantia nigra and locus caeruleus, and enlarged ventricles. The qualitative description of modern CTE characterises the neuropathological change as also including fronto-temporal lobe atrophy with extensive tau pathology distributed throughout the neocortex, medial temporal lobe, diencephalon, brainstem and spinal cord. In the 51 cases reviewed by Gavett *et al*⁷ 61 diffuse amyloid plaques were found in 24 (47%), neuritic amyloid plaques in 13 (27%) and amyloid angiopathy in 3 (6%). In the Boston study,⁸ amyloid deposition was noted in 44% of the CTE cases.

Although a number of similarities in neuropathological findings have been reported across the modern CTE cases, there are also some important differences. McKee *et al*³ have reported a marked accumulation of tau-immunoreactive astrocytes, but this has not been observed in any of the cases examined by Omalu *et al*.⁶² Omalu *et al*⁶³ have reported the skip-phenomenon, with a propensity to lobar cortical distribution in the absence of prominent periventricular topographic distribution, which has not been described in the McKee *et al*³ cases.

A defining neuropathological feature of the modern CTE entity is abundant filamentous tau lesions, of which the patterns of expression are considered unique, occurring in the absence (or at least relative scarcity) of β -amyloid deposits.³ 8 Prominent filamentous tau inclusions and brain degeneration in the absence of β -amyloid deposits are also considered the sine qua non of a number of neurodegenerative tauopathies. It is worth noting that a number of the degenerative tauopathies, such as behavioural variant fronto-temporal dementia, share many of the clinical and pathological features of 'modern' CTE.^{64–69}

DISCUSSION

The recent autopsy cases differ from the classic CTE description across a number of characteristics including age of onset, progression, assumed predominate (clinical) features and diagnostic criteria. It has also been reported that these recent cases of modern CTE, speculated to be a consequence of concussive and subconcussive blows, are characterised by a distinct neuropathological profile³ 61 63 and manifest primarily as a tauopathy.⁸ Although many of the reported macroscopic neuropathological features are common among the original and newer descriptions of CTE, the different distribution of tau-immunoreactive astrocytes distinguishes the newer description with preferential involvement of the superficial cortical layers occurring on a background of relative scarcity of β -amyloid plaques.

At present, there are no published epidemiological, cross-sectional or prospective studies relating to modern CTE. Owing to the nature of the published studies, being case reports or pathological case series, it is not possible to determine the causality or risk factors with any certainty. As such, the speculation that repeated concussion or subconcussive impacts cause CTE remains unproven.⁷⁰

The extent to which age-related changes, psychiatric or mental health illness, alcohol/drug use or coexisting dementing illnesses contribute to this process is largely unaccounted for in the published literature. In addition, consideration for the potential genetic risk in those athletes with a family history of neurodegenerative disease and the extent to which this contributes to the clinical and pathological profiles also require further investigation.⁴¹

SUMMARY

At present, the interpretation of causation in the modern CTE case studies should proceed cautiously. The causal assumptions require further prospective or longitudinal studies on the topic. Ultimately, scientific research might establish that participation in contact sports leads to a distinct neuropathological syndrome, and this neuropathology causes psychiatric, cognitive and physical problems, but this cause and effect relationship remains to be shown scientifically.

Author affiliations

- ¹The Florey Institute of Neuroscience and Mental Health, Heidelberg, Australia
²Faculty of Kinesiology, Sport Injury Prevention Research Centre, Calgary, Alberta, Canada
³Faculty of Medicine, Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada
⁴Department of Neurology, University of Michigan, Ann Arbor, Michigan, USA
⁵Burke Rehabilitation Hospital, White Plains, New York, USA
⁶Neuropsychiatry Service, HNE Mental Health Service, Newcastle, New South Wales, Australia

Competing interests See the supplementary online data for competing interests (<http://dx.doi.org/10.1136/bjsports-2013-092248>).

Provenance and peer review Commissioned; internally peer reviewed.

REFERENCES

- Gregory S. The problem with football. Our favorite sport is too dangerous. How to make the game safer. *Time* 2010;175:36–43.
- Guskiewicz KM, Marshall SW, Bailes J, *et al*. Association between recurrent concussion and late-life cognitive impairment in retired professional football players. *Neurosurgery* 2005;57:719–26.
- McKee AC, Cantu RC, Nowinski CJ, *et al*. Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury. *J Neuropathol Exp Neurol* 2009;68:709–35.
- Omalu BI, Hamilton RL, Kambh MI, *et al*. Chronic traumatic encephalopathy (CTE) in a National Football League Player: case report and emerging medicolegal practice questions. *J Forensic Nurs* 2010;6:40–6.
- Cantu RC. Chronic traumatic encephalopathy in the National Football League. *Neurosurgery* 2007;61:223–5.
- McCroly P. Sports concussion and the risk of chronic neurological impairment. *Clin J Sport Med* 2011;21:6–12.
- Gavett BE, Cantu RC, Shenton M, *et al*. Clinical appraisal of chronic traumatic encephalopathy: current perspectives and future directions. *Curr Opin Neurol* 2011;24:525–31.
- McKee AC, Stein TD, Nowinski CJ, *et al*. The spectrum of disease in chronic traumatic encephalopathy. *Brain* 2012;136:43–64.
- Chen JK, Johnston KM, Collie A, *et al*. A validation of the post concussion symptom scale in the assessment of complex concussion using cognitive testing and functional MRI. *J Neurol, Neurosurg Psychiatry* 2007;78:1231–8.
- Schwenk TL, Gorenflo DW, Dopp RR, *et al*. Depression and pain in retired professional football players. *Med Sci Sports Exercise* 2007;39:599–605.
- Makdissi M, Darby D, Maruff P, *et al*. Natural history of concussion in sport: markers of severity and implications for management. *Am J Sports Med* 2010;38:464–71.
- King N. Mild head injury: neuropathology, sequelae, measurement and recovery. *Br J Clin Psychol/Br Psychol Soc* 1997;36(Pt 2):161–84.
- Sullivan KA, Edmed SL, Cunningham LC. A comparison of new and existing mild traumatic brain injury vignettes: recommendations for research into post-concussion syndrome. *Brain Inj* 2013;27:19–30.
- King NS, Kirwilliam S. Permanent post-concussion symptoms after mild head injury. *Brain Inj* 2011;25:462–70.
- Guskiewicz KM, Marshall SW, Bailes J, *et al*. Recurrent concussion and risk of depression in retired professional football players. *Med Sci Sports Exercise* 2007;39:903–9.

- 16 De Beaumont L, Theoret H, Mongeon D, *et al.* Brain function decline in healthy retired athletes who sustained their last sports concussion in early adulthood. *Brain* 2009;132(Pt 3):695–708.
- 17 Kraus MF, Susmaras T, Caughlin BP, *et al.* White matter integrity and cognition in chronic traumatic brain injury: a diffusion tensor imaging study. *Brain* 2007;130(Pt 10):2508–19.
- 18 Giza CC, Hovda DA. The neurometabolic cascade of concussion. *J Athl Train* 2001;36:228–35.
- 19 Metting Z, Rodiger LA, De Keyser J, *et al.* Structural and functional neuroimaging in mild-to-moderate head injury. *Lancet Neurol* 2007;6:699–710.
- 20 Zhang L, Heier LA, Zimmerman RD, *et al.* Diffusion anisotropy changes in the brains of professional boxers. *Am J Neuroradiol* 2006;27:2000–4.
- 21 Chen JK, Johnston KM, Collie A, *et al.* Behavioural and functional imaging outcomes in symptomatic concussed athletes measured with cogsport and functional MRI. *BJ Sport Med* 2004;38:659.
- 22 Waites AB, Briellmann RS, Saling MM, *et al.* Functional connectivity networks are disrupted in left temporal lobe epilepsy. *Ann Neurol* 2006;59:335–43.
- 23 Vagnozzi R, Signoretti S, Tavazzi B, *et al.* Temporal window of metabolic brain vulnerability to concussion: a pilot 1H-magnetic resonance spectroscopic study in concussed athletes—part III. *Neurosurgery* 2008;62:1286–95.
- 24 Garnett MR, Blamire AM, Corkill RG, *et al.* Early proton magnetic resonance spectroscopy in normal-appearing brain correlates with outcome in patients following traumatic brain injury. *Brain* 2000;123(Pt 10):2046–54.
- 25 Harvey RJ, Skelton-Robinson M, Rossor MN. The prevalence and causes of dementia in people under the age of 65 years. *J Neurol Neurosurg Psychiatry* 2003;74:1206–9.
- 26 Bazarian JJ, Cernak I, Noble-Haesslein L, *et al.* Long-term neurologic outcomes after traumatic brain injury. *J Head Trauma Rehabil* 2009;24:439–51.
- 27 Gonzalez HM, Tarraf W, Whitfield KE, *et al.* The epidemiology of major depression and ethnicity in the United States. *J Psychiatr Res* 2010;44:1043–51.
- 28 Olin JT, Katz IR, Meyers BS, *et al.* Provisional diagnostic criteria for depression of Alzheimer disease: rationale and background. *Am J Geriatr Psychiatry* 2002;10:129–41.
- 29 Steinberg EM, Rubinow DR, Bartko JJ, *et al.* A cross-sectional evaluation of perimenopausal depression. *J Clin Psychiatry* 2008;69:973–80.
- 30 Lehmann EJ, Hein MJ, Baron SL, *et al.* Neurodegenerative causes of death among retired National Football League players. *Neurology* 2012;79:1–5.
- 31 McKee AC, Gavett BE, Stern RA, *et al.* TDP-43 proteinopathy and motor neuron disease in chronic traumatic encephalopathy. *J Neuropathol Exp Neurol* 2010;69:918–29.
- 32 White L. Brain lesions at autopsy in older Japanese-American men as related to cognitive impairment and dementia in the final years of life: a summary report from the Honolulu-Asia aging study. *J Alzheimers Dis* 2009;18(Pt 3):713–25.
- 33 Robinson JL, Geser F, Corrada MM, *et al.* Neocortical and hippocampal amyloid-beta and tau measures associate with dementia in the oldest-old. *Brain* 2011;134:3708–15.
- 34 Gelber RP, Launer LJ, White LR. The Honolulu-Asia Aging Study: epidemiologic and neuropathologic research on cognitive impairment. *Curr Alzheimer Res* 2012;9:664–72.
- 35 Snowden JS, Gibbons ZC, Blackshaw A, *et al.* Social cognition in frontotemporal dementia and Huntington's disease. *Neuropsychologia* 2003;41:688–701.
- 36 Sonnen JA, Larson EB, Crane PK, *et al.* Pathological correlates of dementia in a longitudinal, population-based sample of aging. *Ann Neurol* 2007;62:406–13.
- 37 Da Cruz S, Cleveland DW. Understanding the role of TDP-43 and FUS/TLS in ALS and beyond. *Curr Opin Neurobiol* 2011;21:904–19.
- 38 Scheltens P, Rockwood K. How golden is the gold standard of neuropathology in dementia? *Alzheimers Demen* 2011;7:486–9.
- 39 Martland HS. Punch drunk. *J Am Med Assoc* 1928;19:1103–7.
- 40 Roberts AH. *Brain damage in boxers: a study of the prevalence of traumatic encephalopathy among ex-professional boxers.* London: Pitman, 1969.
- 41 McCrory P. Boxing and the risk of chronic brain injury. *BMJ* 2007;335:781–2.
- 42 Johnson J. Organic psychosyndromes due to boxing. *Br J Psychiatr* 1969;115:45–53.
- 43 Johnson J. The EEG in the traumatic encephalopathy of boxers. *Psychiatr Clin* 1969;2:204–11.
- 44 Roberts GW, Allsop D, Bruton C. The occult aftermath of boxing. *J Neurol Neurosurg Psychiatry* 1990;53:373–8.
- 45 Nowak LA, Smith GG, Reyes PF. Dementia in a retired world boxing champion: case report and literature review. *Clin Neuropathol* 2009;28:275–80.
- 46 Corsellis JA, Bruton CJ, Freeman-Browne D. The aftermath of boxing. *Psychol Med* 1973;3:270–303.
- 47 De Calignon A, Polydoro M, Suarez-Calvet M, *et al.* Propagation of tau pathology in a model of early Alzheimer's disease. *Neuron* 2012;73:685–97.
- 48 Liu L, Drouet V, Wu JW, *et al.* Trans-synaptic spread of tau pathology in vivo. *PLoS ONE* 2012;7:e31302.
- 49 Frost B, Jacks RL, Diamond MI. Propagation of tau misfolding from the outside to the inside of a cell. *J Biol Chem* 2009;284:12845–52.
- 50 Clavaguera F, Goedert M, Tolnay M. Induction and spreading of tau pathology in a mouse model of Alzheimer's disease. *Med Sci* 2010;26:121–4.
- 51 Omalu BI, Bailes J, Hammers JL, *et al.* Chronic traumatic encephalopathy, suicides and parasuicides in professional American athletes: the role of the forensic pathologist. *Am J Forensic Med Pathol* 2010;31:130–2.
- 52 Clausen H, McCrory P, Anderson V. The risk of chronic traumatic brain injury in professional boxing: change in exposure variables over the past century. *Br J Sports Med* 2005;39:661–4.
- 53 Ahmed OH, Sullivan SJ, Schneiders AG, *et al.* iSupport: do social networking sites have a role to play in concussion awareness? *Disabil Rehabil* 2010;32:1877–83.
- 54 Omalu BI, DeKosky ST, Hamilton RL, *et al.* Chronic traumatic encephalopathy in a national football league player: part II. *Neurosurgery* 2006;59:1086–92; discussion 92–3.
- 55 Omalu BI, DeKosky ST, Minster RL, *et al.* Chronic traumatic encephalopathy in a National Football League player. *Neurosurgery* 2005;57:128–34; discussion 34..
- 56 Omalu BI, Fitzsimmons RP, Hammers J, *et al.* Chronic traumatic encephalopathy in a professional American wrestler. *J Forensic Nurs* 2010;6:130–6.
- 57 Jordan B. Neurologic aspects of boxing. *Arch Neurol* 1987;44:453–9.
- 58 Guterman A, Smith RW. Neurological sequelae of boxing. *Sports Med (Auckland, NZ)* 1987;4:194–210.
- 59 Jordan BD. Chronic traumatic brain injury associated with boxing. *Semin Neurol* 2000;20:179–85.
- 60 Costanza A, Weber K, Gandy S, *et al.* Review: contact sport-related chronic traumatic encephalopathy in the elderly: clinical expression and structural substrates. *Neuropathol Appl Neurobiol* 2011;37:570–84.
- 61 Gavett BE, Stern RA, McKee AC. Chronic traumatic encephalopathy: a potential late effect of sport-related concussive and subconcussive head trauma. *Clin Sports Med* 2011;30:179–88, xi.
- 62 Omalu B, Hammers JL, Bailes J, *et al.* Chronic traumatic encephalopathy in an Iraqi war veteran with posttraumatic stress disorder who committed suicide. *Neurosurg Focus* 2011;31:E3.
- 63 Omalu B, Bailes J, Hamilton RL, *et al.* Emerging histomorphologic phenotypes of chronic traumatic encephalopathy in American athletes. *Neurosurgery* 2011;69:173–83; discussion 83..
- 64 Rascovsky K, Salmon DP, Ho GJ, *et al.* Cognitive profiles differ in autopsy-confirmed frontotemporal dementia and AD. *Neurology* 2002;58:1801–8.
- 65 Rascovsky K, Hodges JR, Knopman D, *et al.* Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* 2011;134(Pt 9):2456–77.
- 66 Rascovsky K, Hodges JR, Kipps CM, *et al.* Diagnostic criteria for the behavioral variant of frontotemporal dementia (bvFTD): current limitations and future directions. *Alzheimer Dis Assoc Disord* 2007;21:S14–8.
- 67 Hodges JR. Frontotemporal dementia (Pick's disease): clinical features and assessment. *Neurology* 2001;56(11 Suppl 4):S6–10.
- 68 Hodges JR, Davies RR, Xuereb JH, *et al.* Clinicopathological correlates in frontotemporal dementia. *Ann Neurol* 2004;56:399–406.
- 69 Hodges JR, Miller B. The classification, genetics and neuropathology of frontotemporal dementia. Introduction to the special topic papers: part I. *Neurocase* 2001;7:31–5.
- 70 Gaetz W, Macdonald M, Cheyne D, *et al.* Neuromagnetic imaging of movement-related cortical oscillations in children and adults: age predicts post-movement beta rebound. *Neuroimage* 2010;51:792–807.

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
CONSENSUS STATEMENT WRITING GROUP		
Paul McCrory MBBS PhD	Associate Professor, The Florey Institute of Neuroscience and Mental Health, Heidelberg, Australia	Paul McCrory is a co-investigator, collaborator, or consultant on grants relating to mild TBI funded by several governmental organizations. He is Co-Chair of the Australian Centre for Research into Sports Injury and its Prevention (ACRISP), which is one of the International Research Centres for Prevention of Injury and Protection of Athlete Health supported by the International Olympic Committee (IOC). He has a clinical and consulting practice in general and sports neurology. He receives book royalties from McGraw-Hill and was employed in an editorial capacity by the British Medical Journal Publishing Group from 2001-2008. He has been reimbursed by the government, professional scientific bodies, and sporting bodies for presenting research relating to mild TBI and sport-related concussion at meetings, scientific conferences, and symposiums. He received consultancy fees in 2010 from Axon Sports (US) for the development of educational material (which was not renewed) and has received research funding since 2005 from CogState Inc. He is a cofounder and shareholder in two biomedical companies (involved in eHealth and Compression garment technologies) but does not hold any individual shares in any company related to concussion or brain injury assessment or technology. He did not receive any form of financial support directly related to this manuscript.
Willem H. Meeuwisse, MD, PhD	Professor and Co-Chair, Sport Injury Prevention Research Centre, Faculty of Kinesiology and Hotchkiss Brain Institute, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada	Willem Meeuwisse, MD, PhD has received research grant support through the University of Calgary from the Canadian Institutes of Health Research, Alberta Innovates Health Solutions, the International Football Association (FIFA), Alberta Children’s Hospital Research Institute and the Hotchkiss Brain Institute. He is Co-Chair of the Sport Injury Prevention Research Centre, which is one of the International Research Centres for Prevention of Injury and Protection of Athlete Health supported by the International Olympic Committee (IOC). He has a clinical and consulting practice in sport medicine at the University of Calgary Sport Medicine Centre with a focus on sport related concussion. He has received travel funding from FIFA and the Medical Commission of the IOC. From 1998-2012 he was the Editor-in-Chief of the Clinical Journal of Sport Medicine and received editorial support funding from Lippincott-Wolters Kluwer. He receives compensation as a medical consultant to the National Hockey League, may receive royalty from BKIN Technologies, and is a shareholder of Safebrain Canada and PrivIT Healthcare. He did not

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
		receive any form of financial support directly related to this manuscript.
Mark Aubry MD, Dip.Sport Med.	Chief Medical Officer International Ice Hockey Federation, Switzerland; Member of the IOC Medical Commission Games Group , Co-Director Ottawa Sport Medicine Centre, Ottawa, Ontario, Canada	Mark Aubry M.D.Dip Sport Med. receives travel funding from the IIHF and IOC Medical Commission Games Group for meetings, IIHF Championships and Olympic Games. He did not receive any form of financial support directly related to this manuscript.
Robert C. Cantu, M.A., M.D.	Clinical Professor, Neurosurgery and Co-Director Center for the Study of Traumatic Encephalopathy, Boston University Medical Center, Boston, MA USA	Robert C. Cantu, M.A., M.D serves as Vice President, National Operating Committee on Standards for Athletic Equipment (NOCSAE), Co-Founder and Chairman Medical Director Sports Legacy (SLI), Waltham, MA, USA; Senior Advisor to NFL’s Head, Neck and Spine Committee, Expert witness Trial Testimony. He did not receive any form of financial support directly related to this manuscript.
Jiří Dvořák, MD, PhD	Professor of Neurology at the University of Zurich, Senior consultant Schulthess Clinic Zurich, Switzerland, Chairman F-MARC (FIFA Medical Assessment and Research Center)	Jiri Dvorak, MD, PhD, received compensation from his work at FIFA as FIFA Chief medical officer, no other compensations or grants have been received. He did not receive any form of financial support directly related to this manuscript.
Ruben J. Echemendia, Ph.D	Psychological and Neurobehavioral Associates, Inc., State College, PA, USA, Adjunct Associate Professor of Psychology, University of Missouri – Kansas City, USA	Ruben J. Echemendia, Ph.D. receives financial compensation as a consultant to the National Hockey League, Major League Soccer, and the U.S. Soccer Federation. He also receives financial compensation as a consultant to Princeton University and has served as a PI or co-PI on grants funded by NOCSAE, AMSSM, and the New Jersey Commission on Brain Injury Research. Dr. Echemendia maintains a private practice focused on clinical neuropsychology, sports neuropsychology, and forensic applications of neuropsychology. He did not receive any form of financial support directly related to this manuscript.
Lars Engebretsen MD PhD	Professor, Department of Orthopaedic Surgery, Oslo University Hospital and	Lars Engebretsen, MD, PhD has received research grant support through the University of Oslo from the Norwegian Institutes of Health Research, The Health South East, the

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
	Faculty of Medicine, University of Oslo, Norway. Co-chair Oslo Sports Trauma Research Center, Oslo, Norway. Head Scientific Activities, International Olympic Committee, Lausanne, Switzerland	International Football Association (FIFA), He is Co-Chair of the Sport Injury Prevention Research Centre, which is one of the International Research Centres for Prevention of Injury and Protection of Athlete Health supported by the International Olympic Committee (IOC). He is the professor and Chair of the Oslo University Orthopaedic department. He has been Editor-in-Chief of the BJSM IPHP From 2008. From January 2012 he has been co-editor of The Journal of Bone and Joints. He did not receive any form of financial support directly related to this manuscript.
Karen M. Johnston, MD, Ph.D	Neurosurgeon, Division of Neurosurgery, University of Toronto; Concussion Management Program Athletic Edge Sports Medicine, Toronto, Canada	Karen M. Johnston MD, Ph.D is a neurosurgeon subspecialized in brain trauma and has a clinical and research focus in sport concussion. She has received research funding through the CIHR, ONF and Innovation Fund, American College of Surgeons, McGill University, Pashby Sport Safety Fund and Think First Canada. She has independently consulted widely with many sport organizations however she has not and has never had a formal reimbursed affiliation with any sport team or sport governing body. She did not receive any form of financial support directly related to this manuscript.
Jeffrey S. Kutcher, MD	Associate Professor and Director, Michigan NeuroSport, Departement of Neurology, University of Michigan, Ann Arbor, MI, USA	Jeffrey S. Kutcher, MD is the Director of Michigan NeuroSport, the University of Michigan's academic sports neurology program. He has received research grant support form the National Collegiate Athletic Association and EIMindA, Ltd. He is the Director of the National Basketball Association's Concussion Program and is a paid consultant for the National Hockey League Players' Association. He is a current member of the National Football League Players' Association's Mackey-White TBI Committee. He is also a consulting neurologist for the United States Ski and Snowboard Association. He was a founding member and the first Chairperson of the Sports Neurology Section of the American Academy of Neurology. He did not receive any form of financial support directly related to this manuscript.
Martin Raftery MB BS FACSP	Chief Medical Officer, International Rugby Board, Dublin, Ireland	No disclosures
Allen Sills, MD, FACS	Associate Professor of Neurosurgery, Orthopaedic Surgery and	Allen Sills, MD, FACS serves on the medical board of the Federation Equestre Internationale (FEI) and has received travel funding from that organization. He

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
	Rehabilitation; Vanderbilt Sports Concussion Center, Vanderbilt University Medical Center, Nashville, Tennessee, USA	also serves as unpaid neurosurgical consultant to the US Equestrian Foundation, the Southeastern Conference (SEC) Working Group on Concussion, and the Nashville Predators (NHL) hockey team. He has a clinical practice in neurosurgery at the Vanderbilt University Medical Center with a focus on neurological problems in athletes. He did not receive any form of financial support directly related to this manuscript.
CONSENSUS STATEMENT COAUTHORS		
Brian W. Benson, MD, PhD	Adjunct Research Assistant Professor, Department of Clinical Neurosciences, Faculty of Medicine; Clinical Assistant Professor, Department of Family Medicine; and Sport Medicine Centre, Faculty of Kinesiology, University of Calgary, Calgary, Alberta, Canada	Brian W. Benson, MD, PhD has a clinical and consulting practice in sport medicine at the University of Calgary Sport Medicine Centre with a focus on sport-related concussion. He also serves as Director of the Sport Concussion Clinic at the University of Calgary Sport Medicine Centre and may receive royalty from BKIN Technologies. He did not receive any form of financial support directly related to this manuscript.
Gavin A. Davis, MB.BS	Department of Neurosurgery, Austin and Cabrini Hospitals & The Florey Institute of Neuroscience and Mental Health, Melbourne, Victoria, Australia	Gavin A. Davis, MB.BS. has received travel funding from FIFA. He did not receive any form of financial support directly related to this manuscript
Kevin M. Guskiewicz, PhD, ATC 	Kenan Distinguished Professor and Chair, Department of Exercise and Sport Science; Co-Director, Matthew Gfeller Sport-Related Traumatic Brain Injury Research Center, University of North Carolina at Chapel Hill, Chapel Hill, NC; USA	Kevin Guskiewicz, PhD, ATC, serves on the NCAA's Health and Safety Advisory Committee for Concussion, the NFL's Head Neck and Spine Committee, and the NFLPA's Mackey-White Committee; has received funding for travel and honoraria for lectures on sports concussion for professional organizations, including the IOC and FIFA; has given expert testimony on traumatic brain injury/concussion cases; and has received research funding from the National Institutes of Health, Centers for Disease Control and Prevention, National Operating Committee for Standards in Athletic Equipment, National Collegiate Athletic Association, NFL Charities, NFL Players' Association, USA Hockey, and National Athletic Trainers' Association. He did not receive any form of financial support directly related to this manuscript.

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
Stanley A. Herring, MD	Clinical Professor Departments of Rehabilitation Medicine, Orthopaedics and Sports Medicine and Neurological Surgery, University of Washington; Co-Medical Director Seattle Sports Concussion Program, Team Physician Seattle Seahawks and Seattle Mariners, Seattle, Washington, USA	Stanley A. Herring, MD serves as a Member National Football League Head, Neck and Spine Committee; Member Football and Wellness Committee USA Football; Member Medical Advisory Board Pop Warner Football; Chair of Medical Advisory Board X2IMPACT, a non-financial position. Clinical and consulting practice at Harborview Medical Center focusing on the diagnosis and management of neurological and musculoskeletal injuries with particular interest in spinal disorders in active people and athletes and sports-related concussions. No form of financial support directly related to this manuscript was received
Grant Iverson PhD	Neuropsychology Outcome Assessment Laboratory; Professor, Department of Psychiatry, University of British Columbia, Vancouver, BC, Canada	Grant Iverson, Ph.D. has been reimbursed by the government, professional scientific bodies, and commercial organizations for discussing or presenting research relating to mild TBI and sport-related concussion at meetings, scientific conferences, and symposiums. He has a clinical and consulting practice in forensic neuropsychology involving individuals who have sustained mild TBIs (including professional athletes). He has received research funding from several test publishing companies, including ImPACT Applications, Inc., CNS Vital Signs, and Psychological Assessment Resources (PAR, Inc.). He did not receive any form of financial support directly related to this manuscript.
Barry D. Jordan MD, MPH	Associate Professor of Clinical Neurology at Weill Medical College of Cornell University. New York, New York USA. Assistant Medical Director of the Burke Rehabilitation Hospital, White Plains, New York. USA. Chief Medical Officer of the New York State Athletic Commission. New York, New York USA. National Football League Players Association (NFLPA) Mackey-White Traumatic Brain Injury Committee.	Dr. Jordan has received financial compensation from the National Football League (NFL) Benefits Association for the establishment of a Neurocognitive Disability Benefit for retired players. He also has received financial compensation from Innovative CEUs for developing a medical-based high school coaching education program. In his position as the Chief Medical Officer of the New York State Athletic Commission, Dr. Jordan is a governor's appointed public officer of the State of New York. He maintains an inpatient and outpatient clinical practice at the Burke Rehabilitation Hospital specializing in traumatic brain injury, dementia and sports neurology. Dr. Jordan did not receive any financial support related to this manuscript.
James Kissick MD, CCFP, Dip Sport Med	Sport Medicine Physician, Ottawa Sport Medicine Centre, Ottawa, Canada; Clinical	James Kissick has a clinical and consulting practice in sport and exercise medicine at the Ottawa Sport Medicine Centre. He did not receive any form of financial support directly

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
	Lecturer, Dept of Family Medicine, University of Ottawa, Ottawa, Canada. Team Physician, Canadian National Men's Sledge Hockey Team.	related to this manuscript.
Michael McCrea, PhD, ABPP	Professor & Director of Brain Injury Research, Departments of Neurosurgery and Neurology, Medical College of Wisconsin, WI, USA	Michael McCrea, PhD has been funded by the United States Department of Defense, U.S. Defense Veterans Brain Injury Center (DVBIC), NFL Charities, National Collegiate Athletic Association (NCAA), National Operating Committee on Standards for Athletic Equipment (NOCSAE), National Federation of High School Associations (NFHS), and other funding organizations over the past decade. Dr. McCrea is a member of the NFL Head, Neck and Spine Committee and has served on several U.S. national panels for scientific review and policy making relevant to sport-related concussion. He has been funded by BrainScope, Inc. for prior studies, but receives no direct compensation from BrainScope or other industry entities or sports organizations. Dr. McCrea is co-author of the Standardized Assessment of Concussion (SAC). He receives no financial compensation or support from publication or distribution of the SAC. He is currently principal investigator (PI) on a study by the U.S. Department of Defense comparing several computerized neurocognitive assessment tools for the assessment of sport-related concussion. He did not receive any form of financial support directly related to this manuscript.
Andrew S McIntosh, BAppSci (PT), MBiomedE, PhD	Adjunct Associate Professor Australian Centre for Research into Injury in Sports and its Prevention, Monash Injury Research Institute, Monash University, Adjunct Associate Professor, Transport and Road Safety Research, Faculty of Science, the University of New South Wales, Director, McIntosh Consultancy and Research Pty Ltd. Sydney, Australia	Andrew McIntosh in a self-employed consultant. In the area of sport and sports injury research he has received grant support through UNSW from the Australian Research Council, NSW Racing, the IRB, the ARU, and the AFL. As a consultant in the area of sport and sports injury he has undertaken paid work for the Australian Racing Board, McLaren F1, the IRB and Albion. He has received travel funding from the International Rugby Board, IIHF, FIFA and the Medical Commission of the IOC. He did not receive any form of financial support directly related to this manuscript.
David L. Maddocks, LLB, PhD	Barrister & Solicitor, Perry Maddocks Trollope Lawyers, Melbourne, Australia.	David Maddocks, LLB, PhD has received research grant support through the Australian Football League (AFL). He provides consultancy medico-legal advice to the AFL. He has

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
	Neuropsychologist in private practice.	received travel funding from FIFA. He did not receive any form of financial support directly related to this manuscript.
Michael Makdissi, MBBS, PhD	Research Fellow, The Florey Institute of Neuroscience and Mental Health, Melbourne Brain Centre, Austin Campus, Melbourne, Australia. Research Fellow, Centre For Health Exercise and Sports Medicine, Melbourne Physiotherapy Department, University of Melbourne, Melbourne, Australia	Michael Makdissi, MBBS, PhD, has received research grant support through The Florey Institute of Neurosciences and Mental Health from the National Health and Medical Research Council, the Australian Football League (AFL) and AFL Medical Officers Association. He has received travel funding from FIFA. He has a clinical and consulting practice in sports medicine. He did not receive any form of financial support directly related to this manuscript.
Laura Purcell, MD, FRCPC	Associate Clinical Professor, Department of Pediatrics, McMaster University, Hamilton ON, Canada; Pediatric Sport Medicine Physician, David Braley Sport Medicine and Rehabilitation Centre, McMaster University, Hamilton ON, Canada; President, Paediatric Sport and Exercise Medicine Section, CPS	No disclosures
Margot Putukian, MD	Director of Athletic Medicine, Head Team Physician, Princeton University, Associate Clinical Professor, Robert Wood Johnson, University of Medicine and Dentistry of New Jersey (UMDNJ), USA	Margot Putukian MD has received research grant support through the new Jersey Commission on Brain Injury, the American Medical Society for Sports Medicine (AMSSM) and the National Operating Committee on Standards for Athletic Equipment (NOCSAE). She is the chair of the US Lacrosse Sports Science & Safety Committee and also serves on the National Football League (NFL) Head Neck and Spine Committee, chairing the Return to Play subcommittee for the NFL. She works as a team physician for Princeton University, as well as team physician for the US Men's National Lacrosse Team and for US Soccer. She is a consultant on concussion for the Center for Disease Controls (CDC) and the National Collegiate Athletic Association (NCAA). She is a past president for the AMSSM and currently is president of the AMSSM Foundation. Dr. Putukian does not receive any financial

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
		compensation for her work with the AMSSM, NFL, US Lacrosse, CDC or NCAA. She did not receive any form of financial support directly related to this manuscript.
Kathryn Schneider, PT, PhD	Sport Injury Prevention Research Centre, University of Calgary, Calgary, Alberta, Canada	Kathryn Schneider, PT, PhD has received research grant support through the University of Calgary from the Alberta Centre for Child, Family and Community Research. She has a clinical and consulting practice in physiotherapy at Evidence Sport and Spinal and Centric Health with a focus on sport related concussion. She did not receive any form of financial support directly related to this manuscript.
Charles H. Tator, MD, PHD	Professor of Neurosurgery, Toronto Western Hospital and University of Toronto, Canada; Senior Scientist, Krembil Neuroscience Centre, Toronto, Canada; Founder, ThinkFirst Canada; Board Member, Parachute Canada.	Charles Tator is a consultant to ImpaKt Protective, and is co-chair of the Concussion Awareness and Recognition Committee of the Ontario Neurotrauma Foundation. He is Chair of ThinkFirst's Concussion Education and Awareness Committee, and Project Director of the Canadian Sports Concussion Project at the Krembil Neuroscience Centre. He is Director of the Ontario Concussion Centres Consortium. He has received research grants from many public sources including the Canadian Institute for Health Research, and the Ontario Brain Institute. He did not receive any financial support related to this manuscript.
Michael Turner MB BS	Chief Medical Adviser, British Horseracing Authority, London, United Kingdom	Michael Turner is employed as Chief Medical Adviser by the British Horseracing Authority and the Lawn Tennis Association. He has received travel funding from FIFA, the IOC, the ACSM and Gruppoforte (Physiomind) for attending conferences and lecturing. In addition to travel funding, most conference organisers also provided free Registration and support for accommodation. He did not receive any form of financial support directly related to this manuscript.
OTHER REVIEW AUTHORS		
Mick Molloy MD	Department of Medicine, University College, Cork, Ireland	No disclosures
Johna Register-Mihalik PhD	Senior Research Associate, Emergency Services Institute, WakeMed Health and Hospitals, Raleigh, NC; USA; Adjunct	Johna Register-Mihalik, PhD, LAT, ATC serves on the Board of Directors for the Brain Injury Association of North Carolina; has received funding for travel and honoraria for lectures on concussion in sport for professional organizations and has received funding from the

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
	Assistant Professor, Department of Exercise and Sport Science, University of North Carolina at Chapel Hill, Chapel Hill, NC; USA	National Operating Committee for Standards in Athletic Equipment, National Football League Charities, and center grant funding from the National Institutes of Health (North Carolina Translational and Clinical Science Institute). She did not receive any form of financial support directly related to this manuscript.
Stephen A Macciocchi PhD	Shepherd Center, Atlanta, Georgia, USA	No disclosures
Gerard A Gioia PhD	Children’s National Medical Center & George Washington University School of Medicine, Rockville, Maryland, USA	Gerard A. Gioia, Ph.D. has received research grant support through the CDC and NIH as well as the USA Hockey Foundation. He serves as a consultant to multiple school systems in the Baltimore-Washington region. Dr. Gioia maintains a clinical practice focused on clinical neuropsychology and sports neuropsychology. He receives royalties as a co-author of the Behavior Rating Inventory of Executive Function (BRIEF) and Tasks of Executive Control (TEC). He is a co-author of the Pediatric ImPACT test battery, funded through the CDC, and has not received direct financial compensation for its development. He did not receive any form of financial support directly related to this manuscript.
Paul Comper PhD	Kinesiology and Physical Education & Graduate Department of Rehabilitation Science, University of Toronto, Ontario, Canada	PC receives financial compensation as a consultant to the National Hockey League Players’ Association. He did not receive any form of financial support directly related to this manuscript.
Steven P. Broglio PhD	Michigan NeuroSport, School of Kinesiology, University of Michigan, MI, USA	Dr Broglio is a member of Michigan NeuroSport and currently has research support from EIMindA, Ltd, the National Athletic Trainers’ Association – Research and Education Foundation, the National Institutes of Health – National Institute of Neurological Disorders and Stroke, National Collegiate Athletic Association, and the Women’s Sports, Health and Activity Research and Policy (SHARP) Center. These funds were not used to support this manuscript.
Alain Ptito PhD	Department of Psychology, McGill University Health Centre, Cognitive Neuroscience Unit, Montreal Neurological Institute and Hospital & Department of Neurology and Neurosurgery, McGill	Alain Ptito is Director of the Department of Psychology at the MUHC and Professor of Neurology and Neurosurgery at McGill University. He has received research grants from several public sources including the Canadian Institute for Health Research, and the Ontario Neurotrauma Foundation. He is also a consultant neuropsychologist specializing in traumatic brain injury/concussion in athletes and motor vehicle accident victims. He did

Affiliation and disclosure statement for all authors in the BJSM 47:5 April IPHP issue

NB: All panelists received travel funding and accommodation from the Concussion in Sport Group to attend the Consensus meeting in Zurich, Switzerland

Name	Affiliations	Disclosures
	University, Montreal, Canada.	not receive any financial support related to this manuscript.
Carolyn A Emery PhD	Faculty of Kinesiology, Sport Injury Prevention Research Centre, University of Calgary, Calgary, Alberta, Canada	CAE is supported by a Population Health Investigator Award from the Alberta Innovates Health Solutions. She also holds a Professorship in Pediatric Rehabilitation funded through the Alberta Children’s Hospital Foundation. She co-chairs the Sport Injury Prevention research Centre at the University of Calgary funded by the International Olympic Committee (IOC).
Jon Patricios MD	Sports Concussion South Africa, Johannesburg, South Africa The Section of Sports Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa	Dr Jon Patricios received funding from the South African Rugby Union through its BokSmart injury prevention programme to attend the Concussion in Sport Consensus meeting.
Christine Providenza PhD	Parachute (Formerly Safe Communities Canada, Safe Kids Canada, SMARTRISK, and ThinkFirst Canada), Toronto, Ontario, Canada	No disclosures
Andrew Gardner PsyD	Centre for Translational Neuroscience and Mental Health, School of Medicine & Public Health, University of Newcastle, NSW, Australia.	No disclosures
Richard G. Ellenbogen, MD, FACS	Professor and Chairman Theodore S. Roberts Endowed Chair Department of Neurological Surgery University of Washington Seattle, WA, USA Co-Chair, NFL Head, Neck and Spine Medical Committee	None (National Football League, Volunteer Consultant to Commissioner . No financial remunerations received.)