



Editor's choice
Scan to access more
free content

Current physical activity guidelines for health are insufficient to mitigate long-term weight gain: more data in the fitness versus fatness debate (The HUNT study, Norway)

Trine Moholdt,^{1,2} Ulrik Wisløff,¹ Stian Lydersen,³ Javaid Nauman¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2014-093416>)

¹Department of Circulation and Medical Imaging, Faculty of Medicine, KG Jebsen Center of Exercise in Medicine, Norwegian University of Science and Technology, Trondheim, Norway
²Department of Obstetrics and Gynecology, St. Olav's Hospital, Trondheim, Norway
³Faculty of Medicine, Regional Centre for Child and Youth Mental Health and Child Welfare, Norwegian University of Science and Technology, Trondheim, Norway

Correspondence to

Dr Trine Moholdt, Department of Circulation and Medical Imaging, Faculty of Medicine, KG Jebsen Center of Exercise in Medicine, Prinsesse Kristinas gt 3/ PO Box 8905, 7491 Trondheim 7022, Norway; trine.moholdt@ntnu.no

Accepted 3 April 2014
Published Online First
29 April 2014

ABSTRACT

Background To promote and maintain health, all adults are recommended to do moderate-intensity aerobic activity a minimum of 30 min on 5 days, or vigorous-intensity activity of 20 min on 3 days, each week. Whether these levels prevent long-term weight gain is uncertain.

Objective To assess the relationship between physical activity and long-term weight gain.

Study design An observational prospective cohort study.

Methods Weight and physical activity were measured in the Nord-Trøndelag Health Study in 1984–1986, 1995–1997 and 2006–2008. Participants (n=19 127) were classified based on physical activity into inactive, below recommended level, recommended level or above recommended level. We carried out adjusted mixed model regression analyses with weight as outcome.

Results Men maintaining physical activity above the recommendations for 33 years increased 5.6 kg, while inactive men increased 9.1 kg. For women, corresponding numbers were 3.8 kg in those above recommended physical activity levels, and 9.5 kg in inactive. In adjusted analyses, physical activity above the recommendations was associated with 2.1 kg (95% CI 1.8 to 2.4) less weight gain in men over any 11-year period, compared with inactive. Women exceeding the recommendations gained 1.8 kg (CI 1.5 to 2.2) less than inactive. Compared with inactive, the ORs of gaining meaningful weight of ≥ 2.3 kg were 0.79 (CI 0.69 to 0.91) and 0.70 (CI 0.60 to 0.81) if exceeding the recommendations in men and women, respectively.

Conclusions Physical activity above the current recommendations for health benefits was associated with significantly lower risk of weight gain.

INTRODUCTION

The western world is facing an obesity epidemic, and the prevalence of obesity among US adults has now reached 35%.¹ During the past 15 years, obesity prevalence has increased in Europe, and predictions suggest a sizeable further increase in the European population by 2015.² Being overweight or obese is associated with increased overall mortality and cardiovascular disease incidence, as well as with reduced physical and social functioning and quality of life.^{3–5}

The significant public health burden of increased prevalence of overweight and obesity strongly emphasises the need for effective strategies for weight gain prevention. As energy expended during

physical activity potentially affects body weight, we need concise guidelines about the required amount of physical activity to prevent weight gain. The current recommendations on physical activity to promote and maintain health from the American College of Sports Medicine state that all healthy adults aged 18–65 should do moderate-intensity aerobic activity for a minimum of 30 min on 5 days each week, or vigorous activity for a minimum of 20 min on 3 days each week.⁶ The same amount of physical activity is recommended for older adults.⁷ Regarding cardiovascular mortality risk, there are some studies indicating that even smaller amounts of physical activity is adequate to significantly lower the risk.^{8,9} Although not being the main goal of exercise,¹⁰ weight gain prevention is acknowledged as one of the beneficial effects. It should be evaluated, however, if the current recommendations are sufficient to combat the ongoing obesity epidemic.^{11–13}

Lee *et al*¹¹ recently found that complying with the recommended activity levels (30 min of moderate activity on most of the days) was insufficient for weight gain prevention over 13 years in middle-aged women. Only when physically active for 60 min or more per day, women were less likely to gain weight. Importantly, this association was only found in women of normal weight, and no significant association between weight gain and physical activity was observed in women who were already overweight or obese. In contrast, Hankinson *et al*¹² found that 30 min of daily physical activity was enough to prevent weight gain in young adults.

Our aim was, therefore, to examine the associations of physical activity with weight gain in men and women throughout the adult lifespan, and to investigate potential interactions with age and body mass index (BMI). Furthermore, we argue that leisure time and occupational physical activity should be considered to further explore the association between weight gain and physical activity. Our hypotheses were that higher levels of leisure time and occupational physical activity were associated with lower weight gain, and that complying with the current recommendations for physical activity would not be sufficient to prevent weight gain.

METHODS

Study population

We used data from a Norwegian population study, the Nord-Trøndelag Health Study (HUNT), parts 1–3. Data collection in the HUNT study was



CrossMark

To cite: Moholdt T, Wisløff U, Lydersen S, *et al*. *Br J Sports Med* 2014;**48**:1489–1496.

carried out in the Nord-Trøndelag county in the middle part of Norway in three waves, first in 1984–1986 (HUNT 1, H1), second in 1995–1997 (HUNT 2, H2) and the third in 2006–2008 (HUNT 3, H3). Data collection in the HUNT studies is described in detail elsewhere.¹⁴ Briefly, all inhabitants ≥ 20 years in the county were invited to participate, with approximate overall participant rates in H1, H2 and H3 of 88%, 70% and 56%, respectively. The participants came for clinical visits where anthropometry, resting heart rate, blood pressure, blood lipids and glucose were measured. They also filled out questionnaires about lifestyle variables (such as physical activity, smoking, alcohol consumption) and background variables (such as education, occupation, marital status). For this study, we selected the participants with data on age, weight and physical activity at baseline in H1, and who participated in all three HUNT waves. Participants with cardiovascular disease during the follow-up period ($n=4\ 347$) were excluded, leaving 8433 men and 10 783 women (figure 1).

Assessment of leisure time physical activity

At baseline in H1, the participants were asked about their exercise frequency (never, less than once per week, once per week, two to three times per week or four or more times per week), duration (less than 15, 15–30, 31–60 or more than 60 min) and intensity (no sweat, sweat or exhausted). The questions on duration and intensity have been validated against direct measurements of maximal oxygen uptake.¹⁵ The ‘no sweat’ category corresponded to an intensity of about 55% of maximal oxygen uptake, regardless of the exercise duration up to 90 min, whereas the ‘sweat’ and ‘exhausted’ categories corresponded to intensities of 71–75% and 76–90% of maximal oxygen uptake, respectively, with decreased relative intensity with increasing duration of exercise.¹⁵ In H2, the participants were asked about their weekly amount of light physical activity and hard physical activity, with alternatives of ‘none’, ‘less than an hour’, ‘1–2 h’ and ‘3 h or more’. The validity of the H2 questions was found to be poorer than for the H1 questions, but the hard physical activity category was reasonably well correlated with VO_{2peak} and with variables in the widely accepted International Physical Activity Questionnaire (IPAQ).¹⁶ In H3, the questions on physical activity were identical as in H1.

We constructed a four-level leisure time physical activity index to match with the current recommendations to promote and maintain health⁶; moderate-intensity aerobic activity a minimum of 30 min on 5 days, or vigorous-intensity activity of 20 min on 3 days, each week. Moderate-intensity and vigorous self-reported activity were taken into account when categorising participants, and physical activity was updated over time to allow for individual changes in activity level. The ‘inactive’ level was the one reporting no leisure time physical activity. Participants who reported physical activity below the recommendations were classified as ‘below recommended’. The ‘recommended’ level was participants reporting physical activity corresponding to the current recommendations, either as moderate-intensity exercise for 150 min each week, or as 60 min of vigorous intensity, or a combination of these, whereas participants who reported to be more active than the recommendations were classified as ‘above recommended’.

Other measurements

Occupational physical activity was assessed in all three HUNT waves and classified as ‘sedentary’, ‘walk and lift’ and ‘heavy manual work’. In the analyses, we also used data on age, gender, height, weight, BMI, sitting time, smoking status and alcohol intake. Height and weight were measured with direct measurements. BMI is weight in kilograms divided by height in metres squared. Sitting time was only measured in H3, with the question: “On a regular day, how many hours do you spend sitting?” If the participants did not answer this question, we used the sum in hours of these three questions: “How many hours per day do you usually spend in front of a computer at work?”, “How many hours per day do you usually spend in front of a computer in leisure time?” and “How many hours per day do you usually watch TV/DVD/video?”

Statistical analysis

Physical activity level differs between men and women¹⁷; therefore, we present sex-specific analyses. We compared the baseline characteristics for the participants according to the four levels of physical activity, using linear regression, linear-by-linear test for association, Kruskal-Wallis test and the Cochran-Armitage test for trend for continuous, ordinal, nominal and dichotomous




Figure 1 Flow chart of participants in the study (CVD; cardiovascular disease).

variables, respectively. Linear mixed models with random intercept and slope were used to examine changes in body weight over time according to physical activity levels, using inactive participants as reference. Physical activity level was updated over time. Three models were used; model 1 adjusted for age (continuous), model 2 additionally adjusted for diabetes status (yes or no), smoking (never, former or current) and alcohol (abstainers, <1 time/14 days, 1–4 times/14 days or ≥ 5 times/14 days) and time period indicators of HUNT wave (H1, H2 or H3), and model 3 also adjusted for occupational physical activity (sedentary, walk and lift or heavy manual work).

We also performed stratified analyses with interaction terms to assess whether the prospective changes in body weight by physical activity levels could be modified by other factors. We investigated the potential effect modification by categorised age (<40, 40–59 or ≥ 60 years), BMI (<18.5, 18.5–24.9, 25.0–29.9 or ≥ 30 kg/m²), smoking status (current, former or never smokers) and occupational physical activity (sedentary, walk and lift or heavy manual work). Age categories were selected to give information about young adults, middle-aged and older participants.

We then assessed the OR for meaningful weight gain over time, defined as gaining ≥ 2.3 kg¹⁸ associated with the four physical activity categories using mixed model logistic regression analyses. Stratified analyses were conducted to assess the potential interactions with age, BMI, smoking or occupational physical activity. In separate analyses, we assessed weight change in a subgroup of 2673 participants who reported no change in physical activity over time.

We also performed sensitivity analyses to assess the robustness of our findings. For example, the association between body weight and physical activity could be influenced by time spent in a sedentary state. We examined changes in body weight associated with sedentary time only in H3 because the former two waves did not have information about sedentary behaviour. In another sensitivity analysis, we adjusted for baseline body weight measured at H1. Ninety-five per cent CIs are reported where relevant, and two-sided, *p* values of <0.05 were considered significant. The statistical analyses were conducted using Stata V.12.1 (StataCorp).

RESULTS

Baseline characteristics of men and women by level of physical activity are presented in table 1. Age, weight, BMI, smoking, alcohol consumption, hypertension and occupational activity varied across physical activity groups at baseline, where active participants had favourable cardiovascular profile compared with inactive participants. The mean overall increase in weight during 22 years of follow-up was 7.7 kg (CI 7.4 to 8.1) for men and 8.5 kg (CI 8.2 to 8.8) for women. The weight gain was largest between the first and second wave, with 5.3 kg (CI 5.0 to 5.6) increase for men and 6.1 kg (CI 5.8 to 6.4) increase for women.

In multadjusted analyses for age, smoking, alcohol and diabetes status, we found a clear dose–response association (*p*<0.001 for trend) between changes in body weight according to physical activity levels after allowing physical activity to be updated over an 11-year period in men and women. Compared with inactive men, above recommended level of physical activity was associated with 2.1 kg (CI 1.8 to 2.4) less weight gain, whereas recommended level of physical activity was associated with 0.7 kg (CI 0.5 to 0.9) less weight gain (*p* for trend <0.001). The corresponding estimates of lower weight gain among women were 1.8 kg (CI 1.5 to 2.2) and 0.5 kg (CI 0.3 to

0.7) for above recommended and recommended levels, respectively (*p* for trend <0.001).

We assessed modifying effects of age, BMI, smoking status and occupational activity on the association between physical activity and weight change, and found significant interactions with age and BMI in men and women. However, occupational activity and smoking modified the association only in men, and we found no evidence of interaction among women (table 2). In stratified analyses of BMI, there was a clear dose–response relationship between physical activity and weight gain only for overweight/obese men (BMI >25 kg/m²) and obese women (BMI >25 kg/m²), with a *p* for trend <0.001 for both. We observed less weight gain associated with above recommended levels, that is, >150 min of moderate-intensity activity or >60 min of vigorous activity each week, in overweight/obese (BMI >25 kg/m²; *p*<0.001) men and obese (BMI >30 kg/m²) women (table 2). Obese men and women also had significantly lower weight gain with recommended level of physical activity, that is, 150 min moderate-intensity or 60 min of vigorous activity weekly, compared with the reference category of inactive participants (table 2).

We also examined ORs of gaining ≥ 2.3 kg (meaningful weight gain) associated with physical activity updated over time. Compared with inactive participants, the ORs of meaningful weight gain in the above recommended category were 0.79 (CI 0.69 to 0.91) and 0.70 (CI 0.60 to 0.81) in men and women, respectively (table 3). There was a significant interaction with age for the odds of meaningful weight gain across levels of physical activity for both sexes, with lower ORs of gaining meaningful weight when physically active in older participants. In men >40 years, there was significantly reduced OR of meaningful weight gain with above recommended levels of physical activity. This was also true for women 40–59 years, whereas women >60 years had reduced OR of meaningful weight gain with recommended and above recommended levels of physical activity. BMI modified the association in women (*p* values for interaction <0.001), and only normal weight (BMI 18.5–24.9 kg/m²) and overweight (BMI 25.0–29.9 kg/m²) women were less likely to gain meaningful weight associated with high levels of physical activity (table 3). In men, all but the lowest BMI category (<18.5 kg/m²) had lower ORs for meaningful weight gain associated with above recommended levels of physical activity.

In subgroup analyses of men and women who reported to maintain their physical activity over the years, we observed that men who maintained above recommended levels of physical activity had increased 5.6 kg (CI 3.4 to 7.9) in body weight, while those who remained inactive had increased 9.1 kg (CI 7.9 to 10.3). Recommended levels of physical activity were associated with 7.2 kg (CI 5.5 to 8.8) increase in men. For women, maintaining above recommended levels of physical activity was associated with an increase of 3.8 kg (CI –0.2 to 7.8), compared with inactive women who increased 9.4 kg (CI 8.1 to 10.8). Recommended levels of physical activity in women were associated with an increase of 8.6 kg (CI 6.9 to 10.3; figure 2).

In separate sensitivity analyses, the results were not substantially different from the main analyses. When we adjusted for body weight at baseline, we found that the results were similar to the results obtained without this adjustment (see online supplementary table S1). In a cross sectional analysis of body weight and physical activity from H3, we also adjusted for sitting time in addition to the variables that were already included in the multivariable models. We observed similar dose–response inverse associations between body weight and levels of physical activity after additional adjustments of sitting time (see online supplementary table S2). Among men, the body weight

Table 1 Characteristics of study participants according to baseline (HUNT-1) physical activity levels (N=19 127)

	Men (n=8344)				p Value*	Women (n=10 783)				p Value*
	Inactive (n=3652)	Below recommended (n=2457)	Recommended (n=1578)	Above recommended (n=657)		Inactive (n=4500)	Below recommended (n=4245)	Recommended (n=1723)	Above recommended (n=315)	
Age, mean (range), years	37.9 (20–71)	39.3 (20–79)	38.7 (21–72)	34.9 (20–71)	<0.001	38.5 (20–73)	39.1 (20–72)	39.4 (20–68)	38.7 (20–69)	0.01
Weight, kg (SD)	78.7 (10.3)	79.2 (9.6)	78.2 (9.0)	77.3 (8.4)	<0.001	64.9 (10.6)	64.4 (9.6)	64.6 (9.5)	64.6 (10.6)	0.01
Height, cm (SD)	178.2 (6.3)	178.4 (6.4)	178.5 (6.1)	178.9 (6.1)	0.01	164.8 (5.6)	165.0 (5.6)	165.0 (5.4)	165.3 (6.2)	0.30
Body mass index, kg/m ² , N (%)										
<18.5	21 (0.6)	5 (0.2)	3 (0.2)	3 (0.5)	<0.001	141 (3.1)	76 (1.8)	24 (1.4)	4 (1.3)	<0.001
18.5–24.9	1982 (54.3)	1377 (56.0)	981 (62.2)	439 (66.8)		2952 (65.6)	3001 (70.7)	1187 (68.9)	225 (71.4)	
25.0–29.9	1443 (39.5)	969 (39.4)	554 (35.1)	207 (31.5)		1075 (23.9)	940 (22.1)	408 (23.7)	70 (22.2)	
≥30.0	206 (5.6)	106 (4.3)	40 (2.5)	8 (1.2)		332 (7.4)	228 (5.4)	104 (6.0)	16 (5.1)	
Smoking status, N (%)†										
Never	1127 (30.9)	957 (38.9)	788 (49.9)	353 (53.7)	<0.001	1619 (36.0)	1868 (44.0)	788 (45.7)	148 (47.0)	<0.001
Former	961 (26.3)	741 (30.2)	440 (27.9)	151 (23.0)		870 (19.3)	889 (20.9)	396 (23.0)	80 (25.4)	
Current	1317 (36.1)	622 (25.3)	244 (15.5)	106 (16.1)		1669 (37.1)	1206 (28.4)	419 (24.3)	63 (20.0)	
Occupational physical activity, N (%)†										
Sedentary	446 (12.2)	392 (16.0)	320 (20.3)	132 (20.1)	<0.001	435 (9.7)	475 (11.2)	258 (14.9)	43 (13.7)	<0.001
Walk and lift	2660 (72.8)	1799 (73.2)	1095 (69.4)	422 (64.2)		3439 (76.4)	3272 (77.1)	1243 (72.1)	222 (70.5)	
Heavy	412 (11.3)	161 (6.6)	80 (5.1)	66 (10.0)		410 (9.1)	279 (6.6)	109 (6.3)	27 (8.6)	
Alcoholic drinks, N (%)†‡										
Abstainer	169 (4.6)	89 (3.6)	50 (3.2)	33 (5.0)	0.85	421 (9.4)	293 (6.9)	117 (6.8)	27 (8.6)	<0.001
<1	1061 (29.1)	685 (27.9)	459 (29.1)	194 (29.5)		2317 (51.5)	2088 (49.2)	787 (45.7)	153 (48.6)	
1–4	2106 (57.7)	1490 (60.6)	954 (60.5)	387 (58.9)		1569 (34.9)	1704 (40.1)	746 (43.3)	113 (35.9)	
≥5	298 (8.2)	188 (7.7)	111 (7.0)	39 (5.9)		149 (3.3)	111 (2.6)	58 (3.4)	14 (4.4)	
Hypertension status, N (%)†										
	1330 (36.4)	940 (38.3)	511 (32.4)	206 (31.4)	0.001	936 (20.8)	877 (20.7)	381 (22.1)	69 (21.9)	0.33
Diabetes status, N (%)†										
	19 (0.5)	12 (0.5)	7 (0.4)	2 (0.3)	0.48	15 (0.3)	14 (0.3)	8 (0.5)	2 (0.6)	0.34

*For linear trend, regression analyses were used for continuous variables; χ^2 tests for proportions of categorical variables.

†Data are missing for this category.

‡Based on consumption over a 2-week period.

HUNT, Nord-Trøndelag County Health Study.

Table 2 Difference in body weight in kilograms, mean (95% CI) over any 11-year period by physical activity levels

	N*	Physical activity categories				p Value†	p Value‡
		Inactive	Below recommended	Recommended	Above recommended		
Men							
Model 1	8344	Ref.	0.02 (−0.18 to 0.22)	−0.16 (−0.38 to 0.06)	−1.46 (−0.38 to −1.16)	<0.001	
Model 2	8334	Ref.	−0.12 (−0.31 to 0.07)	−0.70 (−0.91 to −0.48)	−2.10 (−2.39 to −1.81)	<0.001	
Model 3	8230	Ref.	−0.21 (−0.41 to −0.01)	−0.75 (−0.97 to −0.52)	−1.80 (−2.10 to −1.49)	<0.001	
Stratified analyses§							
Age, years							
<40	5042	Ref.	0.25 (−0.22 to 0.72)	−0.33 (−0.84 to 0.18)	−1.08 (−1.72 to −0.43)	0.003	<0.001
40–59	3096	Ref.	−0.44 (−0.73 to −0.16)	−1.06 (−1.37 to −0.76)	−1.89 (−2.32 to −1.46)	<0.001	
≥60	206	Ref.	−0.32 (−0.87 to 0.23)	−0.66 (−1.24 to −0.80)	−1.68 (−2.43 to −0.94)	<0.001	
Occupational activity							
Sedentary	1290	Ref.	−0.20 (−0.67 to 0.26)	−0.95 (−1.44 to −0.46)	−2.80 (−3.47 to −2.13)	<0.001	0.001
Walk and life	5976	Ref.	−0.17 (−0.47 to 0.13)	−0.73 (−1.07 to −0.40)	−1.78 (−2.26 to −1.30)	<0.001	
Heavy	719	Ref.	−0.40 (−0.99 to 0.18)	−0.88 (−1.50 to −0.26)	−1.07 (−1.79 to −0.34)	0.001	
Smoking status							
Never	3225	Ref.	−0.31 (−0.61 to −0.004)	−1.08 (−1.41 to −0.76)	−2.45 (−2.88 to −2.02)	<0.001	0.005
Former	2293	Ref.	−0.12 (−0.48 to 0.23)	−0.62 (−1.01 to −0.23)	−2.18 (−2.69 to −1.66)	<0.001	
Current	2289	Ref.	0.08 (−0.35 to 0.50)	−0.31 (−0.82 to 0.19)	−1.62 (−2.34 to −0.89)	0.001	
BMI categories, kg/m ²							
<18.5	32	Ref.	−2.37 (−5.34 to 0.60)	0.47 (−2.82 to 3.75)	−0.35 (−4.28 to 3.58)	0.64	<0.001
18.5–24.9	4779	Ref.	0.68 (0.43 to 0.93)	0.51 (0.24 to 0.77)	−0.19 (−0.53 to 0.16)	0.60	
25.0–29.9	3173	Ref.	0.07 (−0.15 to 0.29)	−0.13 (−0.37 to 0.10)	−0.97 (−1.29 to −0.65)	<0.001	
≥30.0	360	Ref.	−1.24 (−1.90 to −0.58)	−1.41 (−2.14 to −0.68)	−2.53 (−3.60 to −1.46)	<0.001	
Women							
Model 1	10 783	Ref.	−0.23 (−0.42 to −0.04)	−0.18 (−0.40 to 0.03)	−1.41 (−1.76 to −1.06)	<0.001	
Model 2	10 744	Ref.	−0.07 (−0.26 to 0.11)	−0.49 (−0.70 to −0.27)	−1.84 (−2.19 to −1.50)	<0.001	
Model 3	10 413	Ref.	−0.16 (−0.36 to 0.03)	−0.69 (−0.91 to −0.45)	−2.10 (−2.48, −1.72)	<0.001	
Stratified analyses§							
Age, years							
<40	6245	Ref.	−0.21 (−0.62 to 0.19)	−0.42 (−0.92 to 0.08)	−0.78 (−1.70 to 0.13)	0.04	<0.001
40–59	4163	Ref.	−0.31 (−0.59 to −0.03)	−1.16 (−1.15 to −0.85)	−2.24 (−2.86 to −1.83)	<0.001	
≥60	375	Ref.	−0.44 (−0.94 to 0.07)	−1.09 (−1.68 to −0.50)	−3.31 (−4.22 to −2.41)	<0.001	
Occupational activity							
Sedentary	1211	Ref.	−0.25 (−0.76 to 0.24)	−1.04 (−1.57 to −0.50)	−2.75 (−3.62 to −1.88)	<0.001	0.90
Walk and life	8176	Ref.	−0.22 (−0.46 to 0.02)	−0.76 (−1.03 to −0.48)	−2.12 (−2.59 to −1.66)	<0.001	
Heavy	825	Ref.	−0.52 (−1.82 to 0.78)	−0.51 (−2.12 to 1.10)	−2.76 (−5.11 to −0.41)	0.05	
Smoking status							
Never	4423	Ref.	−0.16 (−0.44 to 0.12)	−0.45 (−0.77 to −0.12)	−1.77 (−2.29 to −1.25)	<0.001	0.09
Former	2235	Ref.	−0.17 (−0.57 to 0.23)	−0.79 (−1.24 to −0.34)	−2.44 (−3.12 to −1.76)	<0.001	
Current	3357	Ref.	−0.07 (−0.42 to 0.29)	−0.43 (−0.86 to −0.01)	−1.36 (−2.15 to −0.58)	0.002	
BMI categories, kg/m ²							
<18.5	245	Ref.	−0.04 (−0.82 to 0.75)	0.59 (−0.49 to 1.67)	2.34 (0.22 to 4.46)	0.09	<0.001
18.5–24.9	7365	Ref.	0.19 (0.02 to 0.37)	0.42 (0.22 to 0.62)	−0.16 (−0.47 to 0.15)	0.05	
25.0–29.9	2493	Ref.	−0.13 (−0.37 to 0.09)	−0.19 (−0.45 to 0.06)	−0.34 (−0.76 to 0.08)	0.07	
≥30.0	680	Ref.	−0.99 (−1.58 to −0.39)	−1.78 (−2.45 to −1.12)	−2.32 (−3.52 to −1.11)	<0.001	

Model 1, adjusted for age; model 2 also for diabetes status (yes or no), smoking (never, former or current), and alcohol (abstainers, <1 time/14 days, 1–4 times/14 days or ≥5 times/14 days) and time period-dummies of HUNT wave (H1, H2 or H3) and model 3 additionally for occupational physical activity (sedentary, walk and life or heavy manual work).

*Number of participants at baseline (HUNT-1).

†p Value for trend across categories.

‡p Value for interaction.

§Analytical model 2 is used in all stratified analyses.

BMI, body mass index; HUNT, Nord-Trøndelag County Health Study.

was 4.4 kg (CI 3.4 to 5.3) lower in those who reported above recommended levels of physical activity, compared with inactive men. Similarly, among women who reported above recommended levels of physical activity, body weight was 6.7 kg (CI 5.7 to 7.7) lower compared with women who were inactive (see online supplementary table S2).

DISCUSSION

In this prospective study of physical activity and body weight over three decades, we found a strong dose-dependent inverse association between activity levels and weight gain in men and women. There was an overall weight gain over time, and participants reporting of physical activity corresponding to or

Table 3 ORs (95% CI) for ≥ 2.3 kg increase in body weight over any 11-year period by physical activity levels

	N*	Physical activity categories				p Value†	p Value‡
		Inactive	Below recommended	Recommended	Above recommended		
Men							
Model 1	8344	Ref.	1.09 (0.99 to 1.19)	1.06 (0.97 to 1.17)	0.81 (0.71 to 0.92)	0.07	
Model 2	8334	Ref.	1.02 (0.92 to 1.13)	0.98 (0.88 to 1.09)	0.79 (0.69 to 0.91)	0.01	
Model 3	8230	Ref.	1.00 (0.90 to 1.11)	0.97 (0.87 to 1.08)	0.79 (0.69 to 0.91)	<0.01	
Stratified analyses§							
Age, years							
<40	5042	Ref.	1.04 (0.89 to 1.20)	0.92 (0.79 to 1.08)	0.89 (0.72 to 1.09)	0.18	0.002
40–59	3096	Ref.	0.99 (0.87 to 1.13)	0.97 (0.85 to 1.11)	0.77 (0.64 to 0.92)	0.03	
≥ 60	206	Ref.	0.98 (0.72 to 1.33)	0.96 (0.71 to 1.31)	0.41 (0.28 to 0.62)	<0.01	
Occupational activity							
Inactive	1290	Ref.	0.87 (0.69 to 1.10)	0.87 (0.69 to 1.10)	0.64 (0.47 to 0.87)	0.01	0.83
Walk and life	5976	Ref.	0.98 (0.86 to 1.12)	0.92 (0.80 to 1.06)	0.72 (0.59 to 0.89)	<0.01	
Heavy	719	Ref.	1.17 (0.86 to 1.59)	0.91 (0.66 to 1.26)	0.77 (0.53 to 1.11)	0.16	
Smoking status							
Never	3 225	Ref.	0.98 (0.83 to 1.15)	0.88 (0.75 to 1.04)	0.77 (0.63 to 0.95)	0.01	0.74
Former	2 293	Ref.	1.01 (0.85 to 1.20)	0.99 (0.83 to 1.19)	0.72 (0.57 to 0.92)	0.05	
Current	2 289	Ref.	1.05 (0.87 to 1.27)	1.02 (0.82 to 1.26)	0.75 (0.55 to 1.02)	0.32	
BMI categories, kg/m²							
<18.5	32	Ref.	0.66 (0.11 to 4.04)	1.31 (0.19 to 9.26)	0.97 (0.05 to 18.8)	0.93	0.27
18.5–24.9	4779	Ref.	0.92 (0.79 to 1.06)	0.96 (0.82 to 1.12)	0.78 (0.64 to 0.95)	0.05	
25.0–29.9	3173	Ref.	1.07 (0.93 to 1.23)	1.01 (0.87 to 1.16)	0.81 (0.67 to 0.98)	0.12	
≥ 30.0	360	Ref.	1.13 (0.83 to 1.53)	0.96 (0.69 to 1.33)	0.54 (0.34 to 0.86)	0.06	
Women							
Model 1	10 783	Ref.	1.34 (1.24 to 1.44)	1.54 (1.42 to 1.67)	1.18 (1.04 to 1.36)	<0.001	
Model 2	10 744	Ref.	1.04 (0.96 to 1.13)	1.03 (0.93 to 1.13)	0.70 (0.60 to 0.81)	0.02	
Model 3	10 413	Ref.	1.01 (0.92 to 1.10)	0.97 (0.88 to 1.07)	0.69 (0.59 to 0.82)	0.006	
Stratified analyses§							
Age, years							
<40	6245	Ref.	0.97 (0.86 to 1.10)	0.86 (0.74 to 0.99)	0.81 (0.62 to 1.07)	0.02	0.03
40–59	4163	Ref.	1.03 (0.92 to 1.16)	1.02 (0.90 to 1.15)	0.60 (0.49 to 0.73)	<0.01	
≥ 60	375	Ref.	0.98 (0.78 to 1.23)	0.73 (0.57 to 0.94)	0.31 (0.21 to 0.44)	<0.001	
Occupational activity							
Inactive	1211	Ref.	0.93 (0.73 to 1.18)	0.89 (0.69 to 1.14)	0.48 (0.32 to 0.72)	0.01	0.76
Walk and life	8176	Ref.	1.02 (0.92 to 1.14)	0.97 (0.86 to 1.09)	0.70 (0.57 to 0.85)	0.02	
Heavy	825	Ref.	0.72 (0.45 to 1.14)	0.84 (0.47 to 1.49)	0.68 (0.30 to 1.54)	0.27	
Smoking status							
Never	4423	Ref.	1.04 (0.91 to 1.18)	1.01 (0.87 to 1.16)	0.68 (0.54 to 0.85)	0.05	0.68
Former	2235	Ref.	0.99 (0.83 to 1.18)	0.84 (0.70 to 1.01)	0.55 (0.41 to 0.72)	<0.001	
Current	3357	Ref.	1.02 (0.88 to 1.18)	1.11 (0.94 to 1.32)	0.60 (0.44 to 0.82)	0.42	
BMI categories, kg/m²							
<18.5	245	Ref.	0.49 (0.22 to 1.06)	1.24 (0.41 to 3.73)	0.83 (0.11 to 6.49)	0.69	<0.001
18.5–24.9	7365	Ref.	0.91 (0.81 to 1.01)	0.91 (0.81 to 1.03)	0.73 (0.60 to 0.88)	0.005	
25.0–29.9	2493	Ref.	1.14 (1.00 to 1.31)	1.17 (1.00 to 1.36)	0.77 (0.60 to 0.98)	0.86	
≥ 30.0	680	Ref.	1.09 (0.87 to 1.37)	1.17 (0.90 to 1.52)	0.64 (0.40 to 1.03)	0.97	

Abbreviations: BMI, body mass index, calculated as body weight in kilograms divided by height in meters squared.

Model 1, adjusted for age; model 2 also for diabetes status (yes or no), smoking (never, former or current), and alcohol (abstainers, <1 time/14 days, 1–4 times/14 days or ≥ 5 times/14 days) and time period-dummies of HUNT wave (H1, H2 or H3) and model 3 additionally for occupational physical activity (sedentary, walk and life or heavy manual work).

*Number of participants at baseline (HUNT-1).

†p Value for trend across categories.

‡p Value for interaction.

§Analytical model 2 is used in all stratified analyses.

BMI, body mass index; HUNT, Nord-Trøndelag County Health Study.

exceeding recommended levels of physical activity had significantly less pronounced weight gain, compared with the inactive reference category. Furthermore, only physical activity exceeding current recommendations of 150 min of moderate-intensity or 60 min of vigorous activity each week was required to prevent significant weight gain, defined as ≥ 2.3 kg over any 11-year

period. Besides, maintaining such high levels (above recommendations) of physical activity for 22 years was associated with 3.5 and 5.6 fewer kg of weight gain compared with staying inactive during this time period in men and women, respectively.

Although extensive research has been carried out to assess the amount of physical activity required for weight loss among




Figure 2 Weight at Nord-Trøndelag County Health Study (HUNT) 1, HUNT 2 and HUNT 3 in men and women who maintained the same level of physical activity during the whole follow-up period.

already overweight or obese participants, fewer studies have quantified how much physical activity is needed for weight gain prevention.¹⁷ Our results are in line with some previous studies^{11–13 19–23} to show an overall favourable effect of physical activity for weight gain prevention. Lee *et al*¹¹ followed middle-aged women for in average 13 years, and found physical activity to be associated with less weight gain only among those with BMI lower than 25 and among women younger than 65 years. Furthermore, they compared two lower levels of physical activity, one equivalent to <150 min/week and one 150–420 min of moderate-intensity physical activity, to a reference category of ≥ 420 min of moderate-intensity activity. They found only the highest category, that is, ≥ 420 min of moderate-intensity activity, to prevent weight gain.

Our results contrast with the findings of Lee *et al*¹¹ as we found that moderate amounts of physical activity, equivalent to current recommendations of 150 min of moderate-intensity activity, might be sufficient to reduce overall weight gain in men and women. We found, however, greater effects of physical activity above the recommendations.

Interestingly, we found that only physical activity above the recommended level was required to lessen the OR for meaningful weight gain (ie, ≥ 2.3 kg) over time, in line with findings of Lee *et al*.¹¹ In our study, this association was present also for overweight women (BMI 25–29.9 kg/m²) and overweight and obese men (BMI ≥ 25 kg/m²), as opposed to Lee *et al*¹¹ who found significantly lower weight gain with high levels of physical activity in normal weight women only.

In regard to different age groups, our results are partly different than the results of Lee *et al*¹¹ as we found significantly lower weight gain with higher levels of physical activity also in participants ≥ 65 years. Shiroma *et al*¹³ studied the association of physical activity and weight gain in men at the mean age of 65 years at baseline, and found that being inactive was associated with meaningful weight gain over time compared with being moderately active for ~ 70 min/day. In line with this, we found the OR for meaningful weight gain over 11 years to be significantly reduced in men > 60 years who exceeded 150 min of moderate activity or 60 min of vigorous activity. However, women > 60 years also had a significantly reduced meaningful weight gain when doing 150 min of moderate-intensity activity/60 min of vigorous activity.

To summarise, our findings in older participants do not fully support other data^{11 13} as we found physical activity in older men and women to be associated in similar inverse dose-response relationship with weight gain as in young participants.

Some of the differences between the mentioned studies might be caused by the different physical activity questionnaires used. In our study, we also asked about high-intensity activity, allowing categorisation of physical activity according to the recommendations proposed by Haskell *et al*⁶ stating that 150 min of moderate-intensity activity, or 60 min of vigorous activity each week, are needed for health benefits.

In a subgroup analysis of 2673 participants, we found that maintenance of physical activity above the recommended level over 22 years was associated with significantly smaller gains in weight among men and women, whereas complying with recommendations was not.

Using data from the Coronary Artery Risk Development in Young Adults (CARDIA) study, Hankinson *et al*¹² found less weight gain in young adults who maintained an activity level corresponding to 150 min of moderate-intensity activity each week throughout young adulthood. However, our data extend these findings as we included different age groups, BMI categories, smoking status and occupational activity strata. Of note, in our study, the most pronounced difference in weight gain was seen when comparing the participants who maintained an activity level above the current recommendations and the other three groups.

Our findings suggest that women more than men have greater control of weight gain by physical activity, and this is consistent with previous results in showing sex differences for the associations between physical activity and weight gain.^{20 22 23} One plausible explanation for the observed sex differences may be more over-reporting of physical activity among men compared with women. In line with this, the relationship between body weight and physical activity is often found to be stronger for women when physical activity is assessed by self-report,^{24 25} whereas the relationship is similar or stronger in men when an objective measure like doubly labelled water is used.²⁶

The adjusted and stratified analyses of daily sitting time and occupational activity suggest that physical activity at leisure time is independently associated with changes in body weight over the years. We observed similar dose-dependent association of weight gain prevention relating to above recommended levels of physical activity after additional adjustments of sitting time and occupational activity. As occupational activity, and especially the effects of sedentary time at work, would potentially affect total energy expenditure greatly, more studies are warranted to further investigate these interactions.²⁷

Although our results showed less pronounced weight gain when exceeding current recommendations, participants in all activity

levels gained weight over 22 years. Based on our data, some weight gain throughout adult life seems to be inevitable, but a high level of physical activity reduces the magnitude of weight gain in men and women. Besides, even if physical activity above the current recommendations for physical activity might be needed to prevent unhealthy weight gain, there is a vast body of literature showing that the recommended 150 min of weekly moderate activity lower the risk of chronic diseases and mortality.^{28–30}

The main strengths of our study include a large sample size of men and women with a wide age range of 20–100 years, who were recruited from three total health surveys in one county with a high participation rate. This allowed for the prospective design with a more than 20 years of follow-up period, and with the ability to adjust for changes in physical activity levels over time. Furthermore, directly measuring body weight in a standardised way is regarded another strength as some participants tend to underestimate their weight when self-reported.³¹ We were able to adjust for many variables potentially confounding the relation between physical activity and weight gain, and also included occupational activity in the analyses.

Limitations include self-reported physical activity and different questionnaires in subsequent HUNT waves. However, physical activity as measured in H1 and H2 shows associations with morbidity and mortality as expected.^{8 9 32 33} Another limitation was that we did not have detailed information about diet in this population. Although a decline in energy expenditure and over-nutrition may explain the increasing prevalence of obesity, it is suggested that the main reason for increased body weight is declined physical activity levels rather than increased energy intake.³⁴ However, dietary habits are hard to analyse in population studies, and food data from large scale epidemiological studies should be interpreted with caution.³⁵

Our data do not include waist circumference in H1. As weight per se does not distinguish between fat and muscle, and muscle tissue has a higher weight density than fat, we argue that the effects of physical activity could be higher if we could assess changes in waist circumference and not merely weight change. Finally, as the participants in the study were primarily Caucasian, reflecting the inhabitants of Nord-Trøndelag County during the study period, the generalisability of the study findings may be limited.

In conclusion, being physically active mitigated weight gain in men and women above 20 years of age in a dose–response manner. Only activity levels exceeding the current recommendations of physical activity for health benefits, that is, ≥ 150 min of moderate-intensity or ≥ 60 min of vigorous activity each week, were associated with lower OR of meaningful weight gain over

What are the new findings?

- ▶ Physical activity above the current recommendations for health benefits, that is, ≥ 150 min of moderate-intensity or ≥ 60 min of vigorous activity each week, was associated with lower OR of meaningful weight gain over time.
- ▶ Maintaining a level of physical activity above the current recommendations over the adult lifespan was associated with 3.5 and 5.6 fewer kg of weight gain compared with staying inactive in men and women, respectively.
- ▶ Physical activity level and weight gain were inversely associated in a dose-dependent way in all age groups, in men and women.

How might it impact on clinical practice in the near future?

- ▶ Healthcare providers who advise patients about physical activity should be aware of the level of physical activity needed to prevent long-term weight gain. The amount required is greater than the ‘guideline’ amount for health benefits.
- ▶ Although the recommended level of physical activity for health benefits reduces the risk of chronic diseases, greater levels of physical activity are required to prevent weight gain; this should be incorporated into future physical activity recommendations.
- ▶ Policy-makers should therefore know of the requested level of physical activity to prevent weight gain for public health initiatives.

time. Although the recommended level of physical activity for health benefits reduces the risk of chronic diseases, the additional impact of physical activity volume and intensity for obesity prevention warrants further investigations, and should be incorporated into future recommendations.

Acknowledgements The Nord-Trøndelag Health Study (The HUNT Study) is a collaboration between HUNT Research Centre (Faculty of Medicine, Norwegian University of Science and Technology NTNU), Nord-Trøndelag County Council, Central Norway Health Authority, and the Norwegian Institute of Public Health. We are indebted to the participants of the HUNT study.

Contributors TM had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. TM, UW and JN were involved in the study concept and design and drafting of the manuscript. TM, UW, SL and JN were involved in analysis and interpretation of the data and critical revision of the manuscript for important intellectual content. TM and JN were involved in acquisition of data. TM, SL and JN were involved in statistical analysis. TM and JN were involved in study supervision. UW obtained funding.

Funding The present study was supported by grants from K.G.Jebesen Foundation, and by the Norwegian Fund for Postgraduate Training in Physiotherapy.

Competing interests None.

Ethics approval The study was approved by the Regional Committee for Medical and Health Research Ethics (REK 2012/2196).

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Flegal KM, Carroll MD, Kit BK, *et al.* Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA* 2012;307:491–7.
- 2 von Ruesten A, Steffen A, Floegel A, *et al.* Trend in obesity prevalence in European adult cohort populations during follow-up since 1996 and their predictions to 2015. *PLoS ONE* 2011;6:e27455.
- 3 Manson JE, Colditz GA, Stampfer MJ, *et al.* A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med* 1990;322:882–9.
- 4 Visscher TL, Seidell JC. The public health impact of obesity. *Ann Rev Public Health* 2001;22:355–75.
- 5 Whitlock G, Lewington S, Sherliker P, *et al.* Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009;373:1083–96.
- 6 Haskell WL, Lee IM, Pate RR, *et al.* Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation* 2007;116:1081–93.
- 7 Nelson ME, Rejeski WJ, Blair SN, *et al.* Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Circulation* 2007;116:1094–105.
- 8 Wisloff U, Nilsen TI, Droyvold WB, *et al.* A single weekly bout of exercise may reduce cardiovascular mortality: how little pain for cardiac gain? ‘The HUNT study, Norway’. *Eur J Cardiovasc Prev Rehabil* 2006;13:798–804.

- 9 Moholdt T, Wisloff U, Nilsen TI, *et al.* Physical activity and mortality in men and women with coronary heart disease: a prospective population-based cohort study in Norway (the HUNT study). *Eur J Cardiovasc Prev Rehabil* 2008;15:639–45.
- 10 King NA, Hopkins M, Caudwell P, *et al.* Beneficial effects of exercise: shifting the focus from body weight to other markers of health. *Br J Sports Med* 2009;43:924–7.
- 11 Lee IM, Djousse L, Sesso HD, *et al.* Physical activity and weight gain prevention. *JAMA* 2010;303:1173–9.
- 12 Hankinson AL, Daviglius ML, Bouchard C, *et al.* Maintaining a high physical activity level over 20 years and weight gain. *JAMA* 2010;304:2603–10.
- 13 Shiroma EJ, Sesso HD, Lee IM. Physical activity and weight gain prevention in older men. *Int J Obes (Lond)* 2012;36:1165–9.
- 14 Krokstad S, Langhammer A, Hveem K, *et al.* Cohort profile: the HUNT Study, Norway. *Int J Epidemiol* 2012;42:968–77.
- 15 Kurtze N, Rangul V, Hustvedt BE, *et al.* Reliability and validity of self-reported physical activity in the Nord-Trøndelag Health Study: HUNT 1. *Scand J Public Health* 2008;36:52–61.
- 16 Kurtze N, Rangul V, Hustvedt BE, *et al.* Reliability and validity of self-reported physical activity in the Nord-Trøndelag Health Study (HUNT 2). *Eur J Epidemiol* 2007;22:379–87.
- 17 Kumanyika SK, Obarzanek E, Stettler N, *et al.* Population-based prevention of obesity: the need for comprehensive promotion of healthful eating, physical activity, and energy balance: a scientific statement from American Heart Association Council on Epidemiology and Prevention, Interdisciplinary Committee for Prevention (formerly the expert panel on population and prevention science). *Circulation* 2008;118:428–64.
- 18 St Jeor ST, Brunner RL, Harrington ME, *et al.* A classification system to evaluate weight maintainers, gainers, and losers. *J Am Diet Assoc* 1997;97:481–8.
- 19 Owens JF, Matthews KA, Wing RR, *et al.* Can physical activity mitigate the effects of aging in middle-aged women? *Circulation* 1992;85:1265–70.
- 20 Littman AJ, Kristal AR, White E. Effects of physical activity intensity, frequency, and activity type on 10-y weight change in middle-aged men and women. *Int J Obes (Lond)* 2005;29:524–33.
- 21 Waller K, Kaprio J, Kujala UM. Associations between long-term physical activity, waist circumference and weight gain: a 30-year longitudinal twin study. *Int J Obes (Lond)* 2008;32:353–61.
- 22 Droyvold WB, Holmen J, Midthjell K, *et al.* BMI change and leisure time physical activity (LTPA): an 11-y follow-up study in apparently healthy men aged 20–69 y with normal weight at baseline. *Int J Obes Relat Metab Disord* 2004;28:410–17.
- 23 Wenche DB, Holmen J, Kruger O, *et al.* Leisure time physical activity and change in body mass index: an 11-year follow-up study of 9357 normal weight health women 20–49 years old. *J Womens Health (Larchmt)* 2004;13:55–62.
- 24 Ball K, Owen N, Salmon J, *et al.* Associations of physical activity with body weight and fat in men and women. *Int J Obes Relat Metab Disord* 2001;25:914–19.
- 25 Wier LT, Ayers GW, Jackson AS, *et al.* Determining the amount of physical activity needed for long-term weight control. *Int J Obes Relat Metab Disord* 2001;25:613–21.
- 26 Shetty P. Energy requirements of adults. *Public Health Nutr* 2005;8:994–1009.
- 27 Owen N, Salmon J, Koohsari MJ, *et al.* Sedentary behaviour and health: mapping environmental and social contexts to underpin chronic disease prevention. *Br J Sports Med* 2014;48:174–7.
- 28 Hu G, Tuomilehto J, Silventoinen K, *et al.* The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes (Lond)* 2005;29:894–902.
- 29 Leitzmann MF, Park Y, Blair A, *et al.* Physical activity recommendations and decreased risk of mortality. *Arch Intern Med* 2007;167:2453–60.
- 30 Hu G, Tuomilehto J, Silventoinen K, *et al.* Joint effects of physical activity, body mass index, waist circumference and waist-to-hip ratio with the risk of cardiovascular disease among middle-aged Finnish men and women. *Eur Heart J* 2004;25:2212–19.
- 31 Connor Gorber S, Tremblay M, Moher D, *et al.* A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007;8:307–26.
- 32 Stensvold D, Nauman J, Nilsen TI, *et al.* Even low level of physical activity is associated with reduced mortality among people with metabolic syndrome, a population based study (the HUNT 2 study, Norway). *BMC Med* 2011;9:109.
- 33 Nilsen TI, Romundstad PR, Vatten LJ. Recreational physical activity and risk of prostate cancer: a prospective population-based study in Norway (the HUNT study). *Int J Cancer* 2006;119:2943–7.
- 34 Pescatello LS, VanHeest JL. Physical activity mediates a healthier body weight in the presence of obesity. *Br J Sports Med* 2000;34:86–93.
- 35 Liu K. Statistical issues related to semiquantitative food-frequency questionnaires. *Am J Clin Nutr* 1994;59:262S–5S.