



Editor's choice
Scan to access more
free content

Competing with injuries: injuries prior to and during the 15th FINA World Championships 2013 (aquatics)

Margo Mountjoy,^{1,2,3} Astrid Junge,^{4,5,6} Sarah Benjamin,⁷ Kevin Boyd,^{2,8} Mohamed Diop,² David Gerrard,^{2,9} Cees-Rein van den Hoogenband,² Saul Marks,^{2,10} Enrique Martinez-Ruiz,¹¹ Jim Miller,^{2,12} Kyriakos Nanousis,² Farhad Moradi Shahpar,^{2,13} Jose Veloso,^{2,14} William van Mechelen,¹⁵ Evert Verhagen^{15,16}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2014-093991>).

For numbered affiliations see end of article.

Correspondence to

Dr Margo Mountjoy; FINA c/o Michael G. DeGroot School of Medicine, McMaster University Waterloo Regional Campus, 10-B Victoria Street South, Kitchener, Ontario, Canada N2G 1C5; mmsportdoc@mcmaster.ca

Accepted 26 September 2014
Published Online First
17 October 2014

ABSTRACT

Background Injury and illness surveillance is the foundation for the development of prevention strategies.

Objective To examine injuries among the aquatic disciplines in the 4 weeks prior to and during the 2013 FINA World Championships.

Methods The study was comprised of two components: (1) a retrospective athlete survey recording injuries in the 4 weeks prior to the Championships and (2) a prospective recording of injuries and illnesses by the medical teams of the participating countries and the local host medical team.

Results One-third of the 1116 responding athletes reported an injury/physical complaint in the 4 weeks prior to the Championships. Significantly more women (36.7%) than men (28.6%) reported injuries. Divers reported the highest rate of injury/physical complaints (55.7%). At the start of the Championships, 70% of injured respondents (n=258) were still symptomatic; however, full participation was expected by 76%. During the Championships, 186 new injuries were reported (8.3/100 registered athletes) with the highest injury incidence rate in water polo (15.3/100 registered athletes). The most common injured body part was the shoulder (21%). A total of 199 illnesses were reported during the Championships (9.0/100 registered athletes) with the most common diagnosis of illness being gastrointestinal infection. Environmental exposure (allergy, otitis and jellyfish stings) was responsible for 27% of all illnesses in open water swimming.

Conclusions Injuries pose a significant health risk for elite aquatic athletes. A prospective study would improve understanding of out-of-competition injuries. Future injury and illness surveillance at FINA World Championships is required to direct and measure the impact of prevention strategies.

INTRODUCTION

From the recreational to the elite level, aquatic sports are universally popular around the world. In 2013, FINA, the international federation governing aquatic sports, held its 15th World Championships in Barcelona, Spain involving 2223 athletes from 177 countries competing in 68 events in the disciplines of swimming, water polo, diving, synchronised swimming and open water swimming. The 30 m platform diving event was introduced at these Championships.

FINA implemented its first injury surveillance at its World Championships in Rome 2009.¹ Water

polo was included in the injury surveillance study of team sports during the 2004 Olympic Games,² and all aquatic disciplines were surveyed in the 2008³ and 2012⁴ Olympic Games. Illness surveillance was implemented in the FINA World Championships 2009¹ and in the 2012 Olympic Games.⁴

Results from the 2009 survey revealed that overuse (37.5%) was the most common mechanism for injury.¹ Similarly, at the 2012 Olympic Games, diving had the highest incidence of overuse injuries (73%) compared to all sports.⁴ The injury registration methodology utilised in the FINA and IOC injury surveillance studies, however, only records newly acquired injuries without capturing injuries incurred during training. Athletes may continue to train and compete despite injury and other physical complaints, especially in the time period before major competitions.

To facilitate the development of future prevention strategies for aquatic athletes, this study (1) assessed the presence and characteristics of injuries and physical complaints in the 4 weeks prior to and at the start of the FINA World Championships 2013 and (2) described the incidence and nature of new-onset injuries and illnesses incurred during competition.

METHODS

The study was implemented at the 2013 FINA World Championships in Barcelona, Spain. The Championships consisted of competitions in six disciplines, and the study population included all registered athletes. Informed consent was obtained via the athlete entry form. Ethical approval was obtained from the University of Oslo, School of Medicine Ethical Committee and the Col·legi Oficial de Metges de Barcelona.

The study involved two components: (1) a retrospective questionnaire on injuries and physical complaints in the 4 weeks prior to and at the start of the Championships to be completed by the athletes and (2) a prospective survey of new-onset injuries and illnesses reported by the local organising committee (LOC) and national medical teams.

Retrospective questionnaire on prior complaints

A retrospective survey on athletes' injuries and physical complaints during the 4 weeks prior to the Championships was developed to capture the presence, number, location and duration of injuries and



CrossMark

To cite: Mountjoy M, Junge A, Benjamin S, et al. *Br J Sports Med* 2015;**49**:37–43.

physical complaints, as well as their effect on training and competition performance (see online supplementary appendix 1). It included a modified, extended version of the four questions of the Oslo Sports Trauma Research Centre questionnaire.⁵

Athletes were asked to report all injuries or other physical complaints that they experienced in the 4 weeks prior to the Championships, regardless of the consequences for participation in normal training and/or competition. These outcomes are subsequently referred to as 'prior complaints'. The questionnaire was available in English, French and Spanish. Athletes received the questionnaires from their team physician, and returned them prior to or during the event either in person to a member of the research team or via secure locked drop boxes located in the training and competition venues and the FINA office.

Prospective survey on new-onset injuries and illnesses

The prospective injury and illness surveillance was based on the IOC protocol,⁶ previously implemented at the 2009 FINA World Championships.¹ The injury and illness report form was available in English, French, Spanish, Italian and Russian. Daily reporting by the team physician and LOC medical staff was accomplished electronically or via a secured drop box.

Implementation

The purpose and logistics of the study were communicated electronically to the National Federations and team physicians registered in the FINA database 3 months prior to the Championships. The FINA Bureau Sports Medicine Liaison (MM) undertook a pre-event site visit to train the LOC medical staff. Prior to the opening ceremony and prior to the start of the second week, an educational session was held for all national federation and LOC physicians. The study logistics were also presented at the technical meetings of each discipline.

During the competitions, members of the FINA Study Group attended every competition event to encourage reporting compliance. Injury and illness data gathered by the LOC from the training/competition sites, medical clinics, athlete hotels and the Championship hospital network were obtained daily from the Chief Medical Officer's utilisation report database. There was a daily review of the quality of the data and an analysis of reporting compliance. The medical staff of non-compliant countries were subsequently contacted to encourage reporting. Duplication of data entries of injuries and illnesses was resolved by the consensus of MM and AJ.

Definitions and calculation of exposure data and incidences

A discipline was defined as one of the six aquatic sports (swimming, diving, high diving, open water swimming, synchronised swimming and water polo). An event was defined as a competition activity if medals were awarded (eg, 1, 3 and 10 m diving). The number of competing athletes in the individual events was calculated from the official start lists.⁷ Athletes starting in different events were included for each event. Athletes who did not finish or were disqualified were included as competing athletes.

The number of starts was calculated using the official results list.⁷ For the individual sports, all results (including disqualifications and athletes who did not finish) were each included as a start; for example, an athlete competing in the heats, semifinals and finals of a swimming event was counted as three starts. Athletes who started in different disciplines (eg, swimming and open water swimming) were counted for each discipline. For swimming relays, synchronised swimming events, synchronised diving and open water swimming team events, the number of starts was calculated by multiplying the number

Table 1 Population characteristics and exposure during the FINA World Championships 2013

	Swimming	High diving	Diving	Synchronised swimming	Open water	Water polo	Total
Number of registered athletes (male/female)	1071 (608/463)	21 (15/6)	228 (116/112)	306 (0/306)	171 (96/75)	426 (211/215)	2223 (1046/1177)
Mean age of athletes (SD)	21.5 (3.84)	30.0 (4.48)	21.7 (4.03)	20.1 (3.50)	22.6 (4.58)	25.0 (4.79)	22.2 (4.42)
Number of events	40	2	10	7	7	2	68
Number of competing athletes (male/female)	2222 (1212/1010)	20 (14/6)	369 (185/184)	643 (0/643)	354 (198/156)	416 (208/208)	4024 (1817/2207)
Number of starts (male/female)	3102 (1652/1450)	20 (14/6)	609 (305/304)	1039 (0/1039)	354 (198/156)	1232 (616/616)	6356 (2785/3571)
Duration in days (male/female)	8 (8/8)	3 (2/1)	9 (9/9)	8 (0/8)	5 (4/4)	14 (7/7)	17 (17/17)
Number of athletes' competition days	8568	63	2052	2448	855	5964	19 950

Table 2 Number and characteristics of responders, injuries and complaints of in the 4 weeks prior to and at the start of the FINA World Championships 2013

	Swimming	High diving	Diving	Synchronised swimming	Open water	Water polo	Total*
Number of responding athletes (# male/female)	464 (250/214)	17 (14/3)	88 (38/50)	122 (0/122)	87 (45/42)	329 (153/176)	1116 (500/610)
Percentage of responders (male/female)	43.3 (41.1/46.2)	81.0 (93.3/50.0)	38.6 (32.8/44.6)	39.9 (0/39.9)	50.9 (46.9/56.0)	77.2 (72.5/81.9)	50.2 (47.8/51.8)
Percentage of responders with prior complaint (male/female)	23.5 (21.6/25.7)	23.5 (28.6/0)	55.7 (57.9/54.0)	42.6 (0/42.6)	17.4 (15.6/19.0)	41.9 (36.6/46.6)	33.0 (28.6/36.7)
Mean age of responders (SD)	21.9 (3.84)	29.3 (4.82)	21.4 (3.87)	19.7 (3.27)	22.1 (4.77)	24.5 (4.23)	22.5 (4.35)
Mean age of injured athletes (SD)	22.1 (4.08)	29.3 (5.74)	20.9 (3.69)	18.9 (3.06)	23.8 (4.51)	24.2 (3.85)	22.4 (4.31)
Participation in previous 4 weeks							
Full (%)	84.0	75.0	62.3	78.4	80.0	72.9	76.0
Reduced (%)	14.2	0	34.7	19.6	20.0	26.3	22.3
Unable (%)	1.8	25.0	3.0	2.0	0	0.8	1.7
Average number of body parts (SD)	1.28 (0.80)	1.25 (0.50)	1.17 (0.52)	1.65 (0.93)	1.53 (0.92)	1.43 (0.85)	1.38 (0.82)
Days of limited performance in previous 4 weeks							
0 (%)	15.6	25.0	20.4	18.4	50.0	13.3	17.0
1-3 (%)	18.8	-	13.0	20.4	-	17.2	16.7
4-7 (%)	38.5	50.0	26.1	22.4	8.3	26.6	29.2
8-14 (%)	14.6	-	15.0	24.5	8.3	14.8	15.8
15-27 (%)	7.3	-	15.0	6.1	25.0	13.3	11.0
28-30 (%)	5.2	25.0	10.5	8.2	8.3	14.8	10.4
Median days (IQR)*	5.0 (8.0)	6.0 (8.5)	5.5 (11)	6.0 (11)	2.0 (20)	7.0 (12)	6.0 (11.5)
Symptoms at start of competition							
None (%)	34.7	-	20.8	26.9	57.0	31.6	30.9
Minor (%)	48.7	50.0	54.2	42.4	21.5	48.6	47.5
Moderate (%)	15.8	-	20.8	26.9	14.4	16.9	18.3
Major (%)	0.8	25.0	2.1	1.9	7.1	2.9	2.5
No participation (%)	-	25.0	2.1	1.9	-	-	0.8
Performance affected at start of competition							
Not at all (%)	55.5	25.0	43.8	50.0	64.3	55.9	53.4
Minor (%)	37.6	-	33.4	40.4	7.1	31.6	33.4
Moderate (%)	6.9	25.0	12.5	5.8	21.5	8.1	8.7
Major (%)	-	25.0	8.3	-	7.1	4.4	3.4
No participation (%)	-	25.0	2.0	3.8	-	-	1.1

*Sum might be less than the total due to missing values.

of athletes on a team by the number of entries. For water polo, the number of starts was calculated as the number of player-matches (2 teams \times 7 players \times the number of matches played).

The number of athlete competition days was calculated by multiplying the number of registered athletes in each discipline by the total number of days of competition for the respective discipline as per the official competition schedule.⁷

Calculation of incidences and data analysis

Incidence rates were calculated as the number of injuries or illnesses per 100 registered or competing athletes in the respective discipline.⁶

Data were processed using Excel and SPSS. The statistical methods applied were descriptive statistics, cross tabulations, χ^2 and t tests. For incidence rates, 95% CIs were calculated. Statistical significance was accepted at a p value of 0.05 or lower.

RESULTS

Number and exposure of athletes

A total of 2223 athletes from 177 countries registered for the 2013 FINA World Championships that resulted in almost 20 000 athlete competition days. As most athletes, except high divers and water polo players, competed in more than one event, the number of competing athletes was almost double the number of registered athletes. Overall, more female (53%) than male athletes (47%) participated. The athletes' ages range from 12 to 41 years with an average of 22.2 years (SD 4.42). For further details on the characteristics of the athletes and their exposure see table 1.

Retrospective questionnaire on prior complaints

A total of 1116 athletes (610 women and 500 men) completed the retrospective survey of prior complaints, representing an overall response rate of 50.2%, varying by discipline from 38.6%

(diving) to 81% (high diving). Survey responders and non-responders were similar in age and gender, but were statistically different with regard to distribution by discipline (table 2).

Approximately one-third of the responding athletes (n=368; 33%) reported a physical complaint in the 4 weeks prior to the Championships. The duration of prior complaints varied from 7 days to 10 years with a median duration of 85.3 days. The mean age of athletes with and without prior complaints was similar. Significantly more females (n=224; 36.7%) than males (n=143; 28.6%) reported prior complaints ($\chi^2=7.83$; $p<0.005$), and the proportion of athletes with a prior complaint differed with statistical significance between disciplines ($\chi^2=68.2$; $p<0.001$). More than half of the divers (55.7%) reported prior complaints compared to only 17.4% of open water swimmers.

More than three quarters of athletes with prior complaints (n=273; 76%) reported full participation in training in the 4 weeks leading up to the Championships, but only 57 (17%) reported that their performance had not been affected due to injury. The median number of days of limited sport participation was 6. High diving and water polo had the greatest number of days of limited performance.

At the start of the Championships, less than one-third of athletes with prior complaints (30.9%) reported no current symptoms, 47.5% minor symptoms, 18.3% moderate symptoms and 2.5% severe symptoms. More than half (53.4%) reported no effect on performance from their prior complaints at the beginning of the Championships. In contrast, four athletes were unable to participate in the Championships and 1.4%, despite being severely affected, still competed.

Prospective survey on new-onset injuries and illnesses during the Championships

The medical staff from 55 (31.1%) countries submitted 587 injury and illness report forms covering a total of 1571 athletes

Table 3 Number and incidence (95% CI) of injuries and illnesses during the FINA World Championships 2013

	Swimming	High diving	Diving	Synchronised swimming	Open water	Water polo	Total (95% CI)*†
All injuries (male/female)	65 (31/34)	1 (1/–)	26 (10/16)	5 (–/5)	20 (11/9)	65 (33/32)	186 (88/98)
Per 100 registered athletes	6.1	4.8	11.4	1.6	11.7	15.3	8.3 (1.19)
Per 100 athletes' competition days	0.7	1.5	1.3	0.2	2.3	1.1	0.5
Training injuries	46	0	12	1	4	18	81
Per 100 registered athletes	4.3	0	5.3	0.3	2.3	4.2	3.6 (0.78)
Competition injuries	16	1	14	4	16	47	98
Per 100 competing athletes	0.7	5.0	3.8	0.6	4.5	11.3	2.4 (0.48)
Per 100 starts	0.5	5.0	2.3	0.4	4.5	3.8	1.5 (0.3)
Time-loss injuries (male/female)	12 (2/10)	0	7 (2/5)	0	2 (1/1)	11 (9/2)	32 (14/18)
Per 100 registered athletes	1.1	0	3.1	0	1.2	2.6	1.4 (0.49)
Time-loss training injuries	8	0	4	0	0	4	16
Per 100 registered athletes	0.75	0	1.75	0	0	0.94	0.72 (0.35)
Time-loss competition injuries	4	0	3	0	2	7	16
Per 100 competing athletes	0.18	0	0.81	0	0.56	1.68	0.40 (0.2)
Per 100 starts	0.1	0	0.5	0	0.6	0.6	0.3
All illnesses (male/female)	95 (51/44)	0	9 (6/3)	23 (–/23)	37 (21/16)	35 (14/21)	199 (92/107)
Per 100 registered athletes	8.9	0	3.9	7.5	21.6	8.2	9.0 (1.25)
Per 100 athletes' competition days	1.1	0	0.4	0.9	4.3	0.6	0.5
Time-loss illnesses (male/female)	8 (4/4)	0	3 (2/1)	2 (–/2)	2 (1/1)	10 (5/5)	25 (12/13)
Per 100 registered athletes	0.75	0	1.3	0.7	1.1	2.3	1.1 (0.43)
Per 100 athletes' competition days	0.1	0	0.1	0.1	0.2	0.2	0.1

*Sum might be less than the total due to missing values.

†If one athlete started in different disciplines, they were counted for each discipline.

(70.7%; table 3). A total of 186 injuries were reported, representing an incidence of 8.3/100 registered athletes (95% CI ± 1.19). The highest injury incidence was observed in water polo, followed by open water swimming and diving, and the lowest incidence was seen in synchronised swimming. The incidence of injuries was higher in training (3.61; 95% CI ± 0.78) than in competition (2.4; 95% CI ± 0.48). No other differences were found. Thirty-two injuries (17%) were reported to result in time loss, equivalent to an incidence of 1.4/100 athletes (95% CI ± 0.49). Information on time loss was missing for 58 (31.2%) injuries. The rate of time-loss injuries was lower in male athletes (1.34; 95% CI ± 0.7) than in female athletes (1.53; 95% CI ± 0).

The most common injured body part was the shoulder (n=39; 21%; figure 1). Almost half (49%) of the shoulder injuries and two-thirds of the time-loss shoulder injuries were reported in swimming. Water polo was responsible for




Figure 1 Injury location for new-onset injuries during the FINA World Championships 2013.

two-thirds (n=23) of all injuries to the head/neck. The most common injury type was contusion (n=38; 20.4%), and the most frequent injury type was tendinosis of the shoulder (n=17). The most common cause of injury was contact with another athlete (n=46; 24.7%). Eight per cent of the reported injuries resulted in time loss of 1–2 weeks duration, and one-quarter of all time-loss injuries were to the shoulder (n=28). Data on time loss were missing in 53 (28%) cases.

A total of 199 illnesses were reported during the Championships, resulting in an incidence of 9/100 registered athletes (95% CI ± 1.25). There was no significant statistical difference between the number of illnesses reported in male (8.8; 95% CI ± 1.8) and female (9.1; 95% CI ± 1.72) athletes. The highest rate of illness was reported in open water swimmers while no illnesses were reported in high divers (table 3). The most common diagnosis of illness was gastrointestinal infection, with environmental exposure resulting in allergic reactions, otitis and jellyfish stings in open water swimming accounting for 27% of all reported illnesses (figure 2). Twenty-five illnesses were reported to result in time loss, resulting in a rate of 1.1 (95% CI ± 0.43) time-loss injuries per 100 registered athletes. Data on time loss were missing in 64 (32%) cases. The highest rate of time-loss illnesses was observed in water polo.

DISCUSSION

Accurate injury surveillance is a fundamental component of effective risk management in sport.^{8–11} Despite the fact that out-of-competition prospective cohort studies are recommended to monitor injury patterns and risk,^{12–15} no studies of this type have been undertaken in aquatic sport disciplines. Two studies, one prospective study monitoring shoulder pain in Australian swimmers¹⁶ and one descriptive retrospective epidemiological study¹⁷ in NCAA swimming, have been published.

Retrospective questionnaire of prior complaints

The retrospective component of this study is the first to evaluate the presence of out-of-competition injuries in international, elite aquatic athletes. Despite employing a narrow recall period of 4 weeks and a research population comprising only elite athletes, limitations to the accuracy of the retrospective survey exist with a potential for underestimation of injuries related to a potential recall bias.^{18 19} The more severe injuries of athletes who are subsequently unable to qualify for the FINA World Championships were not captured in this study. As such, the injury rates reported in this study are most likely an underestimation of the actual injury rates in elite aquatic athletes. In contrast, a strength of this survey was the recording of athlete symptoms rather than diagnoses, which allowed for greater capture of performance affecting physical complaints by bypassing the potential reporting bias of third party recorders who may interpret differently a recordable problem. This approach provided a more accurate interpretation of the effect that physical complaints may have on athletic performance.^{13 14}

The survey responses demonstrated that almost a third of all athletes reported complaints, with 70% of them reporting prior complaints symptomatic at the commencement of the Championships. Despite these findings, the majority still participated in training, stating that their athletic performance was affected by their prior complaint at the start of the competition. This finding is consistent with the literature outlining that athletes across sport disciplines continue to train despite injury¹³ and demonstrates the need for interventions to address the prevention of non-time-loss injuries. Further research in aquatic




Figure 2 Diagnosis of illnesses during the FINA World Championships 2013.

sports is required to determine whether symptomatic athletes are more vulnerable as previous injury is a known risk factor for injury.⁹ To achieve this end, prospective out-of-competition surveillance together with the standard in-competition protocol is recommended.

Prospective survey on new-onset injuries

The incidence of all and of time-loss injuries during the 2013 FINA World Championships was significantly higher than in 2009.¹ This trend may be due to an increasingly tougher competition or to a busier competition calendar with a consequent decreased recovery period and a subsequent increased injury risk. It may also be attributed to the improved familiarity of team physicians with the injury surveillance system since a higher response rate was achieved in 2013 compared to the

FINA World Championships 2009.¹ A similar trend was observed in the Olympic Games.^{3 4 20} Ongoing injury surveillance is required to decrease this reporting bias.

Consistent with previous published research,^{1 16 21–25} the shoulder was the most commonly injured body part in aquatic sports. Despite the various theories of risk factors and injury mechanisms,^{16 25 26} shoulder injury prevention programmes for aquatic sports have not been validated in the scientific literature. The findings of the present study illustrate the need for a validated shoulder injury prevention programme in the aquatic disciplines.

In the current study, the head and neck regions were the second most commonly reported injury locations. This unexpected finding was influenced by water polo, which reported more than 60% of all head and neck injuries. Water polo data also reflected the highest injury risk of all the aquatic disciplines with a higher number of head and neck injuries reported than shoulder injuries. A pragmatic approach to injury prevention in water polo would include a review of the current rules, de-emphasising contact, highlighting fair play and supporting the capacity of referees to sanction. These measures have been implemented in team sports such as football and rugby.^{27–32}

Prospective survey on illnesses

Statistically significantly more illnesses were reported during the 2013 FINA World Championships than in 2009 (7.1/100 athletes; 95% CI 1.03).¹ This could have been influenced by the environmental risk variability between Barcelona (2013) and Rome (2009), and also as a result of the improved athlete healthcare provision and improved reporting quality. Consistent with other sports, infection was the leading cause of illness.^{4 20 33–36} However, less upper respiratory infections were reported in 2013 (n=36; 18%) than in 2009¹ (n=91; 50.3%), possibly attributed to the presence of hand sterilising stations on site and the education of athletes on personal hygiene.³⁷

Of particular concern was the reported frequency of otitis in all athletes (15% of all illnesses) and jellyfish stings in open water swimmers (38% of illnesses in open water swimmers). These results demonstrate the need for the implementation of targeted preventive measures. The environmental risk posed by jellyfish warrants a closer evaluation of open water venues to develop risk reduction measures and the provision of appropriate medication for the treatment of potential anaphylactic emergencies.

CONCLUSION

This study demonstrates the need to investigate out-of-competition (training) injury patterns in aquatic sports through prospective injury surveillance in aquatic sports. In-competition injury and illness surveillance should continue to gain further insight into trends and to evaluate the efficacy of prevention interventions. While the disciplines of swimming, diving and synchronised swimming have low injury rates relative to other sports, water polo has a uniquely higher propensity for injury distinctively affecting the head, neck and shoulder. Implementation of prevention interventions should target the discipline of water polo, shoulder and head injuries, as well as aquatic-specific illnesses such as jellyfish stings and otitis.

What are the new findings?

- ▶ One-third of all participating athletes reported having suffered physical complaints/injuries in the 4 weeks prior to the 2013 FINA World Championships.
- ▶ Diving and water polo had the highest incidence of physical complaints/injuries in the 4 weeks prior to the Championships relative to the other aquatic disciplines.
- ▶ Female aquatic athletes had a higher rate of physical complaints/injury than male aquatic athletes in the 4 weeks prior to the Championships.
- ▶ The incidence of new-onset injuries during the FINA World Championships was 8.3% with the highest incidence in water polo; and the illness incidence was 9% with the highest incidence in open water swimming.
- ▶ The most common injuries were contusions of the head and overuse injuries of the shoulder, and the most common illnesses were gastrointestinal and respiratory infections.
- ▶ How might it impact on clinical practice in the near future?
- ▶ Injury surveillance programmes in all aquatic disciplines should expand to capture data during the training season specifically in the time period leading up to a major competition.
- ▶ Prevention interventions should be developed to focus on injuries suffered during the out-of-competition (training) period.
- ▶ An analysis of the water polo rules should be implemented to address the injury patterns unique to this aquatic discipline.
- ▶ Injury prevention strategies in all aquatic disciplines should target shoulder and head injuries.
- ▶ Evaluation of environment factors in open water swimming is required.

Author affiliations

- ¹Department of Family Medicine, Michael G. DeGroot School of Medicine, McMaster University, Hamilton, Canada
²Fédération Internationale de Natation (FINA), Lausanne, Switzerland
³International Olympic Committee (IOC), Lausanne, Switzerland
⁴Fédération Internationale de Football Association (FIFA) Medical Assessment and Research Centre (F-MARC), Zürich, Switzerland
⁵Schulthess Klinik, Zürich, Switzerland
⁶Hamburg Medical School (MSH), Germany
⁷University of Southampton, Southampton, Great Britain
⁸University Hospitals of Leicester NHS Trust, Leicester, Great Britain
⁹Medicine Department, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand
¹⁰Department of Psychiatry, University of Toronto, Toronto, Canada
¹¹Catholic University of San Antonio, Murcia, Spain
¹²Medicine Department, University of Virginia, Charlottesville, and Virginia Commonwealth University, Richmond, Virginia, USA
¹³Department of Sport Medicine, Faculty of Physical Education and Sports Sciences, University of Isfahan, Isfahan, Iran
¹⁴Faculty of Biological Sciences, Catholic University of Uruguay, Montevideo, Uruguay
¹⁵Department of Public and Occupational Health, EMGO+ Institute for Health and Care Research, VU University Medical Center, Amsterdam, The Netherlands
¹⁶Australian Centre for Research into Injury in Sport and its Prevention (ACRISP), Federation University Australia, SMB Campus, Ballarat, Victoria, Australia

Acknowledgements The authors highly appreciate the co-operation of all team physicians and medical staff of the FINA World Championships 2013 who volunteered their time to collect the data for this project. They thank Dr Nuria Torres, chief medical officer of BCN2013 for her co-operation in providing the medical data from the host medical clinics and venues and Dr Kathrin Steffen for her assistance with obtaining Ethics Committee approval from the University of Oslo. The authors also gratefully acknowledge FINA for its support and funding of the study.

Contributors MM substantially contributed to conception and design, data collection, interpretation of results, drafting and revising the manuscript and final version to be published. AJ substantially contributed to conception and design, development of the questionnaire and data collection methods, data analysis and interpretation of results, drafting and revising the manuscript, and final approval of the version to be published. SB, JV, KB, MD, C-RvdH, SM, EM-R, KN, FMS substantially contributed to acquisition of the data and final approval of the manuscript to be published. DG contributed to acquisition of the data, revising the manuscript and final approval of the manuscript to be published. JM contributed to acquisition of the data and final approval of the version to be published. WvM contributed to the manuscript and final approval of the version to be published. EV contributed to data collection, revising the manuscript and final approval of the version to be published.

Competing interests None.

Ethics approval University of Oslo, School of Medicine Ethical Committee+Collegi Oficial de Metges de Barcelona.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Mountjoy M, Junge A, Alonso JM, *et al.* Sports injuries & illnesses in the 2009 FINA World Aquatic Championships. *Br J Sports Med* 2010;44:522–7.
- 2 Junge A, Langevoort G, Pipe A, *et al.* Injuries in team sport tournaments during the 2004 Olympic Games. *Am J Sports Med* 2006;34:565–76.
- 3 Junge A, Engebretsen L, Mountjoy M, *et al.* Sports injuries during the Summer Olympic Games 2008. *Am J Sports Med* 2009;37:2165–72.
- 4 Engebretsen L, Soligard T, Steffen K, *et al.* Sports injuries and illnesses during the London Summer Olympic Games 2012. *Br J Sports Med* 2013;0:1–8.
- 5 Clarsen B, Myklebust G, Bahr R. Development and validation of a new method for the registration of overuse injuries in sports injury epidemiology. *Br J Sports Med* 2013;47:495–502.
- 6 Junge A, Engebretsen L, Alonso JM, *et al.* Injury surveillance in multi-sport events—the IOC approach. *Br J Sports Med* 2008;42:413–21.
- 7 Omega Results. <http://www.omegatiming.com/Sport?Sport=AQ>. Retrieved 7 December 2013.
- 8 Fuller C, Drawer S. The application of risk management in sport. *Sports Med* 2004;34:349–56.
- 9 Fuller C. Managing the risk of injury in sport. *Clin J Sport Med* 2007;17:182–7.
- 10 Rogge J. An ounce of prevention? *Br J Sports Med* 2009;43:627.
- 11 Steffen K, Soligard T, Engebretsen L. Health protection of the Olympic athlete. *Br J Sports Med* 2012;46:466–70.
- 12 Fuller C, Bahr R, Dick R, *et al.* A framework for recording recurrences, reinjuries, and exacerbations in injury surveillance. *Clin J Sport Med* 2007;17:197–200.
- 13 Bahr R. No injuries, but plenty of pain? On the methodology for recording overuse symptoms in sports. *Br J Sports Med* 2009;43:966–72.
- 14 Clarsen B, Ronsén O, Myklebust G, *et al.* The Oslo Sports Trauma Research Center Questionnaire on Health Problems: a new approach to prospective monitoring of illness and injury in elite athletes. *Br J Sports Med* 2014;48:754–60.
- 15 Clarsen B, Bahr R, Heymans M, *et al.* The prevalence and impact of overuse injuries in five Norwegian sports: application of a new surveillance method. *Scand J Med Sci Sports* 2014. doi:10.1111/sms.12223
- 16 Walker H, Gabbe B, Wajswelner H, *et al.* Shoulder pain in swimmers: a 12-month prospective cohort study of incidence and risk factors. *Physical Therapy Sport* 2012;13:243–9.
- 17 Wolf B, Ebinger A, Lawler M, *et al.* Injury patterns in division I collegiate swimming. *Am J Sports Med* 2009;37:2037–42.
- 18 Junge A, Dvorak J. Influence of definition and data collection on the incidence of injuries in football. *Am J Sports Med* 2000;28:S40–6.
- 19 Bjørneboe J, Flørens T, Bahr R, *et al.* Injury surveillance in male professional football; is medical staff reporting complete and accurate? *Scand J Med Sci Sports* 2011;21:713–20.
- 20 Engebretsen L, Steffen K, Alonso JM, *et al.* Injuries and illnesses during the Winter Olympic Games 2010. *Br J Sports Med* 2010;44:772–80.
- 21 Wanivenhaus F, Fox A, Chaudhury S, *et al.* Epidemiology of injuries and prevention strategies in competitive swimmers. *Sports Health* 2012;4:246–51.
- 22 Webster M, Morris M, Galna B. Shoulder pain in water polo: a systematic review of the literature. *J Sci Med Sport* 2009;12:3–11.
- 23 Gaunt T, Maffulli N. Soothing suffering swimmers: a systematic review of the epidemiology, diagnosis, treatment and rehabilitation of musculoskeletal injuries in competitive swimmers. *Br Med Bull* 2012;103:45–88.
- 24 Johnson J, Sim F, Scott S. Musculoskeletal injuries in competitive swimmers. *Mayo Clin Proc* 1987;62:289–304.
- 25 Folland J, Archer G. Shoulder injuries in highly trained competitive swimmers. *Br J Sports Med* 2014;48:593.

- 26 Hibberd E, Myers J. Practice habits and attitudes and behaviors concerning shoulder pain in high school competitive club swimmers. *Clin J Sport Med* 2013;23:450–5.
- 27 Rynnänen J, Junge A, Dvorak J, *et al.* Foul play is associated with injury incidence: an epidemiological study of three FIFA World Cups (2002–2010). *Br J Sports Med* 2013;47:986–91.
- 28 Rynnänen J1, Dvorak J, Peterson L, *et al.* Increased risk of injury following red and yellow cards, injuries and goals in FIFA World Cups. *Br J Sports Med* 2013;47:970–3.
- 29 Andersen T, Engebretsen L, Bahr R. Rule violations as a cause of injuries in male Norwegian professional football: are the referees doing their job? *Am J Sports Med* 2004;32(1 Suppl):62S–8S.
- 30 Fuller C, Junge A, Dvorak J. An assessment of football referees' decisions in incidents leading to player injuries. *Am J Sports Med* 2004;32(1 Suppl):17S–22S.
- 31 Comstock R, Fields S. The fair sex? Foul play among female rugby players. *J Sci Med Sport* 2005;8:101–10.
- 32 Bjørneboe J, Bahr R, Dvorak J, *et al.* Lower incidence of arm-to-head contact incidents with stricter interpretation of the Laws of the Game in Norwegian male professional football. *Br J Sports Med* 2013;47:508–14.
- 33 Edouard P, Depiesse F, Branco P, *et al.* Analyses of Helsinki 2012 European Athletics Championships Injury and Illness Surveillance to Discuss Elite Athletes Risk Factors. *Clin J Sport Med* 2014;24:409–15.
- 34 Edouard P, Depiesse F, Hertert P, *et al.* Injuries and illnesses during the 2011 Paris European Athletics Indoor Championships. *Scand J Med Sci Sports* 2013;23:e213–18.
- 35 Alonso JM, Edouard P, Fischetto G, *et al.* Determination of future prevention strategies in elite track and field: analysis of Daegu 2011 IAAF Championships injuries and illnesses surveillance. *Br J Sports Med* 2012;46:505–14.
- 36 Alonso JM, Tscholl PM, Engebretsen L, *et al.* Occurrence of injuries and illnesses during the 2009 IAAF World Athletics Championships. *Br J Sports Med* 2010;44:1100–5.
- 37 Hanstad D, Rønson O, Andersen S, *et al.* Fit for the fight? Illnesses in the Norwegian team in the Vancouver Olympic Games. *Br J Sports Med* 2011;45:571–5.

Appendix # 1. Pre-existing Injury Survey

BCN 2013
BCN JO13

BARCELONA 2013
15th FINA WORLD
CHAMPIONSHIPS
©FINA 2011

SURVEY ON PRE-EXISTING INJURY



Date of birth: <small>dd/mm/yy</small>		Gender:	<input type="checkbox"/> Female	<input type="checkbox"/>
			Male	
Sport/Event:		Country:		

Dear Athlete,

FINA would like to learn more about **injuries or physical complaints** such as pain, ache, stiffness, swelling, instability/giving way, locking or other symptoms, you had in **the 4 weeks prior to this championship**, regardless of whether or not they have had major consequences for your participation in normal training and/or competition. Please answer the following questions. The results will help to improve the care for aquatic athlete and to prevent injury.

1. Did you have any injury/physical complaints in the 4 weeks prior to this championship?

- no (*you do not need to answer further questions*)
- yes, full participation, but with injury/physical complaints
- yes, reduced participation due to injury/physical complaints
- yes, Cannot participate due to injury/physical complaints

If you had several injuries/physical complaints, please begin with your worst problem, and describe the other problems separately on extra forms.

2. Please tick the location of injury/physical complaint referred to above

<input type="checkbox"/> Head/face	<input type="checkbox"/> Hand/fingers	<input type="checkbox"/> Thigh
<input type="checkbox"/> Neck	<input type="checkbox"/> Chest/ribs	<input type="checkbox"/> Knee
<input type="checkbox"/> Shoulder (including clavicle)	<input type="checkbox"/> Abdomen	<input type="checkbox"/> Lower leg
<input type="checkbox"/> Upper arm	<input type="checkbox"/> Thoracic spine	<input type="checkbox"/> Ankle
<input type="checkbox"/> Elbow	<input type="checkbox"/> Lumbar spine	<input type="checkbox"/> Foot/toes
<input type="checkbox"/> Forearm	<input type="checkbox"/> Pelvis and buttock	<input type="checkbox"/> Other (Specify _____)
<input type="checkbox"/> Wrist	<input type="checkbox"/> Hip and groin	

3. How did this injury/physical complaint first occur?

- Due to a clearly identifiable injury event, e.g. a collision or a fall
- Suddenly while performing normal training or competition (no obvious trauma)
- Gradual onset, no single event
- Other, not related to training or competing in aquatics

4. For how long have you been suffering from this injury/complaint? _____ years
_____ months _____ weeks

5. In the 4 weeks prior to this championship, ...	Please tick the appropriate box
...how much did you reduce your training volume due to this injury/physical complaint?	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> not at all minor moderate major no training
...how much did you modify the content / intensity of training due to this injury/physical complaint?	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> not at all minor moderate major no training
...how much did this injury/physical complaint affect your performance during training or competition?	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> not at all minor moderate major no participation
... how severe was this complaint during training or competition?	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> none minor moderate major no participation
... on how many days was your performance affected by this injury /complaint?	_____ days in the past 4 weeks

6. At the start of your competition at this championship ...	
... how severe was this complaint during training or competition?	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> none minor moderate major no participation
...how much was your performance affected due to this injury/physical complaint?	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> not at all minor moderate major no participation

Please ask the FINA researcher for additional forms, if needed, and return the filled in forms to the FINA research assistant or in the research drop box in the medical stations.

Thank you very much for your participation in this survey!