



Editor's choice
Scan to access more
free content

Preparticipation injury complaint is a risk factor for injury: a prospective study of the Moscow 2013 IAAF Championships

Juan-Manuel Alonso,^{1,2} Jenny Jacobsson,^{3,4} Toomas Timpka,³ Ola Ronsen,^{1,5} Alma Kajenienne,^{1,6} Örjan Dahlström,⁷ Armin Spreco,³ Pascal Edouard^{8,9,10}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2014-094359>).

For numbered affiliations see end of article.

Correspondence to

Dr Juan-Manuel Alonso, Sports Medicine Department, Aspetar, Qatar Orthopedics and Sports Medicine Hospital, PO Box 29222, Doha, Qatar; juan-manuel.alonso@aspetar.com

Accepted 7 February 2015
Published Online First
25 February 2015

ABSTRACT

Objectives To determine the health status of athletes before the start of an international athletics championship and to determine whether preparticipation risk factors predicted in-championship injuries.

Methods At the beginning of the 2013 International Association of Athletics Federations (IAAF) World Championships, all registered athletes (n=1784) were invited to complete a preparticipation health questionnaire (PHQ) on health status during the month preceding the championships. New injuries that occurred at the championships were prospectively recorded.

Results The PHQ was completed by 698 (39%) athletes; 204 (29.2%) reported an injury complaint during the month before the championships. The most common mode of onset of preparticipation injury complaints was gradual (43.6%). Forty-nine athletes in the study reported at least one injury during the championships. Athletes who reported a preparticipation injury complaint were at twofold increased risk for an in-championship injury (OR=2.09; 95% CI 1.16 to 3.77); p=0.014). Those who reported a preparticipation gradual-onset injury complaint were at an almost fourfold increased risk for an in-championship time-loss injury (OR=3.92; 95% CI 1.69 to 9.08); p=0.001). Importantly, the preparticipation injury complaint severity score was associated with the risk of sustaining an in-championship injury (OR=1.14; 95% CI 1.06 to 1.22); p=0.001).

Summary and conclusions About one-third of the athletes participating in the study reported an injury complaint during the month before the championships, which represented a risk factor for sustaining an injury during the championship. This study emphasises the importance of the PHQ as a screening tool to identify athletes at risk of injuries before international championships.

INTRODUCTION

During international outdoor athletics championships, the incidence and characteristics of new injuries have been well described.^{1–5} To further expand our knowledge, we aimed to identify possible risk factors for injury during the championships.

Athletes participating in championships are assumed to be healthy, that is, free of health problems at the start of a championship; however, this is not always the case. Athletes may have been injured and/or sick during preparation for the championships. In a recent pilot study of 74 athletes, 54% reported at least one health problem (an injury complaint was reported by 24%, an

illness complaint by 18% and a pain complaint by only 12%) during the 4 weeks before the 2013 European Athletics Indoor Championships.⁶ There have been no studies examining whether carrying injuries into competition predicts injury during competition.

As injury epidemiology has evolved, so have the ways to capture preparticipation health status.⁶ When focusing on injuries, we can collect data on several aspects of an athlete's self-reported injury complaints (eg, all complaints, time-loss complaints, gradual-onset/sudden-onset complaints and severity scores).^{7,8}

Therefore, we aimed to determine the health status (injury complaints) of athletes before the start of an international athletics championship. We hypothesised that certain types of preparticipation injury complaints would predict injury during the championship. Our methodological investigation was to investigate the strength of different measures of preparticipation injury complaints (all complaints, time-loss complaints, a complaint severity score) as well as gender, age and event group as predictors of injury during a championship.

METHODS

Study design and data collection procedures

A total cohort design was used for the study. At the beginning of the International Association of Athletics Federations (IAAF) World Athletics Championships in Moscow in 2013, all registered athletes (n=1784) were invited to complete a preparticipation health questionnaire (PHQ) including individual preparticipation information (personal and training characteristics and health status during the month preceding the championship).

The PHQ was developed by a group of experts consisting of scientists and practitioners (n=6) based on a previously published PHQ.⁶ The overall goal was to identify possible risk factors of sustaining injury during the championships by comparing data from the PHQ with those from injury surveillance during the championships.⁶ The PHQ was developed to collect information directly from the athletes regarding their personal characteristics (gender, age, country, height, weight, event group, time spent in training) and health status ('injuries or physical complaints', which will be called 'injury complaints' from this point on, and 'illnesses') during the month preceding the championships (online supplementary figure S1).⁶

The questions on the functional consequences of the injury complaint were formulated in line with a



CrossMark

To cite: Alonso J-M, Jacobsson J, Timpka T, et al. *Br J Sports Med* 2015;**49**:1118–1124.

questionnaire used for monitoring overuse sports injuries.^{9 10} An injury complaint severity score was calculated using items 3.1, 3.2, 3.3 and 3.4 of the PHQ inspired by the calculation of the Oslo Sports Trauma Research Centre (OSTRC) Overuse Injury Questionnaire severity score from Clarsen *et al*⁹ The response values 0–4 were allocated so that 0 represents no problems and 4 represents the maximum level for each question. An injury complaint severity score of 0 means no problem, and a score of 16 means maximal functional limitation. In the case of more than one injury complaint during the 4-week period, the athletes had to specify the main (worst) injury complaint in the PHQ and then report the additional number of injury complaints. The PHQ was available in a paper format (in English, French, Spanish, Russian), and distributed in the athlete's welcome bag, or by the team physicians or at the antidoping blood sample collection office. Athletes were asked to complete the questionnaire themselves or with the help of team physicians, if needed, and to return it to their team physician or to the designated desk in the warm-up area.

During the period of the championships (10–18 August 2013), newly incurred injuries were recorded by national medical teams (physicians and/or physiotherapists) and/or by physicians on the local organising committee (LOC) using the injury surveillance system (definitions and methods) previously used during international athletics championships.^{2–6 11} To maintain anonymity and enable linking of the PHQ and injury surveillance data, each participant in the study received a specific identification number.

All the national medical teams and LOC physicians were informed about the study 1 month before the championships by email and on the day before the championships during a medical and antidoping meeting.

Preparticipation injury complaints asked for by self-reporting were defined as “injuries or physical complaints (such as pain, ache, stiffness, swelling, instability/giving way, locking or other symptoms) that athletes had had in the four weeks prior to the championship, even if this had not had major consequences for athlete's participation in normal training and/or competition.” In-championship injury was defined as proposed by the International Olympic Committee consensus¹² and used during the three previous IAAF World Championships.^{2–4} An injury was defined as “all musculoskeletal injuries (traumatic and overuse) newly incurred during competition or training regardless of the consequences with respect to the athlete's absence from competition or training.”^{2 12} A time-loss injury was defined as one that “leads to the athlete being unable to take full part in athletics training and/or competition the day after the incident occurred.”¹² Similarly, for a preparticipation injury complaint, a reduced participation injury complaint was defined as one that “leads to the athlete being unable to take full part in athletics training and/or competition the day after the incident occurred”; it corresponds to the responses “Yes, reduced participation” and “Yes, I cannot participate” to question 1 of the PHQ and/or “Minor/Moderate/Major/No training” to questions 3.1 and/or 3.2.

Although the PHQ also included items on illness complaints, this study focuses on injuries: preparticipation injury complaints and in-championship injuries. Data on illness complaints were not included in the present analyses.

Confidentiality and ethical approval

The athletes' gender, date of birth and nationality were used only to avoid duplicate reporting from team and LOC physicians, and to provide descriptive data on athletes' characteristics.

Information about the purpose of the study and the procedure was provided to the athletes in their welcome bags or by medical teams. All athletes were free to refuse the use of their data. All PHQs and injury reports were stored in a locked filing cabinet and were made anonymous after the championships. The confidentiality of all information was ensured so that no individual athlete or national team could be identified. Ethical approval was obtained from the Saint-Etienne University Hospital Ethical Committee (Institutional Review Board Information: IORG0004981).

Data analysis

For the PHQ data, the response rate and completeness of the PHQs were assessed, and the athletes' characteristics and injury complaints were analysed using descriptive statistics.⁶ Analysis of the non-responders was performed by comparing their distribution of home continent, event group, gender and age with the final study group.

For the in-championship injury surveillance, the response rate, coverage, and injury incidence calculations and comparisons were analysed in accordance with previous injury surveillance studies in athletics^{4–6 11} using a list of athletes provided by the IAAF and the internet database (<http://www.iaaf.org/competitions/iaaf-world-championships/14th-iaaf-world-championships-4873>).


For the final risk factor analyses, the PHQ data were recoded into the four variables “any injury complaints” (y/n), “reduced participation injury complaints” (y/n), “gradual-onset injury complaints” (y/n) and the injury complaint severity score (0–16). We first performed analyses with simple models (logistic regression analyses with one explanatory variable) and thereafter analyses with multiple models (logistic regression analyses with several explanatory variables) with any in-championship injury (y/n) and in-championship time-loss injury (y/n) as outcomes. The explanatory variables were gender, age, home continent, event group (following event group categories from Timpka *et al*¹³) and preparticipation injury complaints. The multiple models were calculated using backward elimination of non-significant variables (ie, variables with $p \geq 0.05$ were eliminated stepwise). The Statistical Package for the Social Sciences (SPSS) for Windows V.21.0 was used for the analyses. All statistical tests were two-sided and outcomes with $p < 0.05$ regarded as statistically significant.

RESULTS

Study population

From a total population of 1784 registered athletes, 940 (52.7%) were reached to provide their informed consent to participate in the study; 750 athletes (42% of the total population) accepted. Fifty-two (2.9%) PHQs did not provide sufficient information to classify the athlete (eg, missing date of birth, event); 698 (39.1%) athletes were included in the study (figure 1). Analysis of the non-responders did not show any meaningful differences between the final study group and the non-participant athletes with regard to gender, age category and event groups; differences did exist for continents (slightly increased primary non-participation among Asian, North American and African athletes ($p < 0.001$); online supplementary table S1). The characteristics of the 698 athletes are reported in table 1. With regard to in-championship injury surveillance, 42 of the 102 national teams (41.2%) that had medical teams took part, covering 588 athletes (84.2%). The response rate for the injury report forms was 91%. None of the participating athletes refused to allow their injury data to be used for scientific research.

Figure 1 Flow chart of athlete enrolment and analysis population (PHQ, preparticipation health questionnaire).



Preparticipation injury complaints

Two hundred and four of the 698 athletes (29.2%) reported an injury complaint during the month before the championships; 108 reported having only one injury complaint, 61 reported two injury complaints or more, and 35 did not provide the total number of injury complaints. The characteristics of the main (worst) injury complaints are displayed in table 2, the distribution and incidence according to the event groups in table 3 and body location in table 4. One hundred and twenty-six athletes reporting injury complaints (61.8%) reported that the injury complaint problem did not affect their ability to participate in athletics (table 2). The most frequently reported mode of onset was gradual (49.2%), followed by sudden/overuse (36.5%). Only 9.9% of the preparticipation injury complaints were

caused by trauma. Almost half of the injury complaints (44.6%) had lasted for more than 4 weeks. There was a tendency for a higher proportion of gradual-onset injury complaints (61.7%) among those lasting for more than 4 weeks, and a higher proportion of sudden-onset injury complaints (50%) among those lasting for less than 2 weeks ($p=0.07$). At the start of the championship, 134 (65.6%) of the 204 athletes reported no or minor discomfort during training or competition caused by the injury complaint, 14 reported major discomfort and 3 were unable to participate.

In-championship injuries

Forty-nine (7%) athletes reported at least one in-championship injury. A total of 64 injuries were recorded, representing an incidence of 91.7 injuries per 1000 registered athletes (95% CI 70.3 to 113.1), including 35 time-loss injuries (54.7%). The most frequent location of injury was the thigh (33.3%), followed by the lower leg (20.6%). Hamstring strain (20.3% of injuries and 25.7% of time-loss injuries) was the most common diagnosis. The most commonly reported cause of injury was overuse (57.1%).

Associations between preparticipation risk factors and in-championship injuries

Using all in-championship injuries as the outcome (ie, any injury; y/n), the simple models showed that all categories for reporting an injury complaint in the month before the championship were associated with an increased injury risk: any injury complaint (OR=2.09; 95% CI 1.16 to 3.77; $p=0.014$), reduced participation injury complaints (OR=2.53; 95% CI 1.39 to 4.61; $p=0.002$), gradual-onset injury complaints (OR=2.72; 95% CI 1.38 to 5.36; $p=0.004$), and the injury complaint severity score (OR=1.14; 95% CI 1.06 to 1.22; $p=0.001$; table 5). In the analyses of the multiple models, only the injury complaint severity score remained associated with in-championship injury

Table 1 Characteristics of the 698 athletes who agreed to participate in the PHQ

	Total (n=698)	Females (n=325)	Males (n=373)
Athletes' characteristics, mean (SD)			
Age (years)	26.2 (4.3)	26.2 (4.2)	26.2 (4.4)
Height (cm)	176.7 (10.4)	170.1 (8.2)	182.4 (8.7)
Weight (kg)	68.2 (17.3)	59.3 (12.5)	75.9 (17.3)
Body mass index (kg/m ²)	21.6 (3.6)	20.4 (3.1)	22.5 (3.7)
Self-reported injury complaint in the 4 weeks before the championship, n (%)			
No	494 (70.8)	227 (69.8)	267 (71.6)
Yes	204 (29.2)	98 (30.2)	106 (28.4)

Among the 698 PHQs, 96.4% were complete with regard to the athlete's main characteristics (gender, age, country, height, weight, event group, time spent training; data were missing for females regarding height (n=15), weight (n=16), body mass index (n=20), training time (n=45), and for males regarding height (n=18), weight (n=15), body mass index (n=25), training time (n=66); preparticipation health status data were complete (question 1 of the PHQ). PHQ, preparticipation health questionnaire.

Table 2 Characteristics of preparticipation injury complaints (n=204)

	Total (n=204)	Females (n=98)	Males (n=106)
Level of athletics participation (training or competition) while experiencing an injury complaint, n (%)			
Full participation	126 (61.8)	66 (67.3)	60 (56.6)
Reduced participation (partial time loss)	70 (34.3)	30 (30.6)	40 (37.7)
No participation (full time loss)	8 (3.9)	2 (2.0)	6 (5.7)
Mode of onset, n (%)*			
Sudden onset (traumatic cause)	18 (9.9)	7 (8.0)	11 (11.8)
Sudden onset (overuse cause)	66 (36.5)	34 (38.6)	32 (34.4)
Gradual onset	89 (49.2)	45 (51.1)	44 (47.3)
Other incident, not related to training or competing in athletics	8 (4.4)	2 (2.3)	6 (6.5)
Injury complaint severity score, † mean (SD)	5.4 (3.7)	5.4 (3.6)	5.4 (3.7)
Duration, n (%)‡			
<1 week	27 (14.7)	12 (13.6)	15 (15.6)
1–2 weeks	43 (23.4)	22 (25.0)	21 (21.9)
3–4 weeks	32 (17.4)	16 (18.2)	16 (16.7)
>4 weeks	82 (44.6)	38 (43.2)	44 (45.8)

One preparticipation injury complaint was counted per athlete.

Among the 204 athletes who reported at least a pre-existing injury, the main athlete's characteristics (gender, age, country, height, weight, event group, time spent to training) were 96.4% complete and the injury complaint characteristics were 91.3% complete.

*Missing data for 23 injury complaints: 10 females and 13 males. These complaints were excluded from the calculations.

†The injury complaint severity score was calculated using items 3.1, 3.2, 3.3 and 3.4 of the PHQ inspired by the calculation of the OSTRC Overuse Injury Questionnaire severity score from Clarsen *et al.*⁹

‡Missing data for 20 injury complaints: 10 females and 10 males. These complaints were excluded from the calculations.

OSTRC, Oslo Sports Trauma Research Centre; PHQ, preparticipation health questionnaire.

(OR 1.14; 95% CI 1.06 to 1.22; $p=0.001$; table 5). Restricting the outcome for the examinations to in-championship time-loss injuries only (ie, time-loss injury; y/n), the analyses using simple models still showed that reporting injury complaints in the month before the championships was associated with increased injury risk, but only for gradual-onset complaints (OR=3.92; 95% CI, 1.69–9.08; $P=0.001$) and the injury complaint severity score (OR=1.12; 95% CI, 1.02–1.23; $P=0.001$) (table 5). In the analysis of the multiple models, only the gradual-onset injury complaints category remained with an almost fourfold increased injury risk (OR=3.92; 95% CI, 1.69–9.08; $P=0.001$) (table 5).

DISCUSSION

The main findings of this study were as follows: (1) about one-third of the athletes involved in this study and participating in an international athletics championship reported an injury complaint during the month before the competition; (2) almost two-thirds of these athletes (61.8%) stated that this injury complaint did not influence their participation; (3) athletes who reported a preparticipation injury complaint had twice the risk of an in-championship injury; (4) those who reported a preparticipation gradual-onset injury complaint had an almost fourfold increased risk of an in-championship time-loss injury; (5) the preparticipation injury complaint severity score was associated with an increased in-championship injury risk.

Competing with injury in athletics

About one-third of the athletes involved in this study and participating in this international championship reported at least

Table 3 Number of athletes reporting preparticipation injury complaints and number of athletes reporting in-championship injuries, displayed by event groups and severity (n=698); in the case of several preparticipation injury complaints, only the main (worst) complaint is presented and in the case of several in-championship injuries only the first is presented

	Sprints	Hurdles	Middle distance	Long distance	Marathon	Race walking	Jumps	Throws	Combined events	Total
Registered athletes, n	190	65	52	67	57	75	94	75	23	698
Preparticipation injury complaint										
Athletes reporting preparticipation injury complaint, n (%)	57 (30.0)	25 (38.5)	14 (26.9)	15 (22.4)	11 (19.3)	20 (26.7)	33 (35.1)	21 (28.0)	8 (34.8)	204 (100)
Athletes reporting preparticipation injury complaint per 1000 athletes	300.0	384.6	269.2	223.9	193.0	266.7	351.1	280.0	347.8	292.3
Athletes with reduced preparticipation injury complaint, n (%)	24 (12.6)	10 (15.4)	6 (11.5)	4 (6.0)	3 (5.3)	8 (10.7)	11 (11.7)	11 (14.7)	1 (4.3)	78 (100)
Athletes with reduced preparticipation injury complaint per 1000 athletes	126.3	153.8	115.4	59.7	52.6	106.7	117.0	146.7	43.5	111.7
Injury complaint severity score for athlete category, mean (SD)	1.7 (3.1)	2.6 (4.2)	1.5 (3.1)	0.9 (2.2)	0.7 (2.2)	1.2 (2.7)	2.0 (3.5)	1.5 (3.3)	1.3 (2.6)	1.6 (3.1)
In-championship injury										
Athletes with in-championship injury, n (%)	18 (9.5)	7 (10.8)	1 (1.9)	4 (6.0)	4 (7.0)	3 (4.0)	5 (5.3)	3 (4.0)	4 (17.4)	49 (100)
Athletes with in-championship injury per 1000 athletes	94.7	107.7	19.2	59.7	70.2	40.0	53.2	40.0	173.9	70.2
Athletes with in-championship time-loss injury, n (%)	9 (4.7)	5 (7.7)	1 (1.9)	2 (3.0)	3 (5.3)	1 (1.3)	2 (2.1)	2 (2.7)	2 (8.7)	27 (100)
Athletes with in-championship time-loss injury per 1000 athletes	47.4	76.9	19.2	29.9	52.6	13.3	21.3	26.7	87.0	38.7

Table 4 Body location for the main (worst) preparticipation injury complaint and body location for the first in-championship injury

Body location	Preparticipation injury complaints			In-championship injuries		
	Total, n (%)	Female, n (%)	Male, n (%)	Total, n (%)	Female, n (%)	Male, n (%)
Total	194	92	102	48	18	30
Face/head	10 (5.1)	3 (3.2)	7 (6.8)	0 (0.0)	0 (0.0)	0 (0.0)
Neck/spine	23 (11.9)	7 (7.6)	16 (15.7)	1 (2.1)	1 (5.6)	0 (0.0)
Trunk (except spine)	18 (9.2)	10 (10.9)	8 (7.8)	2 (4.2)	1 (5.6)	1 (3.3)
Shoulder	3 (1.5)	2 (2.2)	1 (1.0)	1 (2.1)	0 (0.0)	1 (3.3)
Arm/hand except shoulder	4 (2.0)	3 (3.3)	1 (1.0)	3 (6.3)	2 (11.1)	1 (3.3)
Hip/groin	21 (10.8)	10 (10.9)	11 (10.8)	0 (0.0)	0 (0.0)	0 (0.0)
Thigh	32 (16.5)	15 (16.3)	17 (16.7)	17 (35.4)	6 (33.3)	11 (36.7)
Knee	24 (12.4)	10 (10.9)	14 (13.7)	3 (6.3)	0 (0.0)	3 (10.0)
Lower leg	24 (12.4)	14 (15.2)	10 (9.8)	9 (18.8)	3 (16.7)	6 (20.0)
Achilles tendon	0 (0.0)	0 (0.0)	0 (0.0)	4 (8.3)	1 (5.6)	3 (10.0)
Ankle	19 (9.8)	10 (10.9)	9 (8.8)	1 (2.1)	0 (0.0)	1 (3.3)
Foot	16 (8.4)	8 (8.7)	8 (7.8)	7 (14.6)	4 (22.2)	3 (10.0)

One preparticipation injury complaint and one in-championship injury was counted per athlete. Information was missing for 10 preparticipation injury complaints (6 females and 4 males) and for 1 in-championship injury (1 male). These complaints were excluded from the calculations. The mean severity scores (SD) were 6.0 (3.5) for preparticipation thigh complaints, 4.9 (4.3) for preparticipation knee complaints, 4.7 (3.4) for preparticipation lower leg complaints, 5.2 (3.9) for preparticipation neck/spine complaints, 5.9 (4.6) for preparticipation hip/groin complaints and 5.9 (3.0) for preparticipation ankle complaints.

one injury complaint during the previous month. This finding extends results from our pilot study at the 2013 European Athletics Indoor Championships during which 54% of the athletes included reported at least one health problem (injury

complaint for 24%, illness complaint for 18% and pain complaint for 30%) during the 4 weeks leading up to the championships.⁶ This is a relatively high percentage given that athletes aim to peak at a championship. The results also correspond to

Table 5 Risk factors for sustaining an in-championship injury or an in-championship time-loss injury presented as ORs (95% CI) calculated by simple and multiple logistic regression analyses (n=698)

	OR (95% CI)			
	In-championship injury		In-championship time-loss injury	
	Simple models	Multiple model	Simple models	Multiple model
Gender	NS	NS	NS	NS
Age	NS	NS	NS	NS
Continent (reference continent was Europe)	NS	NS	NS	NS
Africa	NS	NS	NS	NS
Asia	NS	NS	NS	NS
Australia	NS	NS	NS	NS
North America	NS	NS	NS	NS
South America	NS	NS	NS	NS
Event group (reference group was combined events)	NS	NS	NS	NS
Sprints	NS	NS	NS	NS
Hurdles	NS	NS	NS	NS
Middle distance	NS	NS	NS	NS
Long distance	NS	NS	NS	NS
Marathon	NS	NS	NS	NS
Race walking	NS	NS	NS	NS
Jumps	NS	NS	NS	NS
Throws	NS	NS	NS	NS
Preparticipation injury complaints				
All complaints	2.09 (1.16 to 3.77) (p=0.014)	NS	NS*	NS
Reduced participation injury complaints	2.53 (1.39 to 4.61) (p=0.002)	NS	NS†	NS
Gradual-onset injury complaints	2.72 (1.38 to 5.36) (p=0.004)	NS	3.92 (1.69 to 9.08) (p=0.001)	3.92 (1.69 to 9.08) (p=0.001)
Injury complaint severity score (maximum score 16)	1.14 (1.06 to 1.22) (p<0.001)	1.14 (1.06 to 1.22) (p<0.001)	1.12 (1.02 to 1.23) (p=0.019)	NS

*OR 2.14 (95% CI 0.97 to 4.72; p=0.058).

†OR 2.19 (95% CI 0.98 to 4.94; p=0.058).

NS, not significant.

the findings from a study performed at the 2013 FINA World Championships (Aquatics) where 33% of responding athletes reported a physical complaint in the 4 weeks prior to the championships,¹⁴ and are also in agreement with those from a cohort of Swedish national-level gymnasts, where 58% competed with injury complaints.¹⁵

For the sport of athletics, it is a matter of concern that a high number of athletes seem to live and train with injuries. About half of the athletes reported a preparticipation injury complaint related to overuse and more than half had had their injury complaint for more than 3 weeks. These data are consistent with historical publications where the incidence and prevalence of injuries throughout the athletics season is about 2–3 injuries per athlete per year.^{16–20}

Jacobsson *et al*¹⁹ found that 96% of injuries reported by elite Swedish athletes throughout the season were classified as non-traumatic (caused by overuse). Similarly, in a cohort of 142 Olympic and Paralympic candidates from different sports, Clarsen *et al*¹⁰ reported an average weekly prevalence of overuse injury of 25% at any given point in the season. These results suggest that athletes continue training and compete even though they have injury and/or health problems. To confirm this, prospective whole-season studies must be conducted in athletics to record injuries, including overuse injuries, and examine possible causes and their functional consequences.^{13 19 21}

Bringing injuries to the championships increases risk of injury

Having an injury complaint during the month preceding an international athletics championship represented a risk factor for sustaining an in-championship injury. Our results showed that the injury complaint severity score predicted sustaining any injury during the championships according to the multiple model (logistic regression analyses with several explanatory variables), suggesting that having an injury complaint before the championships interferes with preparations and predisposes athletes to an injury during the championships. Injury complaints during the past month may have delayed the athletes' physical preparation and training and affected their ability to perform optimally at the championships, leaving the athlete more at risk of injury. Therefore, the injury severity score could be a good method for predicting which athletes are at increased risk of sustaining an injury during the championships, but not the severity of that injury.

A gradual-onset injury complaint was strongly associated with the risk for sustaining a time-loss injury at the championships. These measures of preparticipation injury complaints thus seem to have the highest validity with regard to impact and relevance in athletics. These results support the further development of functional consequences assessment of injuries to better define the severity in addition to the duration of time loss.^{9 22} The duration of time loss may not be an appropriate measure of the severity of an injury for athletes,^{20 22} because many athletes continue to train and compete despite being injured.^{9 10 15 22} The severity score (or functional consequences assessment) may help to better define the magnitude of the functional consequences for athletes. However, more research is needed to identify a cutpoint whereby the severity score denotes a clear injury risk.

Limitations

Several potential limitations should be acknowledged when interpreting the results of this study. First, the percentage of athletes included in the study is low. However, the 60.9% of all

registered athletes who did not participate in this study would most probably not change the results, since the study group of 698 athletes is similar to the non-participating athletes on the demographic and in-championship injury characteristics. Considering the methods of preparticipation data collection (circumstances and timing correspond to the last 3 days before and throughout the IAAF World championships), the response rate of 39.1% to our preparticipation questionnaire is acceptable. However, we need to consider how modern technology including tablets and translation mobile applications could help us improve our recruitment of athletes.²³

We faced methodological challenges in reaching out to more than 200 national teams with native languages not covered in our translations of the information and questionnaire. There was also lack of trust in our confidentiality measures which could in part explain the proportion of non-responders. We noted a lower response rate among North American, African and Asian athletes, which indicates that language, social or other cultural disparities were involved to some extent.

Furthermore, some athletes and team medical personnel voiced concerns about filling out forms about personal injuries at a time close to the most important competition of the year. This is hard to circumvent when collecting data during a championship.

Our response rate was lower than that of a previous pilot study (60.6%),⁶ but the pilot study was a European indoor championship with a smaller and more homogeneous study population with less language and cultural differences. For future studies, methodological factors must be mastered in order to increase the response rate (improving the information about the study and its interest for athletes, increasing the number of languages, developing informatics support rather than the paper form).

Second, this study combined two different methods for collecting injury data: athletes' retrospective self-reports and medical attention at the championship. Although a detailed definition of injury complaint was given at the start of the PHQ, athletes could have misunderstood this.⁹ Furthermore, self-reported data might provide a limited description of injuries; data on the exact location (including type, severity and cause) and functional consequences should preferably be reported by medical practitioners. In addition, injuries reported at the championships by national and LOC medical teams might not all be new index injuries. On review of the data (not presented in this study), we noted that athletes reported an injury complaint in the PHQ and then the same problem was reported by the medical team as a new injury incurred during the championship.

This observation does not affect the results in this study but indicates an area where the method of reporting needs to be improved to distinguish between, for example, recurrence and exacerbation, because gradual-onset injury complaints were the best predictor of injury. Also, the complete burden of injuries affecting the athletes is not taken into account fully.

Third, the retrospective design of the PHQ could represent a recall bias, although this should have been minimal as we asked the participants only about the 4 weeks immediately before the competition.⁶ The content and characteristics of the PHQ should be analysed further and validated in a future study.

SUMMARY AND CONCLUSIONS

A self-reported PHQ was feasible to determine the preparticipation health status of athletes before an international championship. Moreover, these data can help to determine preparticipation risk

factors and help identify which athletes may be at increased risk of injuries. About one-third of athletes participating in the study reported an injury complaint during the month before the championships. These athletes were at twice the risk of an in-championship injury. In addition, athletes who reported a pre-participation gradual-onset injury complaint were at an almost fourfold increased risk for an in-championship time-loss injury. The preparticipation injury complaint severity score was associated with a general increased risk for injury.

What is already known on this topic?

- ▶ During previous international outdoor athletics championships, injury incidence has been about 100 injuries per 1000 athletes.
- ▶ The most frequently reported injury has been thigh strain, especially hamstring strain, and overuse has been the predominant cause.
- ▶ To better understand what causes injuries during a championship, the European Athletics Association and the International Association of Athletics Federations encourage teams to administer a valid athlete self-reporting preparticipation health questionnaire.

What are the new findings?

- ▶ During the month before the championships, about one-third of athletes involved in this study were troubled by an injury complaint.
- ▶ Athletes who presented with a preparticipation injury complaint were at twice the risk of sustaining an injury and/or time-loss injury during the championships.
- ▶ Athletes who reported a preparticipation gradual-onset injury complaint were at an almost fourfold increased risk for an in-championship time-loss injury.

How might it impact clinical practice in the near future?

- ▶ Up-to-date information on the health status of athletes is important.
- ▶ National medical teams should obtain health status data during the month before and at the start of the championships.
- ▶ We recommend a routine preparticipation health questionnaire for athletes and medical teams before the championships in order to screen for athletes at risk of being injured.

Author affiliations

¹Medical and Anti-doping Commission, International Association of Athletics Federations (IAAF), Monaco, Monaco

²Sports Medicine Department, Aspetar, Qatar Orthopedics and Sports Medicine Hospital, Doha, Qatar

³Athletics Research Center, Linköping University, Linköping, Sweden

⁴Swedish Athletics Association, Stockholm, Sweden

⁵Aker Solutions, Lysaker, Norway

⁶Institute of Sport, Lithuanian University of Health Sciences, Kaunas, Lithuania

⁷Department of Behavioural Sciences and Learning, Linköping University, Linköping, Sweden

⁸Department of Clinical and Exercise Physiology, Sports Medicine Unity, Faculty of Medicine, University Hospital of Saint-Etienne, Saint-Etienne, France

⁹Laboratory of Exercise Physiology (LPE EA 4338), University of Lyon, Saint-Etienne, France

¹⁰Medical Commission, French Athletics Federation (FFA), Paris, France

Twitter Follow Juan Manuel Alonso at @DrJuanMalonso

Acknowledgements The authors highly appreciate the cooperation of the team physicians and physiotherapists and the local organising committee of the 2013 IAAF World Championships who volunteered their time to collect the data for this project. The authors would also like to thank Jan Kowalski for his help with the data support, Ben Clarsen, Astrid Junge, Margo Mountjoy and Kathrin Steffen for their help with the development of the preparticipation health questionnaire, and Mathieu Oriol and Vincent Pichot for their contribution to the statistical analyses.

Contributors J-MA and PE made substantial contributions to the development of the project. J-MA, JJ, OR and PE made substantial contributions to the conception and design of the study. J-MA, JJ, OR, AK and PE were involved in data collection. JJ, TT, OD, AS and PE were involved in the analysis of the data. J-MA, JJ, TT, OR and PE made substantial contributions to data interpretation. PE made a substantial contribution to the drafting and writing of the article. All authors were involved in revising the manuscript and final approval of the version to be published.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Feddermann-Demont N, Junge A, Edouard P, *et al.* Injuries in 13 international athletics championships between 2007–2012. *Br J Sports Med* 2014;48:513–22.
- 2 Alonso JM, Junge A, Renstrom P, *et al.* Sports injuries surveillance during the 2007 IAAF World Athletics Championships. *Clin J Sport Med* 2009;19:26–32.
- 3 Alonso JM, Tscholl PM, Engebretsen L, *et al.* Occurrence of injuries and illnesses during the 2009 IAAF World Athletics Championships. *Br J Sports Med* 2010;44:1100–5.
- 4 Alonso JM, Edouard P, Fischetto G, *et al.* Determination of future prevention strategies in elite track and field: analysis of Daegu 2011 IAAF Championships injuries and illnesses surveillance. *Br J Sports Med* 2012;46:505–14.
- 5 Edouard P, Depiessis F, Branco P, *et al.* Analyses of Helsinki 2012 European Athletics Championships injury and illness surveillance to discuss elite athletes risk factors. *Clin J Sport Med* 2014;24:409–15.
- 6 Edouard P, Jacobsson J, Timpka T, *et al.* Extending in-competition athletics injury and illness surveillance with pre-participation risk factor screening: a pilot study. *Phys Ther Sport* Published online: 14 Jun 2014. doi:10.1016/j.ptsp.2014.05.003
- 7 Clarsen B, Bahr R. Matching the choice of injury/illness definition to study setting, purpose and design: one size does not fit all! *Br J Sports Med* 2014;48:510–12.
- 8 Timpka T, Jacobsson J, Ekberg J, *et al.* Meta-narrative analysis of sports injury reporting practices based on the Injury Definitions Concept Framework (IDCF): a review of consensus statements and epidemiological studies in athletics (track and field). *J Sci Med Sport* Published online: 5 Dec 2014. doi:10.1016/j.jsams.2014.11.393
- 9 Clarsen B, Myklebust G, Bahr R. Development and validation of a new method for the registration of overuse injuries in sports injury epidemiology: the Oslo Sports Trauma Research Centre (OSTRC) Overuse Injury Questionnaire. *Br J Sports Med* 2013;47:495–502.
- 10 Clarsen B, Ronsén O, Myklebust G, *et al.* The Oslo Sports Trauma Research Center questionnaire on health problems: a new approach to prospective monitoring of illness and injury in elite athletes. *Br J Sports Med* 2014;48:754–60.
- 11 Edouard P, Depiessis F, Hertert P, *et al.* Injuries and illnesses during the 2011 Paris European Athletics Indoor Championships. *Scand J Med Sci Sports* 2013;23:e213–18.
- 12 Junge A, Engebretsen L, Alonso JM, *et al.* Injury surveillance in multi-sport events: the International Olympic Committee approach. *Br J Sports Med* 2008;42:413–21.
- 13 Timpka T, Alonso JM, Jacobsson J, *et al.* Injury and illness definitions and data collection procedures for use in epidemiological studies in Athletics (track and field): consensus statement. *Br J Sports Med* 2014;48:483–90.
- 14 Mountjoy M, Junge A, Benjamin S, *et al.* Competing with injuries: injuries prior to and during the 15th FINA World Championships 2013 (aquatics). *Br J Sports Med* 2015;49:37–43.
- 15 Harringe ML, Lindblad S, Werner S. Do team gymnasts compete in spite of symptoms from an injury? *Br J Sports Med* 2004;38:398–401.
- 16 D'Souza D. Track and field athletics injuries—a one-year survey. *Br J Sports Med* 1994;28:197–202.

- 17 Bennell KL, Crossley K. Musculoskeletal injuries in track and field: incidence, distribution and risk factors. *Aust J Sci Med Sport* 1996;28:69–75.
- 18 Rebella GS, Edwards JO, Greene JJ, et al. A prospective study of injury patterns in high school pole vaulters. *Am J Sports Med* 2008;36:913–20.
- 19 Jacobsson J, Timpka T, Kowalski J, et al. Injury patterns in Swedish elite athletics: annual incidence, injury types and risk factors. *Br J Sports Med* 2013;47:941–52.
- 20 Jacobsson J, Timpka T, Kowalski J, et al. Prevalence of musculoskeletal injuries in Swedish elite track and field athletes. *Am J Sports Med* 2012;40:163–9.
- 21 Edouard P, Branco P, Alonso JM. Challenges in athletics injury and illness prevention: implementing prospective studies by standardised surveillance. *Br J Sports Med* 2014;48:481–2.
- 22 Bahr R. No injuries, but plenty of pain? On the methodology for recording overuse symptoms in sports. *Br J Sports Med* 2009;43:966–72.
- 23 Verhagen EA, Clarsen B, Bahr R. A peek into the future of sports medicine: the digital revolution has entered our pitch. *Br J Sports Med* 2014;48:739–40.