



OPEN ACCESS

Hamstring injuries: prevention and treatment— an update

Peter Brukner

Correspondence to

Dr Peter Brukner, La Trobe University, Melbourne, Victoria, Australia; peterbrukner@gmail.com

Accepted 31 May 2015
Published Online First
23 June 2015

ABSTRACT

Despite increased knowledge of hamstring muscle injuries, the incidence has not diminished. We now know that not all hamstring injuries are the same and that certain types of injuries require prolonged rehabilitation and return to play. The slow stretch type of injury and injuries involving the central tendon both require longer times to return to play. A number of factors have been proposed as being indicators of time taken to return to play, but the evidence for these is conflicting. Recurrence rates remain high and it is now thought that strength deficits may be an important factor. Strengthening exercise should be performed with the hamstrings in a lengthened position. There is conflicting evidence regarding the efficacy of platelet-rich plasma injection in the treatment of hamstring injuries so at this stage we cannot advise their use. Various tests have been proposed as predictors of hamstring injury and the use of the Nordboard is an interesting addition to the testing process. Prevention of these injuries is the ultimate aim and there is increasing evidence that Nordic hamstring exercises are effective in reducing the incidence.

INTRODUCTION

In spite of all the research and additional understanding of hamstring muscle injuries over the past 20–30 years, we have not reduced the incidence of first-time injuries and the recurrence rate is still extremely high. While research published over the past couple of years has led to an increased understanding of these challenging injuries, we still have a long way to go in the management of hamstring muscle injuries.

NOT ALL HAMSTRING INJURIES ARE THE SAME

We now understand that certain types of hamstring injuries are more likely to require prolonged rehabilitation and delayed return to play (RTP). Askling *et al*^{1–3} has proposed two distinctly different types of acute hamstring strains, one occurring during high-speed running and mainly involving the biceps femoris long head, the other during movements leading to extensive lengthening of the hamstrings (such as high kicking, sliding tackle, sagittal split) often involving the free proximal tendon of semimembranosus. He demonstrated that injuries of the slow stretch type, while initially appearing less severe than the sprinting type of injury, actually have a prolonged RTP.^{1–4}

In addition, muscle tendon architecture may be a factor in the development of hamstring injuries. Rehorn and Blemker,⁵ using a three-dimensional model, initially suggested that the aponeurosis morphology of the biceps femoris long head

(BFLh) may play a significant role in determining stretch distributions throughout the muscle. Others have suggested that variability of aponeurosis widths may be important in determining muscle injury susceptibility.⁶ Studies have suggested that a relatively small^{7,8} or narrow⁹ proximal aponeurosis of the BFLh may be predictive of hamstring injury.

This may explain why injuries involving the central tendon have been shown to be associated with prolonged RTP.¹⁰ MRI has enabled us to visualise the intramuscular component of the tendon.

A number of us have for many years promoted the concept that the mild hamstring injury, often presenting as a ‘cramp’ or ‘twinge’ or ‘feeling of impending tearing’, is not due to a hamstring muscle injury, but due to referred pain from the lumbar spine, fascial injury or gluteal trigger points. Now with the advent of MRI showing minimal or no local muscle damage in these cases, the so-called MRI-negative hamstring injury, there has been broader acceptance of this phenomenon and appropriate treatment initiated with resulting early RTP.^{11,12}

WHAT DETERMINES RTP?

There are a number of factors that have been suggested as good indicators of severity and prolonged time to return to play. Askling *et al*³ suggested that the closer the lesion was to the ischial tuberosity the longer the time to RTP. While a relationship between proximity to the ischial tuberosity and time to RTP makes sense based on anatomical knowledge (ie, likelihood to involve tendon), it is worth noting that a recent review¹² found some conflicting evidence. Of the four studies performed to date, three studies reported a significant association between a shorter distance to the ischial tuberosity and a longer time to RTP,^{3,13,14} whereas one study found no association.²

The same review¹² also evaluated the relationship between the size of the hamstring injury lesion on MRI, including the length, cross-sectional area and signal volume, and the time to RTP. The review concluded that there was no strong evidence from any MRI finding that can guide radiologists and sports physicians in predicting prognosis for the time to RTP after an acute hamstring injury. Similarly there was no correlation of ultrasound findings with time to RTP.¹⁵

In a study comparing clinical and MRI indicators of RTP,¹⁶ the clinical parameters of self-predicted time to RTP (TTRTP) and passive straight leg raise deficit were independently associated with the TTRTP. The latter contradicted the findings of two previous studies.^{4,17} MRI parameters in grade 1 and 2 hamstring injuries were not associated with TTRTP. It was, therefore, suggested that for clinical



Open Access
Scan to access more
free content



CrossMark

To cite: Brukner P. *Br J Sports Med* 2015;**49**:1241–1244.

practice, prognosis of the TTRTP in these injuries should be based on key clinical parameters.

RECURRENCE

Reinjury after RTP remains a major problem. It is more common when the injury involves the biceps femoris.¹¹ The number of previous hamstring injuries, active knee extension deficit, isometric knee flexion force deficit at 15°, and presence of localised discomfort on palpation just after RTP are also associated with a higher hamstring reinjury rate.¹⁸

There is increasing evidence that even after RTP, eccentric hamstring strength is reduced, which may be a factor in the high recurrence rate of these injuries.^{19–25} Earlier studies failed to show any differences.^{26 27}

Whereas studies by Silder *et al*²⁸ and Emami *et al*²⁹ have failed to show any difference, Opar *et al*³⁰ demonstrated that previously injured hamstrings displayed lower rate of torque development (RTD) and early contractile impulse (IMP) during slow maximal eccentric contraction compared with the contralateral uninjured limb. Lower myoelectrical activity was confined to the biceps femoris long head.

Regardless of whether these deficits are the cause of or the result of injury, these findings could have important implications for hamstring strain injury and reinjury. Particularly, given the importance of high levels of muscle activity to bring about specific muscular adaptations, lower levels of myoelectrical activity may limit the adaptive response to rehabilitation interventions and suggest that greater attention be given to neural function of the knee flexors after hamstring strain injury.

It has been suggested that the cause of this eccentric weakness is prolonged neuromuscular inhibition at long muscle lengths after hamstring muscle injury.³¹ Pain-driven neuromuscular inhibition of hamstring voluntary activation occurs following hamstring strain injury, and this inhibition has a detrimental effect on hamstring recovery by limiting hamstring exposure to eccentric stimuli at long muscle lengths during rehabilitative exercise.³¹

Mendiguchia *et al*³² examined the effects of an acute hamstring strain injury on sprinting performance, and mechanical properties of sprint running at the time of return to sport and 2 months later. The study showed that despite being cleared to play, soccer players returning from a recent hamstring injury had substantial lower sprinting speed performance and reduced mechanical horizontal properties compared to the uninjured players. The greater magnitude differences in horizontal force compared to maximum velocity suggested that the lower maximal horizontal power observed in the injured player was mainly related to the reduced maximal horizontal force component. Approximately 2 months of regular soccer training after return to sports resulted in substantial improvements in sprinting speed (acceleration) concomitant with an increase in maximal horizontal force and power, whereas the speed component and top speed remained unaltered.

The limited exposure to eccentric stimuli could potentially produce several maladaptations observed following hamstring injury, including chronic eccentric hamstring weakness,^{20 31 33 34} selective hamstring atrophy³⁵ and shifts in the torque joint-angle relationship.²⁶ Timmins *et al*^{36 37} recently demonstrated shorter biceps fascicle length and increased pennation angle as well as reduced eccentric strength in previously injured hamstrings.

IMPLICATIONS FOR CLINICAL MANAGEMENT

What are the implications of this research for clinical practice? Perhaps more emphasis should be placed on reduction of pain

in the early days after hamstring injury to reduce the neuromuscular inhibition associated with pain, while at the same time encouraging early muscle activation, particularly eccentric exercise at longer muscle lengths, and early return to running with rapid progression to high-speed running.

While the concept of eccentric muscle training as an important component of the rehabilitation process has been with us for many years, it now appears that these exercises must be in the lengthened position. This makes sense when you think that the majority of hamstring muscle injuries are located in the long head of biceps femoris, a muscle that straddles both the hip and knee joints. The standard leg curl exercise, therefore, does not work the long head sufficiently. As a result, lengthening eccentric exercises such as the Nordic hamstring exercise (NHE),^{38 39} the Romanian dead lift and Askling's 'extender', 'diver' and 'glider' exercises¹³ are now becoming the mainstay of postinjury rehabilitation.

Two papers have proposed parameters by which hamstring rehabilitation programmes can be created. Malliaropoulos *et al*⁴⁰ suggested the following parameters—*injury mechanism, hip or knee dominant, location, targeted muscle, length rather than strength, training parameters*—should be considered when devising a rehabilitation protocol.

Guex and Millet⁴¹ suggested a conceptual framework for strengthening exercises for hamstring muscles specific to the terminal swing phase of sprinting based on six key parameters (contraction type, load, range of motion, angular velocity, unilateral/bilateral exercises, kinetic chain) for strain prevention. They advocated that in sprinting sports, high-load eccentric contractions should be performed at a slow to moderate angular velocity and focused at the knee joint, while the hip is kept in a large flexion position (80°) in order to expose the hamstrings to a greater elongation stress than occurs in the terminal swing phase. They postulated that as a result, during sprinting, athletes would be better trained to brake knee extension effectively in the whole range of motion without overstretch of the hamstrings.

They also advocated unilateral open kinetic chain exercises based on their functional application. After analysing some of the frequently used hamstring strengthening exercises, they came to the conclusion that the 'optimal exercise had not been designed yet'. Finally, they noted that strain prevention is not only a question of strength, but also depends on the timing of contraction, or a combination of both.

REHABILITATION PROGRAMMES

For a number of years the only RCT comparing different hamstring rehabilitation programmes was Sherry and Best⁴² study, which reported significantly lower reinjury rates in athletes who completed a progressive agility and trunk stabilisation (PATS) programme, compared to those whose rehabilitation programmes focused on isolated hamstring strengthening and stretching.

Silder *et al*¹⁴ demonstrated a similar degree of muscle recovery at the time of return to sport in patients with an acute hamstring strain injury treated with either the PATS programme or a programme with a heavy emphasis on eccentric strengthening (PRES).

Askling *et al*¹³ performed two identical studies, one in footballers and other in sprinters and jumpers,⁴³ and demonstrated that a rehabilitation protocol consisting of mainly lengthening type of exercises (L-protocol) is more effective than a conventional protocol in promoting return to sport after acute hamstring injury. The most conspicuous characteristics of the more successful L-protocol were the systematic attempts to put load on the hamstrings during maximal dynamic lengthening, using

exercises entitled The Extender, The Diver and The Glider. On this basis, they recommended that hamstring injury rehabilitation protocols should be preferentially based on strength and flexibility exercises that primarily involve exercises with high loads at long muscle–tendon lengths.

IS THERE A ROLE FOR PLATELET-RICH PLASMA?

The use of autologous blood injections, such as platelet-rich plasma (PRP) has become widespread in recent years, primarily in the treatment of chondral and tendon problems. Recently the first two high-quality studies have been published examining the use of PRP in muscle injuries. Two studies produced conflicting results.

Using a double-blind, randomised and multicentre approach, Reurink *et al*⁴⁴ recruited 80 recreational athletes with hamstring muscle strain injuries, and infiltrated the injured area with either isotonic saline or a standardised formulation of PRP. Ultimately, the authors found no statistical or clinically significant difference in either of their key outcome measures, RTP duration and reinjury rate, leading them to conclude that PRP is no more effective than a placebo injection of saline.

Hamid *et al*⁴⁵ compared a group who received a PRP injection along with a standard rehabilitation programme with a control group who received the rehabilitation programme alone and found that patients in the PRP group achieved full recovery significantly earlier than controls. Significantly, lower pain severity scores were observed in the PRP group throughout the study. They concluded that a single autologous PRP injection combined with a rehabilitation programme was significantly more effective in treating hamstring injuries than a rehabilitation programme alone.

Of the two studies, the Reurink paper is of substantially higher quality with its use of a placebo saline injection. Previous PRP studies using saline as placebo^{46–47} have also failed to confirm the improvements found in those studies without placebo. There is not at this stage sufficient evidence to advocate the use of PRP in acute muscle injuries.

CAN WE PREDICT HAMSTRING INJURY?

Previous attempts to develop a tool to predict the likelihood of hamstring injuries have been based on isokinetic testing.⁴⁸ More recently, other tests have been proposed.

Freckleton *et al*⁴⁹ demonstrated a significant deficit in pre-season single leg hamstring bridge (SLHB) scores on the right leg of players who subsequently sustained a right-sided hamstring injury. Age, previous knee injury and a history of hamstring injury were other risk factors supported in this study.

Shield and Opar have designed a test, the Nordic board test, to measure hamstring strength based on the NHE. Their study⁵⁰ demonstrated that (1) the experimental device showed high to moderate test–retest reliability for measurements when the NHE was performed bilaterally, but poor reliability during unilateral testing, and (2) elite athletes with a unilateral history of hamstring injury within the previous 12 months displayed significant eccentric knee flexor weakness in their injured limb compared to their uninjured limb and to the limbs of uninjured recreational athletes.

Hamstring weakness in Australian rules footballers demonstrated on the Nordic board test was associated with increased risk of hamstring injury. Higher Nordic hamstring strength offset the risk of increasing age and previous hamstring injury.⁵¹

Another study⁵² examined hamstring strength measured with a hand-held dynamometer and distance achieved in a single leg hop test. They found that lower maximum eccentric hamstring

strength, higher isometric/eccentric hamstring strength ratio, and a lower score on the Sydney Local Health District (SLHD) test were significant risk factors for a subsequent hamstring injury.

Finally, in a case report, Schache *et al*⁵³ looked at the use of regular clinical monitoring of hamstring strength during a season in football players with a history of hamstring strain. It was concluded that measuring postgame hamstring isometric maximal voluntary contraction asymmetry on a weekly basis may be helpful in identifying adverse reactions to load (ie, inhibition, presence of symptoms, or both) that could represent early warning signs for hamstring strain susceptibility.

It is clear that we are some distance away from being able to predict hamstring muscle injury. Many of the issues highlighted by Mendiguchia in his 2012 *BJSM* paper⁵⁴ remain unresolved. The inter-relationships between the various possible risk factors need to be examined more fully.

CAN WE PREVENT HAMSTRING INJURIES?

Prevention is better than cure and there is evidence that a programme of eccentric hamstring exercises, such as the yo-yo curl⁵⁵ or NHE,^{38–56–59} can reduce the incidence of hamstring muscle injuries.

The recommended NHE programme³⁸ of three sessions per week during a 10-week pre-season programme and subsequently, one session a week has been incorporated in the training regimes of many football clubs. An Italian study⁶⁰ demonstrated reduced injuries in an amateur football club using the FIFA 11+ injury prevention programme with addition of NHE.

CONCLUSION

While there has been significant additions to the literature over the past couple of years, we have still not managed to reduce the incidence of hamstring muscle injuries. Further high-quality research is needed.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Asklings CM, Malliaropoulos N, Karlsson J. High-speed running type or stretching-type of hamstring injuries makes a difference to treatment and prognosis. *Br J Sports Med* 2012;46:86–7.
- 2 Asklings CM, Tengvar M, Saartok T, *et al*. Acute first-time hamstring strains during slow-speed stretching: clinical, magnetic resonance imaging, and recovery characteristics. *Am J Sports Med* 2007;35:1716–24.
- 3 Asklings CM, Tengvar M, Saartok T, *et al*. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med* 2007;35:197–206.
- 4 Asklings C, Saartok T, Thorstensson A. Type of acute hamstring strain affects flexibility, strength, and time to return to pre-injury level. *Br J Sports Med* 2006;40:40–4.
- 5 Rehorn MR, Blemker SS. The effects of aponeurosis geometry on strain injury susceptibility explored with a 3D muscle model. *J Biomech* 2010;43:2574–81.
- 6 Fiorentino NM, Blemker SS. Musculotendon variability influences tissue strains experienced by the biceps femoris long head muscle during high-speed running. *J Biomech* 2014;47:3325–33.
- 7 Evangelidis P, Massey G, Pain M, *et al*. Is biceps femoris long head aponeurosis size a risk factor for hamstring strain injury? *Br J Sports Med* 2014;48:591–2.
- 8 Evangelidis PE, Massey GJ, Pain MTG, *et al*. Biceps femoris aponeurosis size: a potential risk factor for strain injury? *Med Sci Sports Exerc* 2014. [Epub ahead of print].

- 9 Fiorentino NM, Epstein FH, Blemker SS. Activation and aponeurosis morphology affect in vivo muscle tissue strains near the myotendinous junction. *J Biomech* 2012;45:647–52.
- 10 Comin J, Malliaras P, Baquie P, et al. Return to competitive play after hamstring injuries involving disruption of the central tendon. *Am J Sports Med* 2013;41:111–15.
- 11 Hallen A, Ekstrand J. Return to play following muscle injuries in professional footballers. *J Sports Sci* 2014;32:1229–36.
- 12 Reurink G, Brilman EG, de Vos RJ, et al. Magnetic resonance imaging in acute hamstring injury: can we provide a return to play prognosis? *Sports Med* 2015;45:133–46.
- 13 Askling CM, Tengvar M, Thorstensson A. Acute hamstring injuries in Swedish elite football: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med* 2013;47:953–9.
- 14 Silder A, Sherry MA, Sanfilippo J, et al. Clinical and morphological changes following 2 rehabilitation programs for acute hamstring strain injuries: a randomized clinical trial. *J Orthop Sports Phys Ther* 2013;43:284–99.
- 15 Petersen J, Thorborg K, Nielsen MB, et al. The diagnostic and prognostic value of ultrasonography in soccer players with acute hamstring injuries. *Am J Sports Med* 2014;42:399–404.
- 16 Moen MH, Reurink G, Weir A, et al. Predicting return to play after hamstring injuries. *Br J Sports Med* 2014;48:1358–63.
- 17 Warren P, Gabbe BJ, Schneider-Kolsky M, et al. Clinical predictors of time to return to competition and of recurrence following hamstring strain in elite Australian footballers. *Br J Sports Med* 2010;44:415–19.
- 18 De Vos RJ, Reurink G, Goudswaard GJ, et al. Clinical findings just after return to play predict hamstring re-injury, but baseline MRI findings do not. *Br J Sports Med* 2014;48:1377–84.
- 19 Brughelli M, Cronin J, Mendiguchia J, et al. Contralateral leg deficits in kinetic and kinematic variables during running in Australian rules football players with previous hamstring injuries. *J Strength Cond Res* 2010;24:2539–44.
- 20 Lee MJ, Reid SL, Elliott BC, et al. Running biomechanics and lower limb strength associated with prior hamstring injury. *Med Sci Sports Exerc* 2009;41:1942–51.
- 21 Opar DA, Williams MD, Timmins RG, et al. Knee flexor strength and bicep femoris electromyographical activity is lower in previously strained hamstrings. *J Electromyogr Kinesiol* 2013;23:696–703.
- 22 Sanfilippo JL, Silder A, Sherry MA, et al. Hamstring strength and morphology progression after return to sport from injury. *Med Sci Sports Exerc* 2013;45:448–54.
- 23 Sole G, Milosavljevic S, Nicholson H, et al. Altered muscle activation following hamstring injuries. *Br J Sports Med* 2012;46:118–23.
- 24 Sole G, Milosavljevic S, Nicholson HD, et al. Selective strength loss and decreased muscle activity in hamstring injury. *J Orthop Sports Phys Ther* 2011;41:354–63.
- 25 Tol JL, Hamilton B, Eirale C, et al. At return to play following hamstring injury the majority of professional football players have residual isokinetic deficits. *Br J Sports Med* 2014;48:1364–9.
- 26 Brockett CL, Morgan DL, Proske U. Predicting hamstring strain injury in elite athletes. *Med Sci Sports Exerc* 2004;36:379–87.
- 27 Croisier JL, Forthomme B, Namurois MH, et al. Hamstring muscle strain recurrence and strength performance disorders. *Am J Sports Med* 2002;30:199–203.
- 28 Silder A, Thelen DG, Heiderscheidt BC. Effects of prior hamstring strain injury on strength, flexibility, and running mechanics. *Clin Biomech (Bristol, Avon)* 2010;25:681–6.
- 29 Emami M, Arab AM, Ghamkhar L. The activity pattern of the lumbo-pelvic muscles during prone hip extension in athletes with and without hamstring strain injury. *Int J Sports Phys Ther* 2014;9:312–19.
- 30 Opar DA, Williams MD, Timmins RG, et al. Rate of torque and electromyographic development during anticipated eccentric contraction is lower in previously strained hamstrings. *Am J Sports Med* 2013;41:116–25.
- 31 Fyfe JJ, Opar DA, Williams MD, et al. The role of neuromuscular inhibition in hamstring strain injury recurrence. *J Electromyogr Kinesiol* 2013;23:523–30.
- 32 Mendiguchia J, Samozino P, Martinez-Ruiz E, et al. Progression of mechanical properties during on-field sprint running after returning to sports from a hamstring muscle injury in soccer players. *Int J Sports Med* 2014;35:690–5.
- 33 Jonhagen S, Nemeth G, Eriksson E. Hamstring injuries in sprinters. The role of concentric and eccentric hamstring muscle strength and flexibility. *Am J Sports Med* 1994;22:262–6.
- 34 Opar DA, Williams MD, Shield AJ. Hamstring strain injuries: factors that lead to injury and re-injury. *Sports Med* 2012;42:209–26.
- 35 Silder A, Heiderscheidt BC, Thelen DG, et al. MR observations of long-term musculotendon remodeling following a hamstring strain injury. *Skeletal Radiol* 2008;37:1101–9.
- 36 Timmins R, Porter K, Williams M, et al. Biceps femoris muscle architecture—the influence of previous injury. *Br J Sports Med* 2014;48:665–6.
- 37 Timmins RG, Shield AJ, Williams MD, et al. Biceps femoris long-head architecture: a reliability and retrospective injury study. *Med Sci Sports Exerc* 2015;47:905–13.
- 38 Petersen J, Thorborg K, Nielsen MB, et al. Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. *Am J Sports Med* 2011;39:2296–303.
- 39 van der Horst N, Smits DW, Petersen J, et al. The preventive effect of the Nordic hamstring exercise on hamstring injuries in amateur soccer players: study protocol for a randomised controlled trial. *Inj Prev* 2014;20:e8.
- 40 Malliaropoulos N, Mendiguchia J, Pehlivanidis H, et al. Hamstring exercises for track and field athletes: injury and exercise biomechanics, and possible implications for exercise selection and primary prevention. *Br J Sports Med* 2012;46:846–51.
- 41 Guex K, Millet GP. Conceptual framework for strengthening exercises to prevent hamstring strains. *Sports Med* 2013;43:1207–15.
- 42 Sherry MA, Best TM. A comparison of 2 rehabilitation programs in the treatment of acute hamstring strains. *J Orthop Sports Phys Ther* 2004;34:116–25.
- 43 Askling CM, Tengvar M, Tarassova O, et al. Acute hamstring injuries in Swedish elite sprinters and jumpers: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med* 2014;48:532–9.
- 44 Reurink G, Goudswaard GJ, Moen MH, et al. Platelet-rich plasma injections in acute muscle injury. *N Engl J Med* 2014;370:2546–7.
- 45 A Hamid MS, Mohamed Ali MR, Yusof A, et al. Platelet-rich plasma injections for the treatment of hamstring injuries: a randomized controlled trial. *Am J Sports Med* 2014;42:2410–18.
- 46 de Jonge S, De Vos RJ, Weir A, et al. One-year follow-up of platelet-rich plasma treatment in chronic achilles tendinopathy: a double-blind randomized placebo-controlled trial. *Am J Sports Med* 2011;39:1623–9.
- 47 de Vos RJ, Weir A, van Schie HTM, et al. Platelet-rich plasma injection for chronic achilles tendinopathy. A randomized controlled trial. *JAMA* 2010;303:144–9.
- 48 Bennell K, Wajswelner H, Lew P, et al. Isokinetic strength testing does not predict hamstring injury in Australian Rules footballers. *Br J Sports Med* 1998;32:309–14.
- 49 Freckleton G, Cook J, Pizzari T. The predictive validity of a single leg bridge test for hamstring injuries in Australian Rules Football Players. *Br J Sports Med* 2014;48:713–17.
- 50 Opar DA, Piatkowski T, Williams MD, et al. A novel device using the Nordic hamstring exercise to assess eccentric knee flexor strength: a reliability and retrospective injury study. *J Orthop Sports Phys Ther* 2013;43:636–40.
- 51 Opar DA, Williams MD, Timmins RG, et al. Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med Sci Sports Exerc* 2015;47:857–65.
- 52 Goossens L, Witvrouw E, Vanden Bossche L, et al. Lower eccentric hamstring strength and single leg hop for distance predict hamstring injury in PETE students. *Eur J Sport Sci* 2015;15:436–42.
- 53 Schache AG, Crossley KM, Macindoe IG, et al. Can a clinical test of hamstring strength identify football players at risk of hamstring strain? *Knee Surg Sports Traumatol Arthrosc* 2011;19:38–41.
- 54 Mendiguchia J, Alentorn-Geli E, Brughelli M. Hamstring strain injuries: are we heading in the right direction? *Br J Sports Med* 2012;46:81–5.
- 55 Askling C, Karlsson J, Thorstensson A. Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. *Scand J Med Sci Sports* 2003;13:244–50.
- 56 Arnason A, Andersen TE, Holme I, et al. Prevention of hamstring strains in elite soccer: an intervention study. *Scand J Med Sci Sports* 2008;18:40–8.
- 57 Brooks JH, Fuller CW, Kemp SP, et al. Incidence, risk, and prevention of hamstring muscle injuries in professional rugby union. *Am J Sports Med* 2006;34:1297–306.
- 58 Gabbe BJ, Branson R, Bennell KL. A pilot randomised controlled trial of eccentric exercise to prevent hamstring injuries in community-level Australian Football. *J Sci Med Sport* 2006;9:103–9.
- 59 Seagrave RAI, Perez L, McQueeney S, et al. Preventive effects of eccentric training on acute hamstring muscle injury in professional baseball. *Orth J Sports Med* 2014;2.
- 60 Melegati G, Tornese D, Gevi M, et al. Reducing muscle injuries and reinjuries in one Italian professional male soccer team. *Muscles Ligaments Tendons J* 2013;3:324–30.