



OPEN ACCESS

'Physical Activity 4 Everyone' school-based intervention to prevent decline in adolescent physical activity levels: 12 month (mid-intervention) report on a cluster randomised trial

Rachel Sutherland,^{1,2,3} Elizabeth Campbell,^{1,2,3} David R Lubans,⁴ Philip J Morgan,⁴ Anthony D Okely,^{5,6} Nicole Nathan,^{1,2,3} Luke Wolfenden,^{1,2,3} Jarrod Wiese,^{1,2} Karen Gillham,^{1,3} Jenna Hollis,⁷ John Wiggers^{1,2,3}

¹Hunter New England Population Health, Wallsend, New South Wales, Australia
²School of Medicine and Public Health, University of Newcastle, Newcastle, New South Wales, Australia
³Hunter Medical Research Institute, Newcastle, New South Wales, Australia
⁴Priority Research Centre in Physical Activity and Nutrition, School of Education, University of Newcastle, Newcastle, New South Wales, Australia
⁵Early Start Research Institute and School of Education, University of Wollongong, Wollongong, New South Wales, Australia
⁶Illawarra Health and Medical Research Institute, Wollongong, New South Wales, Australia
⁷Rowett Institute of Nutrition and Health, University of Aberdeen, Scotland, UK

Correspondence to

Rachel Sutherland, Hunter New England Population Health, Locked Bag No. 10, Wallsend NSW 2287, Australia; Rachel.Sutherland@hnehealth.nsw.gov.au

Accepted 17 June 2015
Published Online First
10 September 2015



Open Access
Scan to access more
free content



CrossMark

To cite: Sutherland R, Campbell E, Lubans DR, et al. *Br J Sports Med* 2016;**50**:488–495.

ABSTRACT

Background Adolescence is a recognised period of physical activity decline, particularly among low-income communities. We report the 12-month (midpoint) effects of a 2-year multicomponent physical activity intervention implemented in disadvantaged secondary schools.

Methods A cluster randomised trial was undertaken in 10 secondary schools located in disadvantaged areas in New South Wales, Australia. Students in Grade 7 were recruited, with follow-up in Grade 8. The intervention was guided by socioecological theory and included seven physical activity strategies, and six implementation adoption strategies. The primary outcome was mean minutes of moderate-to-vigorous physical activity (MVPA) per day assessed using Actigraph GT3X accelerometers. Outcome data were analysed using repeated measures linear mixed models.

Results At baseline, 1150 (93%) students participated in the data collection (mean age 12 years, 48% boys) and 1050 (79%) students participated at 12-month follow-up. By the 12-month follow-up, the six implementation adoption strategies had been used to support schools to deliver four of the seven physical activity elements. There was a significant group-by-time interaction for mean minutes of MVPA per day in favour of the intervention group (adjusted difference between groups at follow-up=3.85 min, 95% CI (0.79 to 6.91), $p \leq 0.01$), including significantly more vigorous physical activity (2.45 min, $p \leq 0.01$), equating to 27 min more MVPA per week.

Summary At 12-month follow-up, the intervention had reduced the decline in physical activity among adolescents from disadvantaged schools. The intervention may assist students to meet physical activity guidelines.

BACKGROUND

Participation in adequate physical activity has numerous physical and psychological health benefits.¹ Despite this, the proportion of adolescents who are adequately active is consistently low, with as few as 20% meeting physical activity guidelines of 60 min MVPA per day.² International data indicate a significant inverse association between physical activity and socioeconomic status (SES), with adolescents from disadvantaged backgrounds experiencing a steeper decline in physical activity.^{3,4}

As physical inactivity tends to track into adulthood, reducing this decline is a public health priority.⁵

Schools provide access to almost all adolescents over extended periods of time.^{6–8} Schools have qualified staff such as physical education (PE) teachers, resources including sporting equipment and facilities, and a mandate to implement curriculum that promotes physical activity.⁸ Based on a number of systematic reviews,^{9–12} there is evidence that school-based interventions are effective in increasing the proportion of students who are physically active, the length of time spent being active, and student fitness levels.^{9,10,13,14} However, such evidence is primarily focused on children of elementary school age (5–12 years), with very few studies focusing on adolescents.⁹

Systematic reviews of physical activity interventions for children and adolescents^{9–12} conclude that interventions were more likely to be successful if they were multicomponent, longer in duration and based on theory.^{9,10} Such reviews recommended that future trials include the use of an objective measure of physical activity, measurement of total daily physical activity, use clear intervention implementation strategies, focus on low-socioeconomic groups, focus on interventions targeting adolescents, have an intervention duration spanning greater than 12 months, and employ longer follow-up.

Of the fourteen interventions targeting school-based physical activity in adolescents, only five studies have specifically targeted disadvantaged secondary school students. Of these, three have tested single sex interventions, and the studies did not demonstrate a positive intervention effect on physical activity.^{16–18} The two remaining intervention trials both used an objective measure of physical activity and were able to demonstrate an intervention effect. However, both interventions were of short duration, 17 weeks¹⁹ and 6 months, respectively.²⁰

Given the limited number of effective interventions targeting greater physical activity among adolescents from disadvantaged backgrounds, the primary aim of this study was to report on the 12-month, mid-intervention impact of a 2-year multicomponent physical activity intervention implemented in disadvantaged secondary schools, which aimed to reduce the decline in physical activity associated with adolescence. Subgroup analyses for sex, baseline weight status and baseline activity level are also reported.

METHODS

Study design and setting

The Physical Activity 4 Everyone (PA4E1) study was a multi-component school-based cluster randomised trial with study assessments conducted at baseline, 12 months (mid-intervention) and 24 months. The trial was conducted in three local government areas (Hunter, Central Coast and Mid-North Coast) in the state of New South Wales (NSW), Australia. The regions have lower average indices of socioeconomic status than the state.²¹ The trial was registered with the Australian New Zealand Clinical Trials Registry (ACTRN1261200038287) and approved by the Hunter New England Area Human Research Ethics Committee (11/03/16/4.0), and University of Newcastle Human Research Ethics Committee (H-2011-0210). A trial protocol has been published elsewhere.²¹ The study adheres to the Consolidated Standards of Reporting Trials (CONSORT) guidelines (<http://www.consort-statement.org>).

Participants and recruitment

Secondary schools

The Socioeconomic Indexes for Areas (SEIFA) of relative socioeconomic disadvantage were used to identify eligible secondary schools.²¹ The SEIFA (scale: 1=lowest to 10=highest) summarises the characteristics of people and households within an area and is based on postcode. Secondary schools were considered eligible if they met the following criteria: Government and Catholic schools; schools with a SEIFA score of 5 or less (bottom 50% of NSW)²²; between 120 and 200 year 7 students (to meet sample size requirements); and were not participating in other physical activity intervention studies. Recruitment of schools occurred from October to December 2011. An invitation to participate was sent to the first 10 randomly selected schools. Thirteen schools were approached to obtain a sample of 10 schools.

Students

All students in Grade 7 (first year of secondary school) at participating schools were invited to take part via an information package sent to their parents. Parental consent was obtained via returned consent form. If a consent form was not obtained, parents were contacted via telephone and asked to provide consent.

Teachers

All PE teachers at intervention schools were invited to complete a pen and paper survey. Consent was obtained via return survey.

Randomisation and allocation

Randomisation and allocation of schools to the intervention or control group occurred after baseline data collection. Using block randomisation (1:1 ratio), schools were allocated based on a random number function in Microsoft Excel. Schools were randomly allocated to receive either a multicomponent intervention that was implemented during school terms and started after baseline data collection in June 2012, or to a control group.

Intervention and comparison

The 24-month PA4E1 intervention was designed as a multicomponent school-based programme guided by social cognitive²³ and social-ecological theories.²⁴ The strategies implemented in the intervention addressed the domains of the WHO's Health Promoting Schools framework targeting the curriculum, school environment and community.^{15 25-28}

The intervention comprised seven physical activity strategies to be implemented in a staged fashion over the intervention period (see figure 1 for strategies and the standards set to be achieved for each). The strategies were: 'Formal Curriculum'— (1) teaching strategies to maximise activity in PE lessons, including pedometer-based lessons,²⁹ (2) development of individual




Figure 1 Intervention delivery baseline to 12-month follow-up.

student physical activity plans, (3) enhanced school sport for all students (to be delivered in Grade 8);^{20–30} ‘School Ethos and Environment’—(4) school physical activity policies, (5) offering physical activity in school breaks (lunch and recess); ‘Partnerships and Services’—(6) linking schools to community physical activity providers, (7) parent engagement. By 12-month follow-up, implementation duration within intervention schools ranged from two to three school terms (each term was 10 weeks in duration). Implementation of four of the seven physical activity strategies started (strategies 1, 2, 5, 7 above).

Figure 1 summarises the physical activity strategies delivered, the adoption strategies used to facilitate their delivery and the desired standard and dose delivered within the first 12 months of the intervention period. In the first 12-month period, two curriculum-based strategies started, including teaching strategies to maximise activity in PE and the development of individualised student physical activity plans. To facilitate adoption, PE teachers were provided training on strategies to maximise physical activity in PE, prompted by the change agent to teach pedometer-based lessons and support students to complete personalised physical activity plans, were given resources (such as templates and instructions for use) to support students in developing personal activity plans, and given feedback on activity levels in PE based on SOFIT observations. One strategy targeting the school ethos and environment started within the first 12-month period. Offering physical activity in school breaks (lunch and/or recess) started in each school twice per week. Schools were provided equipment, including a variety of balls, hoops and ropes in a secure locked box to facilitate the start of these activities. The final strategy to start within the first 12-month period focused on parent engagement, whereby hard copy newsletters and websites were used to provide parents with updates in the programme being implemented at school plus articles about ways to support students to be active outside of the school. To facilitate the adoption of these physical activity strategies, schools established committees to oversee the changes and were provided with feedback reports outlining the schools progress towards programme adoption at the end of each term. Meetings were held with school executives, PE teachers and the school change agent to communicate the content of each feedback report.

Comparison

Schools allocated to the control group participated in the measurement components of the study only. They were asked to continue with their usual physical activity practices, including time table-based Health and Physical Education lessons, school sport, breaks for recess and lunch and any scheduled professional development for teachers.

Data collection procedures

Data collection was undertaken by trained research assistants, blinded to group allocation. Baseline data were collected from March to June 2012, and 12-month follow-up data (mid-intervention) data collected from the same cohort of students 12 months later in March–June 2013. The average duration between baseline and follow-up measurements for all schools was 12 months.

Measures

Outcome measures: physical activity levels

Physical activity was measured using accelerometers (Actigraph GT3X+ and GT3X models).^{31–33} Mean minutes of MVPA per day was the primary outcome. Additional outcome measures

included: (1) percentage of time spent in MVPA per day (calculated to adjust for individual wear time), (2) mean minutes per day and percentage wear time for moderate physical activity, (3) mean minutes per day and percentage of wear time in vigorous physical activity, (4) accelerometer counts per minute (CPM). Counts were collected in 15 s epochs and CPM calculated by dividing the total counts per day by the minutes of wear time. The proportion of students meeting physical activity guidelines of 60 min of MVPA/day has also been reported.

Accelerometers and instructions for use were distributed to students within class time when students also completed an online survey and had anthropometric measures taken. Students were requested to wear the accelerometer over the right hip during waking hours for seven consecutive days. Student and parent mobile numbers were collected via the consent form, and these were used to text daily reminders to wear the accelerometer. Student data were included in the analysis if the accelerometer was worn for ≥ 600 min on ≥ 3 days.³⁴ Non-wear time was defined as 30 min of consecutive zeros.³⁵ The Everson cut-points were used to categorise different intensities of physical activity.³⁶

Anthropometric data

Student anthropometric data, including height, weight (used to calculate body mass index (BMI)) and waist circumference was collected in duplicate using the International Society for the Advancement of Kinanthropometry (ISAK) procedures.³⁷ Weight was measured in light clothing without shoes using a portable digital scale (Model no. UC-321PC, A&D Company Ltd., Tokyo, Japan) to the nearest 0.1 kg. Height was recorded to the nearest 0.1 cm using a portable stadiometer (Model no. PE087, Mentone Educational Centre, Australia). Weight status (BMI) was determined using International Obesity Taskforce definitions.³⁸ Waist measurement was taken at the narrowest point between the inferior rib border and the iliac crest, using a flexible but inelastic tape measure. Waist circumference was recorded to the nearest 0.1 cm.

Student characteristics

Students completed an online survey that assessed student socio-demographic characteristics, including age, sex, Aboriginal or Torres Strait Islander status, and postcode of residence. The online survey also included other measures that were not included in the current paper (eg, physical activity mediators).

Process measures

A process evaluation was conducted to determine if the intervention was delivered (fidelity) and received (reach) as intended. At the 12-month follow-up, PE teachers completed a pen and paper survey that assessed intervention fidelity by asking about delivery of three physical activity strategies; implementing pedometer-based PE lessons and termly student physical activity plans with their classes, and whether the school offered recess and/or lunch activities. The school change agent also retained records of intervention implementation at each school. These records were used to determine if programme strategies were implemented to the desired standard outlined in figure 1. This included records of lessons in which pedometers had been used, personal PA plans developed by students, recess and/or lunch physical activities run at each school, and information in newsletters. Students in the intervention group completed online survey items at 12-month follow-up that aimed to assess the reach of three intervention strategies: pedometer-based PE lessons, termly physical activity plans, and availability of recess

and/ or lunch activities. The school change agent also kept records of the adoption strategies implemented by schools, including committee meetings held, teacher training attendance, equipment/resources received by schools and prompts sent to teachers.

Sample size calculations

Based on an estimate of 120 students per school and 50% of year 7 students consenting, it was estimated each school should yield at least 60 students, providing at least 300 students per group.^{39–40} Based on 65% of the cohort providing usable data at 24-month follow-up, it was estimated that there would be at least 195 students per group at follow-up.⁴¹ Previous studies were used to estimate the SD of mean daily minutes MVPA per group (17.1)⁴² and the intraclass correlation coefficient (ICC; 0.01).⁴³ After adjustment for the design effect of 1.38, the effective sample size was estimated to be at least 141 students per group. With this sample size, 80% power and an α level of 0.05, the study was able to detect a difference in the mean daily MVPA between intervention and control students of ± 5.73 min at follow-up.

Statistical analysis

All analyses were conducted using SAS V9.2 (SAS Institute Inc, Cary, North Carolina, USA). Summary statistics were created for the variables of interest (student sex, age, aboriginality, height, weight, BMI, activity level, SES) and accelerometer wear time. T tests were used to determine if students who provided data at 12-month follow-up differed to those that only provided baseline data on the following characteristics—sex, baseline age, weight status and physical activity level. Significance levels were set at $p \leq 0.05$.

Physical activity change

Analyses followed intention-to-treat principles. Analysis of the primary outcome (minutes of MVPA/day), and of the additional physical activity outcome variables (% of wear time spent in MVPA/day; mean minutes and % wear time in moderate physical activity and vigorous physical activity and accelerometer CPM) were facilitated through a linear mixed model (LMM). These statistical models are preferable as they are robust to the biases of missing data.⁴⁴ A three-level hierarchical model was used to capture the correlations in the data with random intercepts for repeated measures (level-1) on individuals (level-2) and clustering within schools (level-3). LMM analysis was used to determine whether the change in physical activity between intervention and control groups differed significantly after 12 months, assessed through an interaction term between group (intervention vs control) and time (baseline vs follow-up). The data were analysed assuming data were ‘missing at random’. Descriptive statistics were used to describe the proportion of students in each group meeting the physical activity guidelines of 60 min MVPA per day.

Subgroups analyses

Sex, baseline weight status and baseline activity level were the variables chosen a priori as these are common moderators of energy balance interventions.⁴⁵ Students’ baseline BMI were categorised into two groups: ‘underweight/healthy weight’ and ‘overweight/obese’ based on the Cole cut-points.³⁸ Baseline student activity level was categorised as those who obtained 60 min or more of MVPA per day (meeting the guidelines), and those with less than 60 min of MVPA each day (not meeting the physical activity guidelines). We included moderator interaction

terms in the above LMM separately for all potential moderators and presented the results by mediator subgroup if the test for three-way interaction term (group \times time \times moderator) was significant at the liberal 20% threshold.⁴⁶

Process measures

χ^2 Square analyses were used to assess whether student responses to process variables differed by student subgroups of sex, baseline physical activity level and baseline weight status ($p=0.05$).

RESULTS

Sample

Ten schools were recruited to the study, which included four Government and one Catholic secondary school in the intervention group and control group. Thirty-three PE teachers (100%) in intervention schools completed the pen and paper survey. Parental consent was received from 1233 of the 1468 (84%) year 7 students. Figure 2 outlines the flow of participants from recruitment to 12-month follow-up. Baseline characteristics of the 1150 students who wore an accelerometer (93% of those with parental consent) are outlined in table 1.

At baseline, 78% of those students who wore an accelerometer provided at least three days of valid accelerometer data (965/1150). At 12-month follow-up, 1050 students wore an accelerometer and 61% of these students provided at least 3 days of valid accelerometer data (643/1050). We found baseline weight and age were predictive of drop out at 12 months, with higher BMI and younger students more likely to drop out ($p \leq 0.001$ and $p \leq 0.001$, respectively). A sensitivity analysis was conducted on the main outcome, adjusting for baseline weight and age, with minimal difference in the result detected; therefore, unadjusted results are presented.

Individual level physical activity changes


Physical activity outcomes from baseline to 12-month follow-up are presented in table 2. At 12-month follow-up, students in the intervention group participated in statistically significant more minutes per day of MVPA than students in the control group (adjusted difference=3.85 min (0.79 to 6.91), $p=0.01$).

The intervention group spent significantly more time in vigorous activity each day (adjusted difference=2.45 min (0.90 to 4.00), $p \leq 0.01$), but not moderate physical activity. The percent time spent in MVPA (0.5% (0.11 to 0.90)) and vigorous activity (0.3%) (0.12 to 0.52)) also differed significantly between groups at 12 month follow-up ($p=0.01$ and $p \leq 0.01$, respectively) in favour of the intervention group. Mean accelerometer CPM was significantly different between groups at 12-month follow-up in favour of the intervention group (31.02 CPM, (9.05 to 53.00), $p=0.01$). The proportion of students meeting the physical activity guidelines were 33% at baseline and 34% at 12-month follow-up in the intervention group, and 34% at baseline and 28% at 12-month follow-up in the control group.

Changes in physical activity from baseline to follow-up across subgroups (sex, baseline weight status and baseline activity level)

The subgroup interaction term indicated time by intervention effects that differed by subgroup for each variable: sex ($p \leq 0.01$), baseline weight status ($p \leq 0.01$) and baseline physical activity status ($p \leq 0.01$); therefore subgroup analyses were progressed for each. The 12-month physical activity analyses by

Figure 2 CONSORT flow chart describing progress of participants through the study.



subgroup are reported in table 3. A greater effect was observed in male students in the intervention group compared with the control group on mean minutes of MVPA per day (6.47 min (1.24 to 12.95), $p=0.02$) and percentage of wear time spent in MVPA (0.9%, $p=0.02$). No significant differences between groups for females were observed at 12-month follow-up. There were no detected differences between intervention and control based on weight status or activity level detected.

Process measures

Table 4 outlines process evaluation data collected from teachers and students at 12-month follow-up. At 12-month follow-up, 95.5% of teachers reported using pedometers to increase activity levels in PE, 70.3% reported incorporating student personal physical activity plans each term and providing feedback to

students on these plans, and 75% reported the school offered organised physical activity at recess and/or lunchtimes, at least twice per week. The school change agent records showed that all schools had started use of the pedometers in PE classes, with four of the five schools (80%) using them with the desired frequency (figure 1). Similarly, while all schools had administered student physical activity plans at least once, three (60%) had administered these as per the desired termly standard (2 or 3 per student). All schools had implemented recess and/or lunch activities at least once per week, and four (80%) had these implemented at least twice per week. All schools had provided parents with additional information regarding physical activity via newsletters and the school website with the requested termly frequency.

At 12-month follow-up, 92.7% of students reported being offered pedometer-based PE lessons at least twice per term,

Table 1 PA4E1 sample characteristics at baseline

Characteristic	Intervention group	Control group
Number/total participants	645	505
Boys*	290	239
Girls*	317	254
3 vld days	524	435
Mean age (years)	12.0	12.0
Aboriginal and/or Torres Strait Islander (%)	5.3	7.8
Height, (mean m)	157.1	156.8
Weight, (mean kg)	49.3	50.0
<i>Student BMI category, (%)</i>		
Underweight/healthy weight		
All students	77.5%	74.0%
Boys	78.3%	74.4%
Girls	77.2%	73.7%
Overweight/obese		
All students	21.8%	26.0%
Boys	21.9%	24.4%
Girls	22.9%	26.9%
<i>Student activity level</i>		
Active (≥ 60 min MVPA/day)		
All students	33%	33%
Boys	50%	42%
Girls	48%	40%
Low active (< 60 min MVPA/day)		
All students	67%	67%
Boys	50%	58%
Girls	52%	60%
<i>Socioeconomic status</i>		
Low SES	57.8%	61%
<i>Accelerometer wear time</i>		
Mean minutes per day	793.6	804.6

*Does not add to total students (n=645) due to 38 students having gender missing. BMI, body mass index; MVPA, moderate-to-vigorous physical activity; PA4E1, Physical Activity 4 Everyone; SES, socioeconomic status.

51.6% reported completing a personal physical activity plan at least once, and 55.8% reported that the school offered organised physical activity at recess and/or lunch. When the results were compared for male and female students, and for students grouped according to baseline weight and physical activity status, the only statistically significant difference was that male students were more likely than female students to report the school offered recess and/or lunch physical activities (61.5% vs 50.9% ($p=0.03$); table 4).

The adoption strategies outlined in figure 1 were being used consistently in all intervention schools. All schools had formed committees to oversee the implementation of physical activity strategies and had at least 2–3 meetings; the school change agent attended each school for 1 day per week and all schools had at least one staff member attend the professional development (range 1–4 staff). The school change agent sent weekly prompts to PE teachers encouraging pedometer-based PE lessons and completion of student physical activity plans. A range of equipment to facilitate recess and/or lunch activities and a storage box were delivered to each school, and feedback reports outlining progress against each strategy were delivered and discussed with school executives and the head PE teacher at the end of each school term.

DISCUSSION

We report the 12-month mid-intervention findings from a multi-component physical activity intervention implemented in disadvantaged secondary schools. At 12-month follow-up, students attending intervention schools participated in nearly 4 min more MVPA per day than control group students. To some readers, this may not sound like a clinically meaningful difference. However, it represents 27 min more of MVPA over the course of a week.

Small, but clinically significant effect at 1 year

Research in children and adolescents has identified a dose-response relationship between the total volume of MVPA and a reduction in cardiometabolic risk; therefore, any increase in MVPA has public health benefit.⁴⁷ Students in the intervention group participated in significantly more vigorous activity and spent a greater proportion of time in MVPA and vigorous activity each day. We suggest that this magnitude of change in physical activity, particularly the increase in vigorous activity, is clinically meaningful, and may facilitate the prevention of chronic disease such as type 2 diabetes and obesity.^{48–50}

The results displayed at 12 months extend the results described in a meta-analysis of physical activity interventions in children and adolescents;⁵¹ however, most interventions focus on children and few on adolescents. As a result, the effect size seems larger than other school-based interventions targeting adolescents. Of the interventions targeting adolescents that have been effective,^{19 20 52–54} two studies published mid-intervention findings, both of which showed no significant intervention effect.^{53 52} Other school-based interventions targeting adolescents demonstrated positive postintervention findings in favour of the intervention group, with effect sizes

Table 2 Changes in physical activity from baseline to 12-month follow-up (minutes MVPA, % wear time in MVPA, % meeting PA Guidelines)

Outcome	Intervention			Control			Intervention-Control Adjusted difference between treatment group (95% CI)	Group \times time p value
	BASELINE (n=524)	MIDPOINT (n=352)	p Value	BASELINE (n=435)	MIDPOINT (n=288)	p Value		
<i>Minutes of physical activity (mean min/day)</i>								
Total MVPA	53.3	54.2	0.55	53.6	50.8	0.07	3.85 (0.79 to 6.91)	0.01*
Vigorous activity	16.6	18.0	0.07	16.9	16.2	0.37	2.45 (0.90 to 4.00)	0.002
Moderate activity	36.7	36.2	0.52	36.7	34.6	0.02	1.41 (–0.50 to 3.33)	0.15
Counts per minute	482.2	476.2	0.58	486.1	452.3	0.002	31.02 (9.05 to 53.00)	0.01
<i>Percentage of wear time</i>								
Percentage MVPA	6.7	6.9	0.44	6.7	6.4	0.10	0.50 (0.11 to 0.90)	0.01
Percentage vigorous	2.1	2.3	0.05	2.1	2.0	0.40	0.32 (0.12 to 0.52)	0.001
Percentage moderate	4.6	4.6	0.70	4.6	4.4	0.04	0.18 (–0.06 to 0.43)	0.15

Table 3 Changes in physical activity from baseline to follow-up (12 months) by subgroup (gender, weight status at baseline and activity level at baseline) (Mean minutes of MVPA per day, % wear time in MVPA, % meeting physical activity guidelines)

Subgroups	Outcome	Intervention			Control			Intervention-Control Adjusted difference between treatment group (95% CI)	Group× time p value
		BASELINE (n=524)	MIDPOINT (n=352)	p Value	BASELINE (n=435)	MIDPOINT (n=288)	p Value		
Gender									
Males	MVPA (min per day)	62.7 (22.54)	66.3 (27.19)	0.17	59.4 (23.32)	56.6 (19.81)	0.27	6.47 (−1.84 to 14.78)	0.02*
	Percentage of wear time in MVPA	8.0 (2.89)	8.5 (3.72)	0.19	7.5 (2.88)	7.2 (2.57)	0.34	0.90 (−0.13 to 1.93)	0.02*
Females	MVPA (min per day)	46.6 (16.45)	45.7 (15.50)	0.55	48.9 (17.63)	45.8 (17.71)	0.09	−0.94 (−5.62 to 3.74)	0.35
	Percentage of wear time in MVPA	5.8 (2.06)	5.8 (1.98)	0.74	6.1 (2.21)	5.7 (2.17)	0.14	−0.06 (−0.55 to 0.42)	0.34
Weight status at baseline									
Underweight/ healthy weight	MVPA (min per day)	54.9 (21.83)	54.9 (21.98)	0.98	55.4 (21.72)	53.1 (19.22)	0.25	1.82 (−4.22 to 7.87)	0.10
	Percentage of wear time in MVPA	6.9 (2.84)	6.9 (2.89)	0.95	6.9 (2.71)	6.7 (2.48)	0.52	0.28 (−0.39 to 0.95)	0.25
Overweight/obese	MVPA (min per day)	49.8 (17.82)	50.3 (21.23)	0.88	49.3 (19.47)	44.7 (15.68)	0.12	1.74 (−5.80 to 9.29)	0.29
	Percentage of wear time in MVPA	6.2 (2.07)	6.4 (2.79)	0.63	6.2 (2.49)	5.5 (1.88)	0.07	0.28 (−0.57 to 1.12)	0.14
Activity level at baseline									
Active	MVPA (min per day)	76.9 (16.70)	78.7 (21.06)	0.43	77.1 (15.49)	75.4 (13.52)	0.41	4.54 (−1.22 to 10.30)	0.12
	Percentage of wear time in MVPA	9.6 (2.31)	10.1 (2.97)	0.16	9.5 (2.11)	9.3 (1.89)	0.65	0.64 (−0.15 to 1.44)	0.11
Inactive	MVPA (min per day)	41.9 (10.80)	41.7 (11.50)	0.77	41.5 (10.67)	41.2 (11.55)	0.77	−0.21 (−2.42 to 1.99)	0.85
	Percentage of wear time in MVPA	5.4 (1.44)	5.3 (1.54)	0.53	5.3 (1.49)	5.2 (1.52)	0.71	−0.04 (−0.34 to 0.26)	0.81

MVPA, moderate-to-vigorous physical activity.

ranging from 1.9 min of MVPA per day after 2 years of intervention in the TAAG study,⁵³ 3.5 min MVPA per day for males only in the 2-year study by Haerens *et al*,⁵² and 50 accelerometer CPM after the 20-month Health in Adolescence study (HEIA).⁵⁴

More recently, three interventions^{16–18} have specifically targeted adolescents from lower socioeconomic backgrounds; however, none have shown a significant intervention effect on MVPA. Given the challenges in conducting intervention research targeting disadvantaged adolescents and schools, a

positive mid-intervention effect demonstrates potential to intervene with this target group.

Innovations in this study

The PA4E1 intervention differed from previous trials as it focused on students attending schools located in disadvantaged areas, targeted the whole school community while incorporating strategies to engage low-active students. In addition, the PA4E1 intervention was longer in duration, and included a school change agent position within a set of clear adoption

Table 4 Intervention fidelity and reach at 12-months follow-up

Process measure category	Physical activity strategies implemented from baseline—12-month follow-up		
	Formal curriculum		School ethos & environment
	Active PE lessons (%)	Personal physical activity plans (%)	Recess and lunchtime activity (%)
Fidelity (teacher report n=33)	95.5*	70.3†	75.0‡
Reach (student report n=600)			
All students	92.7§	51.6¶	55.8**
Student sex			
Female students	93.2	49.6	50.9*
Male students	95.0	54.0	61.5*
Student activity level			
Active students	96.3	56.8	59.9
Inactive students	93.6	47.8	52.6
Student weight status			
Healthy weight students	95.3	50.3	55.5
Overweight/obese students	94.4	52.8	56.2

*Teacher reports conducting pedometer-based lessons.

†Teacher reports assisting student complete a personal PA plan each school term.

‡School reports running recess and/or lunch activities.

§Students recall using pedometers in PE.

¶Students recall completing personal physical activity plans.

**Students recall having organised recess and/or lunchtime physical activities available.

PA, physical activity; PE, physical education.

strategies. Including explicit implementation strategies, as recommended in systematic reviews,^{10 15} may explain our positive findings.

In particular, the change agent, someone located within the school 1 day per week to support schools in implementing the physical activity strategies (not to deliver them) is novel. Schools often report time and demanding workloads as barriers to implementing intervention strategies.^{55–57} The addition of a change agent aims to overcome these barriers, and maximise intervention reach and fidelity. The addition of a further three physical activity strategies to the PA4E1 intervention in the second phase of intervention, may enhance the likelihood of sustained success. Systematic reviews have concluded that study duration, study size and positive mid-intervention results are associated with a significant intervention effect at follow-up.⁵¹

Limitations

As subgroup analyses were exploratory due to limited power, results should be seen as suggestive and interpreted with caution. At 12-month follow-up, the results were only statistically significant for boys—the intervention appeared to be effective for male students, but there was no significant effect among females. These results are in contrast to a systematic review by Yildirim⁴⁵ finding girls responded better to interventions than boys. If our midpoint assessment holds true, it would have health implications as female students who participate in less MVPA per day are less likely to achieve the daily physical activity guidelines and reduce their activity throughout adolescence at a faster rate.⁵⁸

Although programme records showed that recess and lunch activities were offered in all intervention schools, girls were less likely to report that their school offered organised recess and lunchtime physical activities compared to boys; a substantial proportion of both sexes were not aware of the activities. As physical activity during recess and lunch has been reported to contribute as much as 40% towards daily physical activity recommendations,⁵⁹ the introduction of recess and lunchtime activities that are more evident to students, especially those that appeal to girls, seems an important consideration for future research. However, our mid-intervention results may also indicate girls take longer to respond to interventions than boys.

Strengths

The strengths of this study include the group randomised controlled design, use of an objective measure of physical activity, the focus on disadvantaged populations and the multicomponent socioecological design. However, there are limitations. Obtaining valid accelerometer data in this age group was challenging,⁶⁰ as has been discussed elsewhere.⁶¹

Although a high proportion of students who participated in baseline also participated at midpoint (84%), only 61% of the baseline sample provided at least three days of valid data at 12 months. This decrease, however, seems consistent with other studies for this target group.^{16 54} Lubans *et al*⁶² found that although 79% to 85% of the baseline sample was retained after 12 months, only 53.5% of the sample of disadvantaged girls provided three or more days of valid accelerometer data. Similarly, only 64% of students in the HEIA study in Norway provided useable accelerometer data at the 20-month postintervention.⁵⁴ Although accelerometers are considered the optimal method for measuring physical activity, compliance to protocols among the students, particularly disadvantaged students, has been documented as a challenge.^{13 63} The study did not assess

maturity status, which is known to impact on physical activity levels of adolescents.⁶⁴

In summary, the mid-intervention effects of PA4E1 demonstrates the potential to implement a multicomponent school-based intervention in disadvantaged secondary schools. We will report 24-month follow-up as that is the primary outcome of the study.

What are the new findings?

- ▶ School-based physical activity interventions targeting adolescents from disadvantaged schools are feasible and can produce meaningful physical activity effects.
- ▶ Interventions with positive mid-intervention effects are more likely to have significant effects at follow-up and therefore, the Physical Activity 4 Everyone intervention shows promising signs for impacting on physical activity levels of disadvantaged adolescents.
- ▶ At mid-intervention, the intervention appears to be impacting more on male students.

How might it impact on clinical practice in the near future?

- ▶ Multicomponent school-based interventions that include strategies across the domains of the socioecological framework, such as increasing physical activity level in physical education, recess and/or lunch activities and linking with parents can improve physical activity levels of disadvantaged adolescents.
- ▶ Intervention adoption strategies appear important particularly in this setting/population.

Contributors JW, PJM, DRL, LC, LW and KG obtained funding for the research. All authors contributed to developing the intervention and data collection protocols and materials, and reviewing, editing, and approving the final version of the paper. All authors accept full responsibility for, and have read and approved the final manuscript.

Funding This study is funded through the NSW Ministry of Health, Health Promotion Demonstration grant scheme. In kind support for the study is also provided by the Hunter New England Local Health District. The project also received infrastructure support from the Hunter Medical Research Institute (HMRI). The study has been approved by the Hunter New England Human Research Ethics Committee (Ref No. 11/03/16/4.05), University of Newcastle (Ref No. H-2011-0210), NSW Department of Education and Communities (SERAP 2011111), Maitland Newcastle Catholic School Diocese and Broken Bay Catholic School Diocese. Anthony Okely is supported by a National Heart Foundation Career Development Fellowship.

Competing interests None declared.

Ethics approval Hunter New England Area Human Research Ethics Committee; University of Newcastle Human Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Lee IM, Shiroma EJ, Lobelo F, *et al*. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet* 2012;380:219–29.
- 2 Hallal PC, Andersen LB, Bull FC, *et al*. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet* 2012;380:247–57.

- 3 Borraccino A, Lemma P, Iannotti RJ, *et al.* Socioeconomic effects on meeting physical activity guidelines: comparisons among 32 countries. *Med Sci Sports Exerc* 2009;41:749–56.
- 4 Drummond MDC, Dollman J, Abery L. Physical activity from early childhood to adolescence: a literature review of issues and interventions in disadvantaged populations. *J Student Wellbeing* 2010;4:17–31.
- 5 Telama R, Yang X, Viikari J, *et al.* Physical activity from childhood to adulthood: a 21-year tracking study. *Am J Prev Med* 2005;28:267–73.
- 6 Statistics ABo. Schools, Australia, 2013. Secondary Schools, Australia, 2013 2013. <http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4221.0Main+Features12010>
- 7 World Health Organization (WHO). Global strategy on diet, physical activity and health. Geneva: World Health Organization, 2004.
- 8 Centers for Disease Control and Prevention. *Comprehensive School Physical Activity Programs: A Guide for Schools*. Department of Health and Human Services. Atlanta, Georgia: USA, 2013.
- 9 Dobbins M, Husson H, DeCorby K, *et al.* School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. *Cochrane Database Syst Rev* 2013;2:CD007651.
- 10 Kriemler S, Meyer U, Martin E, *et al.* Effect of school-based interventions on physical activity and fitness in children and adolescents: a review of reviews and systematic update. *Br J Sports Med* 2011;45:923–30.
- 11 Murillo Pardo B, Garcia Bengoechea E, Generelo Lanaspá E, *et al.* Promising school-based strategies and intervention guidelines to increase physical activity of adolescents. *Health Educ Res* 2013;28:523–38.
- 12 van Sluijs EM, McMinn AM, Griffin SJ. Effectiveness of interventions to promote physical activity in children and adolescents: systematic review of controlled trials. *Br J Sports Med* 2008;42:653–7.
- 13 Cohen KE, Morgan PJ, Plotnikoff RC, *et al.* Physical activity and skills intervention: SCORES cluster randomized controlled trial. *Med Sci Sports Exerc* 2015;47:765–74.
- 14 McKay HA, Macdonald HM, Nettlefold L, *et al.* Action schools! BC implementation: from efficacy to effectiveness to scale-up. *Br J Sports Med* 2015;49:210–18.
- 15 van Sluijs EM, McMinn AM, Griffin SJ. Effectiveness of interventions to promote physical activity in children and adolescents: systematic review of controlled trials. *BMJ* 2007;335:703.
- 16 Dewar DL, Morgan PJ, Plotnikoff RC, *et al.* The nutrition and enjoyable activity for teen girls study: a cluster randomized controlled trial. *Am J Prev Med* 2013;45:313–17.
- 17 Casey MM, Harvey JT, Telford A, *et al.* Effectiveness of a school-community linked program on physical activity levels and health-related quality of life for adolescent girls. *BMC Public Health* 2014;14:649.
- 18 Smith JJ, Morgan PJ, Plotnikoff RC, *et al.* Smart-phone obesity prevention trial for adolescent boys in low-income communities: the ATLAS RCT. *Pediatrics* 2014;134:e723–31.
- 19 Wilson DK, Lee Van Horn M, Kitzman-Ulrich H, *et al.* Results of the “Active by Choice Today” (ACT) randomized trial for increasing physical activity in low-income and minority adolescents. *Health Psychol* 2011;30:463–71.
- 20 Lubans DR, Morgan PJ, Callister R, *et al.* Effects of integrating pedometers, parental materials, and E-mail support within an extracurricular school sport intervention. *J Adolesc Health* 2009;44:176–83.
- 21 Sutherland R, Campbell E, Lubans DR, *et al.* A cluster randomised trial of a school-based intervention to prevent decline in adolescent physical activity levels: study protocol for the ‘Physical Activity 4 Everyone’ trial. *BMC Public Health* 2013;13:57.
- 22 Australian Bureau of Statistics (ABS). Technical Paper:Census of Population and Housing: Socio-Economic Indexes For Australia (SEIFA). Cat. no 2039.0.55.001. Canberra; 2001.
- 23 Bandura A. *Social foundations of thought and action: a social cognitive theory*. Englewood Cliffs: Prentice-Hall, Inc, 1986.
- 24 Green LW, Richard L, Potvin L. Ecological foundations of health promotion. *Am J Health Promot* 1996;10:270–81.
- 25 Bandura A. Health promotion by social cognitive means. *Health Educ Behav* 2004;31:143–64.
- 26 Lubans D, Morgan P. Evaluation of an extra-curricular school sport programme promoting lifestyle and lifetime activity for adolescents. *J Sports Sci* 2008;26:519–29.
- 27 Baranowski T, Anderson C, Carmack C. Mediating variable framework in physical activity interventions. How are we doing? How might we do better? *Am J Prev Med* 1998;15:266–97.
- 28 WHO WHO. *Planning meeting in health promoting schools project: background, development and strategy outline of the health promoting schools project*. Copenhagen: WHO, 1991.
- 29 Lonsdale C, Rosenkranz RR, Peralta LR, *et al.* A systematic review and meta-analysis of interventions designed to increase moderate-to-vigorous physical activity in school physical education lessons. *Prev Med* 2013;56:152–61.
- 30 Lubans DR, Morgan PJ, Callister R, *et al.* Exploring the mechanisms of physical activity and dietary behavior change in the program x intervention for adolescents. *J Adolesc Health* 2010;47:83–91.
- 31 Trost SG. State of the art reviews: measurement of physical activity in children and adolescents. *Am J Lifestyle Med* 2007;1:299–314.
- 32 Dollman J, Okely AD, Hardy L, *et al.* A hitchhiker’s guide to assessing young people’s physical activity: deciding what method to use. *J Sci Med Sport* 2009;12:518–25.
- 33 McClain JJ, Tudor-Locke C. Objective monitoring of physical activity in children: considerations for instrument selection. *J Sci Med Sport* 2009;12:526–33.
- 34 Troiano RP, Berrigan D, Dodd KW, *et al.* Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008;40:181–8.
- 35 Yildirim M, Verloigne M, de Bourdeaudhuij I, *et al.* Study protocol of physical activity and sedentary behaviour measurement among schoolchildren by accelerometry—cross-sectional survey as part of the ENERGY-project. *BMC Public Health* 2011;11:182.
- 36 Evenson KR, Catellier DJ, Gill K, *et al.* Calibration of two objective measures of physical activity for children. *J Sports Sci* 2008;26:1557–65.
- 37 Marfell-Jones M, Olds T, Stew A, *et al.* *International standards for anthropometric assessment*. Australia: The International Society for the Advancement of Kinanthropometry, 2006.
- 38 Cole TJ, Bellizzi MC, Flegal KM, *et al.* Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240–3.
- 39 Drenowatz C, Eisenmann JC, Pfeiffer KA, *et al.* Influence of socio-economic status on habitual physical activity and sedentary behavior in 8- to 11-year old children. *BMC Public Health* 2010;10:214.
- 40 Gortmaker SL, Lee RM, Mozaffarian RS, *et al.* Effect of an after-school intervention on increases in children’s physical activity. *Med Sci Sports Exerc* 2012;44:450–7.
- 41 Audrey S, Bell S, Hughes R, *et al.* Adolescent perspectives on wearing accelerometers to measure physical activity in population-based trials. *Eur J Public Health* 2013;23:475–80.
- 42 Lubans DR, Morgan PJ, Dewar D, *et al.* The Nutrition and Enjoyable Activity for Teen Girls (NEAT girls) randomized controlled trial for adolescent girls from disadvantaged secondary schools: rationale, study protocol, and baseline results. *BMC Public Health* 2010;10:652.
- 43 Murray DM, Stevens J, Hannan PJ, *et al.* School-level intraclass correlation for physical activity in sixth grade girls. *Med Sci Sports Exerc* 2006;38:926–36.
- 44 Mallinckrodt CH, Kaiser CJ, Watkin JG, *et al.* The effect of correlation structure on treatment contrasts estimated from incomplete clinical trial data with likelihood-based repeated measures compared with last observation carried forward ANOVA. *Clin Trials* 2004;1:477–89.
- 45 Yildirim M, van Stralen MM, Chinapaw MJ, *et al.* For whom and under what circumstances do school-based energy balance behavior interventions work? Systematic review on moderators. *Int J Pediatr Obes* 2011;6:e46–57.
- 46 Assmann SF, Pocock SJ, Enos LE, *et al.* Subgroup analysis and other (mis)uses of baseline data in clinical trials. *Lancet* 2000;355:1064–9.
- 47 Holman RM, Carson V, Janssen I. Does the fractionalization of daily physical activity (sporadic vs. bouts) impact cardiometabolic risk factors in children and youth? *PLoS ONE* 2011;6:e25733.
- 48 Logan GM, Harris N, Duncan S, *et al.* A review of adolescent high-intensity interval training. *Sports Med* 2014;44:1071–85.
- 49 Thomas AS, Greene LF, Ard JD, *et al.* Physical activity may facilitate diabetes prevention in adolescents. *Diabetes Care* 2009;32:9–13.
- 50 Health. TDo. Australia’s Physical Activity and Sedentary Behaviour Guidelines. Secondary Australia’s Physical Activity and Sedentary Behaviour Guidelines [Webpage] 14th February 2014 2014. <http://www.health.gov.au/internet/main/publishing.nsf/content/health-pubhlth-strateg-phys-act-guidelines>
- 51 Metcalf B, Henley W, Wilkin T. Effectiveness of intervention on physical activity of children: systematic review and meta-analysis of controlled trials with objectively measured outcomes (EarlyBird 54). *BMJ* 2012;345:e5888.
- 52 Haerens L, Deforche B, Maes L, *et al.* Evaluation of a 2-year physical activity and healthy eating intervention in middle school children. *Health Educ Res* 2006;21:911–21.
- 53 Webber LS, Catellier DJ, Lytle LA, *et al.* Promoting physical activity in middle school girls: trial of activity for adolescent girls. *Am J Prev Med* 2008;34:173–84.
- 54 Grydeland M, Bergh IH, Bjelland M, *et al.* Intervention effects on physical activity: the HEIA study—a cluster randomized controlled trial. *Int J Behav Nutr Phys Act* 2013;10:17.
- 55 de Villiers A, Steyn NP, Draper CE, *et al.* “HealthKick”: formative assessment of the health environment in low-resource primary schools in the Western Cape Province of South Africa. *BMC Public Health* 2012;12:794.
- 56 Graham DJ, Lucas-Thompson RG, O’Donnell MB. Jump in! An investigation of school physical activity climate, and a pilot study assessing the acceptability and feasibility of a novel tool to increase activity during learning. *Front Public Health* 2014;2:58.
- 57 Fagen MC, Asada Y, Welch S, *et al.* Policy, systems, and environmentally oriented school-based obesity prevention: opportunities and challenges. *J Prev Interv Community* 2014;42:95–111.
- 58 Camacho-Minano MJ, LaVoie NM, Barr-Anderson DJ. Interventions to promote physical activity among young and adolescent girls: a systematic review. *Health Educ Res* 2011;26:1025–49.

- 59 Parrish AM, Okely AD, Stanley RM, *et al.* The effect of school recess interventions on physical activity: a systematic review. *Sports Med* 2013;43: 287–99.
- 60 Troiano RP, McClain JJ, Brychta RJ, *et al.* Evolution of accelerometer methods for physical activity research. *Br J Sports Med* 2014;48:1019–23.
- 61 Pedisic Z, Bauman A. Accelerometer-based measures in physical activity surveillance: current practices and issues. *Br J Sports Med* 2015;49: 219–23.
- 62 Lubans DR, Morgan PJ, Okely AD, *et al.* Preventing obesity among adolescent girls: one-year outcomes of the nutrition and enjoyable activity for teen girls (NEAT Girls) cluster randomized controlled trial. *Arch Pediatr Adolesc Med* 2012;166:821–7.
- 63 Rich C, Cortina-Borja M, Dezateux C, *et al.* Predictors of non-response in a UK-wide cohort study of children's accelerometer-determined physical activity using postal methods. *BMJ open* 2013;3:e002290.
- 64 Metcalf BS, Hosking J, Jeffery AN, *et al.* Exploring the adolescent fall in physical activity: a 10-yr Cohort Study (EarlyBird 41). *Med Sci Sports Exerc* 2015.