What are the critical elements of sideline screening that can be used to establish the diagnosis of concussion? A systematic review

Jon Patricios,^{1,2} Gordon Ward Fuller,³ Richard Ellenbogen,⁴ Stanley Herring,^{4,5,6} Jeffrey S Kutcher,⁷ Mike Loosemore,⁸ Michael Makdissi,^{9,10} Michael McCrea,¹¹ Margot Putukian,¹² Kathryn J Schneider¹³

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2016-097441>)

For numbered affiliations see end of article.

Correspondence to

Dr Jon Patricios, Morningside Sports Medicine, PO Box 1267, Parklands, Cape Town 2121, South Africa; jpat@mwweb.co.za

Accepted 6 February 2017
Published Online First
7 March 2017

ABSTRACT

Background Sideline detection is the first and most significant step in recognising a potential concussion and removing an athlete from harm. This systematic review aims to evaluate the critical elements aiding sideline recognition of potential concussions including screening tools, technologies and integrated assessment protocols. **Data sources** Bibliographic databases, grey literature repositories and relevant websites were searched from 1 January 2000 to 30 September 2016. A total of 3562 articles were identified.

Study selection Original research studies evaluating a sideline tool, technology or protocol for sports-related concussion were eligible, of which 27 studies were included.

Data extraction A standardised form was used to record information. The QUADAS-2 and Newcastle-Ottawa tools were used to rate risk of bias. Strength of evidence was assessed using the Grades of Recommendation, Assessment, Development and Evaluation Working Group system.

Data synthesis Studies assessing symptoms, the King-Devick test and multimodal assessments reported high sensitivity and specificity. Evaluations of balance and cognitive tests described lower sensitivity but higher specificity. However, these studies were at high risk of bias and the overall strength of evidence examining sideline screening tools was very low. A strong body of evidence demonstrated that head impact sensors did not provide useful sideline concussion information. Low-strength evidence suggested a multimodal, multitime-based concussion evaluation process incorporating video review was important in the recognition of significant head impact events and delayed onset concussion.

Conclusion In the absence of definitive evidence confirming the diagnostic accuracy of sideline screening tests, consensus-derived multimodal assessment tools, such as the Sports Concussion Assessment Tool, are recommended. Sideline video review may improve recognition and removal from play of athletes who have sustained significant head impact events. Current evidence does not support the use of impact sensor systems for real-time concussion identification.

may increase the likelihood of incurring a subsequent head or musculoskeletal injury,³ and repeated concussions could be associated with long-term consequences such as persistent postconcussive symptoms, depression or neurodegenerative disorders.^{1 4 5} Early detection of suspected concussion and removal of the affected player will help prevent these potential adverse sequelae and facilitate further evaluation, management and safe return to play. This systematic review aims to evaluate the critical elements aiding off-field (commonly termed 'sideline') recognition of potential concussions. Specific objectives were to assess the diagnostic accuracy of existing clinical screening and diagnostic tools, determine the utility of technology in detecting SRC and assess integrated head injury assessment protocols currently used in professional collision sports.

METHODS

Study design

Expert consensus guidelines for the conduct of systematic reviews were followed,^{6–8} and a detailed protocol stating an a priori analysis plan was registered before data collection (PROSPERO 2016:CRD42016037831). The review question and inclusion/exclusion criteria are detailed in [table 1](#). Online supplementary details on methodology, including a glossary of technical terms, are presented in the online supplementary file web appendix.

Identification of evidence

An extensive range of electronic information sources were examined including all major bibliographic databases, specialist sports medicine databases, grey literature repositories and relevant websites (see online supplementary web appendix for details). Additional information sources included forward and backward citation searching, author searching, reference checking and contact with experts. Search strategies for bibliographic databases were developed iteratively in conjunction with an information services specialist (University College London) and underwent external peer review (University of Sheffield).

Searches were conducted for original research published between 2000 (corresponding to the modern definition of concussion) and week 4, April 2016, and were otherwise unrestricted. Current awareness searches were conducted in MEDLINE

CrossMark

To cite: Patricios J, Fuller GW, Ellenbogen R, et al. *Br J Sports Med* 2017;**51**:888–895.

INTRODUCTION

Despite a consensus definition of sports-related concussion (SRC) having been well elucidated,¹ its immediate and accurate recognition in a clinical setting remains a challenge.² Sustaining a SRC

Table 1 Review question and inclusion criteria

Primary review question/aim	
What are the critical elements of sideline screening that can be used to establish the diagnosis of concussion or suspected concussion?	
Inclusion criteria	
Population	Athletes competing in sporting activity and sustaining a non-trivial head impact event (includes any nationality, gender, age group or level of performance).
Intervention/index tests	Any sideline* screening assessment used to detect suspected concussion following sports-related significant head impact events (including historical features, symptoms, physical findings, clinical tests or technologies)
Outcome/reference standard	Concussion, clinically diagnosed by a registered medical practitioner.
Study design	Published or unpublished studies of any research design.
Exclusion criteria	
Population	Not related to sport, subjects <13 years, animal studies
Intervention/index tests	Non-sideline testing
Outcome/reference standard	Concussion not examined
Study design	Case reports
Review subtopics/objectives	
Sideline screening tests	Characterise the diagnostic accuracy of sideline clinical tests to detect suspected SRC, including: <ul style="list-style-type: none"> ▶ Symptoms and clinical signs ▶ Balance tests ▶ Oculomotor tests ▶ Cognitive tests ▶ Multimodal assessments (either joint use of individual sideline tests or multifaceted instruments)
Technology	Determine the utility of technology in the detection of suspected SRC.
Integrated head injury assessment protocols	Evaluate integrated protocols for the detection and management of SRC currently used in professional collision sports.

*'Sideline' is used generally to denote testing away from the immediate sporting environment, for example, rink side, track side, locker room, medical room, touch line and so on.
SRC, sports-related concussion.

and Embase (week 4, September 2016) immediately prior to submission.

Selection of evidence and data extraction

Original research studies identified during searches were assessed in a four-stage process by teams of two independent reviewers. First, titles and abstracts were screened for relevance. Second, full-text articles were examined as required to assess eligibility. Third, studies meeting review inclusion criteria were classified into domains pertaining to: sideline screening tests (comprising subtopics of clinical signs and symptoms, balance tests, oculomotor assessments, cognitive tests and multimodal testing strategies); technology; and professional sports-specific head injury assessment protocols (defined in table 1). Finally, data extraction was performed separately for eligible studies within each subtopic. A single unblinded reviewer extracted information on study characteristics, methodology and results using a standardised data extraction form; and a second reviewer

independently checked data for consistency and accuracy. In cases of disagreement at any stage, consultation with a third author was planned, with consensus derived by arbitration.

Risk of bias assessment

Included studies were assessed for risk of bias using peer-reviewed critical appraisal checklists appropriate to study design. The QUDAS-2 tool was used for diagnostic accuracy studies.⁹ Observational studies were evaluated using the Newcastle-Ottawa scale.⁶ A single unblinded reviewer within each subgroup team assessed risk of bias, with a second reviewer independently checking the assessment for validity. Any disagreement between reviewers was resolved by consensus and consultation with a third author with expertise in epidemiology and critical appraisal.

Data synthesis, statistical analyses and assessment of overall quality of evidence

Data synthesis and statistical analysis were performed separately for eligible studies within each subtopic. Results are presented descriptively with reported point estimates and 95% CIs and summarised graphically using Forest plots.¹⁰ Heterogeneity was assessed using the I^2 statistic.¹¹ A narrative synthesis was prespecified in the event that clinically and methodologically homogenous studies at low risk of bias were not identified. References were managed in EndNote (Clarivate Analytics, Berkeley California, USA), extracted data were collated in Excel 2013, and Forest plots were formulated using Meta-DiSc V.1.4 (University of Birmingham, Birmingham, UK). The overall quality of evidence for each outcome was assessed using the consensus Grades of Recommendation, Assessment, Development and Evaluation Working Group (GRADE) approach.¹² GRADE is a systematic method of assessing quality of evidence and strength of recommendations taking into account methodological flaws, consistency of results, generalisability of findings and the effectiveness of treatments. A clinical diagnosis of concussion was the primary outcome for each domain.

RESULTS

Study selection

A total of 3562 citations were screened for eligibility, with the full text of 198 articles retrieved for detailed evaluation. During full text examination, 27 studies were found meeting review inclusion criteria: sideline screening assessment (21 studies); integrated diagnostic protocols (1 study) and technology (5 studies). Figure 1 describes the selection of studies in detail.

Sideline screening tests

Characteristics of included studies

Twenty-one studies met review inclusion criteria and reported interpretable data on the diagnostic accuracy of screening tests, either alone or in combination, to identify suspected SRC. Characteristics of the included studies examining sideline assessments are summarised in table 2.

Risk of bias

Assessment of risk of bias is summarised according to QUDAS-2 domains in table 3 and figure 2. Overall risk of bias was high or unclear for all included studies. The predominant limitation was the use of a 'two-gate' study design using healthy controls, which is known to overestimate estimates of test performance.^{13 14} Other systematic errors included delayed index testing, inaccurate reference standard assessment by a

non-medically trained outcome assessors, and test and diagnostic review, incorporation and attrition biases. Detailed risks of bias evaluations are presented in the online supplementary web appendix.

Results

The diagnostic accuracy of sideline assessments for detecting suspected concussion is summarised in figure 3. Studies examining symptoms, the King-Devick (KD) test and multimodal assessments reported relatively good sensitivity and specificity. Evaluations of balance and cognitive tests described lower sensitivity, but relatively good specificity. However, results were imprecise and heterogeneous for all types of sideline assessments, in addition to the concerns regarding the internal validity. The overall quality of evidence according to GRADE criteria was very low for all classes of sideline tests based on serious concerns regarding inconsistency, imprecision and risk of bias. Detailed results and evaluation of overall quality of evidence for individual tests are provided in the online supplementary web appendix.

Technology

Five studies met review inclusion criteria and reported interpretable data on the use of a technology in sideline screening for SRC, examining head impact sensors (four studies) and sideline video review (one study).^{15–19} Overall risk of bias was low for all studies. Reported results indicated that no clinically significant relationship existed between impact magnitude, or location, and concussion ($p > 0.05$). Fuller *et al*¹⁶ reported that sideline video review contributed to identification of 61.5% of significant

head impact events and influenced sideline evaluation in 20.4% of cases. The overall GRADE quality of evidence was rated as high for head impact sensors and low for sideline video review. Table 4 summarises the characteristics, risk of bias and main results of included technology studies. Further details on risk of bias and GRADE ratings are provided in the online supplementary web appendix.

Integrated head injury assessment protocols

No experimental or comparative effectiveness research was identified evaluating the performance of alternative head injury assessment protocols. However, a single study at low risk of bias was retrieved which evaluated a comprehensive system used at the elite level in Rugby Union.¹⁶ The major finding was the importance of a multimodal, multitime-based concussion evaluation process incorporating video review to identify significant head impact events and delayed onset concussion. The overall GRADE quality of evidence was rated low, secondary to imprecision and potential inconsistency. Further details on existing integrated head injury assessment protocols in professional sports, and the characteristics of Fuller *et al* are provided in the online supplementary web appendix.

DISCUSSION

Summary of key findings

Studies examining symptoms, the KD test and multimodal assessments reported high sensitivity and specificity. Evaluations of balance and cognitive tests described lower sensitivity, but good specificity. However, the overall strength of evidence examining sideline screening tools was of very low quality secondary to high risk of bias, and imprecise and heterogeneous diagnostic accuracy estimates. Studies examining technology provided a high (head impact sensors) or low (video review) strength body of evidence. Head impact sensors did not provide useful information. Conversely, a multimodal, multitime-based concussion evaluation process incorporating video review appeared to be important for the identification of significant head impact events and the delayed onset concussion.

Interpretation

A meta-analysis was not performed due to the absence of studies at low risk of bias and marked heterogeneity; in accordance with the prespecified analysis plan, a narrative synthesis was therefore conducted. Interestingly, no obvious patterns were evident between study results and design characteristics including sample size, setting, performance level, sport or risk of bias. This may be due to the inherent generalisability of findings, but could also be explained by biases operating in different directions and to varying magnitudes across different studies.

Notwithstanding the high risk of systematic error, a wide range of settings, sports and age groups were investigated in eligible studies suggesting good external validity of findings. However, in addition to information on diagnostic accuracy, the feasibility, cost and acceptability of alternative sideline tests may be important in applying these results to different settings. The availability of baseline data, testing environment and influence of the athlete-physician relationship could also affect generalisability. Importantly, in lower levels of competition where medical staff may be limited, an alternative 'recognise and remove' approach is recommended, with exclusion of the sideline screening stage, and immediate and permanent removal from any further participation when there is any suspicion of concussion.^{1 20}

Figure 1 Flow of identification, screening, eligibility and inclusion criteria for the literature review of sideline diagnosis of concussion.

Table 2 Characteristics of included studies examining sideline screening assessments

Study	Setting	Study design	Sample size (n)	Sport(s)	Level	Mean age (years±SE)	Gender (% male)	Index test(s)	Reference standard
Maddocks <i>et al</i> ³³	Australia	PCS	56	Australian football	Professional	NR	100	Individual symptoms, Maddocks questions	Clinical diagnosis
McCrary <i>et al</i> ³⁴	Australia	PCS	303	Australian football	Professional	NR	100	Individual symptoms	Clinical diagnosis
Barr and McCrea ³⁵	USA	PCS	118	American football	Varsity high school	18.1 (NR)	NR	SAC	Clinical diagnosis
Erlanger <i>et al</i> ³⁶	USA	PCS	47	American football, ice hockey, field hockey, wrestling, soccer, basketball	School adolescents	17.6 (SD 2.23)	57	Individual symptoms	NR
McCrea ³⁷	USA	PCS	118	American football	Varsity high school	19.8±1.3	NR	SAC	Clinical diagnosis
McCrea <i>et al</i> ³⁸	USA	PCS	91	American football	Varsity high school	17.5±2.1	NR	SAC	Clinical diagnosis
McCrea <i>et al</i> ³⁹	USA	PCS	150	American football	Collegiate adults	20.04 (SD 1.36)	100	GSC, BESS, SAC	Clinical diagnosis
Echlin <i>et al</i> ⁴⁰	USA	PCS	67	Ice hockey	Junior adolescents	18.2±1.2	100%	BESS, SAC	Clinical diagnosis + SCAT 2
Galetta <i>et al</i> ⁴¹	USA	PCS	39	Boxing, mixed martial arts	Amateur adult	24	97	KD	MACE
Galetta <i>et al</i> ⁴²	USA	PCS	219	American football, soccer, basketball	Collegiate athletics	20.3±1.4	83	KD	Clinical diagnosis
Barr <i>et al</i> ⁴³	USA	PCS	90	American football	High school, collegiate	NR	100	CSI, SAC, BESS	Clinical diagnosis
King <i>et al</i> ⁴⁴	NZ	PCS	50	Rugby league	Amateur adult	22.4±4.1	100	KD	SCAT 2
Galetta <i>et al</i> ⁴⁵	USA	PCS	27	Ice hockey	Professional	25±5	100	KD, SAC	SCAT 2
Dhawan <i>et al</i> ⁴⁶	USA	PCS	141	Hockey	High school athletics	NR	NR	KD	NR
Leong <i>et al</i> ⁴⁷	USA	PCS	34	Boxing	Amateur adult	25.8±8.3	85	KD	MACE
Fuller <i>et al</i> ²¹	UK, RSA, France	PCS	165	Rugby union	Professional adults	NR	100	PSACA1 tool: Maddocks questions, symptoms checklist, mental status assessment, tandem stance test	Clinical diagnosis + SCAT 3
Galetta <i>et al</i> ⁴⁸	USA	PCS	243	Ice hockey, lacrosse, athletics	Amateur youth, collegiate athletics	Youths: 11±3, adults: 20±1	Youths: 84, Adults: 74	KD, timed tandem gait, SAC	Clinical diagnosis
Leong <i>et al</i> ⁴⁹	USA	PCS	127	American football, basketball	Collegiate athletics	19.6±1.2	94	KD	Modified SCAT 2
Marinides <i>et al</i> ⁵⁰	USA	RCS	217	American football, lacrosse, soccer	Collegiate athletics	NR	70	KD, BESS, SAC	Clinical diagnosis
Putukian <i>et al</i> ⁵¹	USA	PCS	263	American football, rugby union, sprint football, crew	Collegiate adults	20.33 (SD 1.74)	67%	SCAT2 symptom checklist, modified BESS, SAC, SCAT2	Clinical diagnosis
Seidman <i>et al</i> ⁵²	USA	PCS	337	American football	High school athletics	15.4±1.3	100	KD	SCAT 3

BESS, Balance Error Scoring System; CSI, Concussion Symptom Inventory; GSC, Graded Symptom Checklist; KD, King-Devick; MACE, Military Acute Concussion Evaluation; NR, not reported; NZ, New Zealand; PCS, prospective cohort study; RCS, retrospective cohort study; RSA, Republic of South Africa; SAC, Standardised Assessment of Concussion; SCAT, Sports Concussion Assessment Tool.

A key concept in sideline assessment is the rapid screening for a suspected concussion, rather than the definitive diagnosis of a head injury. Players manifesting clear on-field observable signs, such as loss of consciousness, ataxia, tonic posturing or

post-traumatic seizures, can immediately be diagnosed with a concussion and removed from sporting participation. Athletes with the possibility of suspected concussion following a significant head impact event can alternatively proceed to sideline

Table 3 Summary of risk of bias across included sideline screening studies

Study	Risk of bias				Overall
	Patient selection	Index test	Reference standard	Flow and timing	
Maddocks <i>et al</i> ³³	H	L	L	L	H
Barr and McCrea ³⁵	H	?	H	L	H
McCroory <i>et al</i> ³⁴	H	?	L	L	H
McCrea ³⁷	H	?	H	L	H
McCrea <i>et al</i> ³⁸	H	?	H	L	H
Erlanger <i>et al</i> ³⁶	H	?	?	L	H
McCrea <i>et al</i> ³⁹	H	H	H	H	H
Echlin <i>et al</i> ⁴⁰	H	H	H	H	H
Galetta <i>et al</i> ⁴¹	L	?	?	H	H
Galetta <i>et al</i> ⁴²	H	L	H	L	H
Barr <i>et al</i> ⁴³	H	H	H	L	H
King <i>et al</i> ⁴⁴	L	?	?	?	?
Galetta <i>et al</i> ⁴⁵	H	?	?	L	H
Dhawan <i>et al</i> ⁴⁶	H	?	?	L	H
Fuller <i>et al</i> ²¹	L	L	H	L	H
Leong <i>et al</i> ⁴⁷	L	L	L	H	H
Galetta <i>et al</i> ⁴⁸	H	?	?	H	H
Leong <i>et al</i> ⁴⁹	H	L	H	L	H
Marinides <i>et al</i> ⁵⁰	H	?	?	H	H
Putukian <i>et al</i> ⁵¹	H	H	L	H	H
Seidman <i>et al</i> ⁵²	H	H	L	H	H

H, high; L, low; ?, unclear risk of bias.

screening, with a later definitive diagnostic evaluation. Clearly, to allow sufficient time and a suitable environment for testing, this should occur away from the sporting environment, and may necessitate a temporary athlete interchange. The importance of off-field testing is exemplified by findings in professional Rugby where the number of players with confirmed concussion returning to play following their head injury dropped from 56% to 13% following the introduction of the Pitchside Suspected Concussion Assessment that superseded an 'on-the-field-and-on-the-run' approach.²¹

Elite contact and collision sports are played at a fast pace in a disorganised environment, where the view of medical staff may be obscured, challenging the evaluation of head impact events. Video review appeared to be helpful in identifying both observable signs of concussion and cases of possible suspected concussion where further assessment off-field is beneficial. Furthermore, evolving and delayed onset concussions have been well described,^{16,22} highlighting the importance of careful follow-up after a significant head impact, regardless of a negative sideline screening test or early diagnostic evaluation. Consequently, implementation of systematic head injury assessment protocols appears to improve detection and management of the full spectrum of SRC.

Concussion can manifest as a diverse range of somatic, cognitive, behavioural or emotional symptoms; and/or physical signs such as vestibulo-ocular deficits, loss of consciousness and ataxia.¹ It would therefore be expected *ex ante* that multimodal assessments, evaluating several of these domains, are necessary to maximise detection of different subtypes of SRC. However, with simultaneous testing a net gain in sensitivity usually occurs at the expense of a net loss in specificity.²³ Interestingly, included multimodal assessment studies reported both high sensitivity and specificity which could suggest either an optimal combination of

Figure 2 Summary of risk of bias in included studies examining sideline screening.

tests, or could reflect study biases. Given the absence of definitive evidence on the performance of sideline tests, expert consensus opinion is necessary to guide practice and strongly recommends the use of a multimodal assessment tool, of which the Sports Concussion Assessment Tool (SCAT; now in its 4th version) is the most established, well developed and studied.²⁴

It is important to note that the pretest probability of concussion will strongly influence the performance of sideline screening tests.²⁵ In settings with high prevalence of concussion, or high test thresholds, the negative predictive value of sideline tests will fall. High sensitivity and specificity would consequently be necessary to ensure the detection of a satisfactory proportion of cases. Conversely, indiscriminate testing, with a lower pretest probability of concussion, would result in higher negative predictive values, but worsening numbers of false positives. Such a safety first approach might be preferred in non-elite settings.

Consistency with other studies or reviews

There have been a large number of narrative reviews, position statements and editorials that have previously examined the role of sideline screening tests or technology in the detection of SRC. Although these articles are inherently limited by a lack of defined inclusion criteria, systematic search strategies and transparent risk of bias assessment, their conclusions are broadly consistent with the current systematic review. For example, Eckner *et al*²⁶ stated that 'multiple assessment tools are available, with no single tool showing clear superiority. Many tools remain based more on expert opinion than rigorous scientific evaluation.'

Six related systematic reviews were also identified during the literature searches, comprising examination of symptom checklists,²⁷ SCAT 2/3,²⁸ Balance Error Scoring System,²⁹ KD test^{30,31} and sideline testing in general.³² Although the review questions were not directly comparable, including delayed non-sideline testing and additional examination of test reliability, similar studies were often included and conclusions concurred with the current study in Alla,²⁷ Yengo-Khan,²⁸ Bell²⁹ and Hunt.³⁰ For example, Alla²⁷ noted that 'There is little information available on the derivation or psychometric properties (eg, sensitivity, reliability, etc) of the various symptom scales', and Yengo-Khan²⁸ observed that 'the sensitivity and specificity of the

Figure 3 Diagnostic accuracy of sideline screening tests for suspected sports-related concussion.

SAC has been reported sparsely. Conversely, in contrast to the current findings, Galetta³¹ and King³² concluded that the KD test can successfully identify SRC on the sideline. This divergent opinion is explained by the absence of any risk of bias

assessment for constituent studies included in their reviews, resulting in the KD test being interpreted as having high sensitivity and specificity whereas the quality of evidence presented did not justify this.

Table 4 Characteristics, risk of bias and primary findings of included technology studies

Study	Setting	Design	Sample size (n)	Sport(s)	Level	Mean age (years±SE)	Technology	Risk of bias/ evidence level	Applicability concerns	Primary finding(s)
Guskiewicz <i>et al</i> ¹⁸	USA	PCS	81		high school	20.2±1.8	HITS	Low	Low	61.5% sensitivity for concussion*
Mihalak <i>et al</i> ¹⁹	USA	PCS	102	American football	Collegiate	19.6±1.6	HITS	Unclear	Low	PPV of 0.35% for concussion†
Greenwald <i>et al</i> ¹⁷	USA	PCS	449	American football	High school	NR	HITS	Low	Low	PPV of 0.3% for concussion*
Broglio <i>et al</i> ¹⁵	USA	PCS	78	American football	High school	16.7±0.8	HITS	Low	Low	PPV of 13.4% for concussion‡
Fuller <i>et al</i> ¹⁶	UK	PCS	49	Rugby union	Professional	26.5 (SD 3.5)	Sideline video review	Low	Low	Contributed to identification of 61% of significant head impact events

*Head impact threshold: linear acceleration >98.9 g.

†Threshold: linear acceleration >80g.

‡Threshold: >5582.3 rad/s² ± 96.1 g linear acceleration ± front/side/top impact.

HITS, head impact telemetry system; NR, not reported; PCS, prospective cohort study; PPV, positive predictive value.

Implications for research

There is an absence of valid research confirming the diagnostic accuracy and impact on improving outcomes of currently used sideline screening tests. Adequately powered diagnostic accuracy studies are therefore recommended that enrol a representative sample of athletes with suspected concussion following non-trivial head impact events. Ideally, once the diagnostic accuracy and optimal threshold of sideline tests have been confirmed, comparative effectiveness studies would investigate whether important outcomes are improved. Further research is also recommended to investigate the impact of integrated head injury assessment protocols and sideline video review for the evaluation of head impact events. Further research could usefully examine novel sideline screening tests such as reaction times, or investigate the utility of tablet software applications as an adjunct to sideline concussion screening.

Review limitations

There are a number of potential methodological weaknesses which could limit the validity of this systematic review. Because of time constraints, hand searching of journals and conference proceedings was not performed and regional bibliographic databases were not included raising the potential for publication bias. Decisions on study relevance, information gathering and validity were unblinded and potentially could have been influenced by preformed opinions. Furthermore, data extraction and risk of bias assessment were not performed in duplicate (ie, two truly independent reviews), with the second reviewer checking the assessment of the first reviewer. Finally, assessment of reference standard bias was challenged by the lack of a convincing diagnostic gold standard.

CONCLUSIONS

Based on this systematic review of the literature, an evidence-based recommendation for any individual screening test or protocol is not possible. The recognition of suspected concussion is therefore best approached using multimodal testing guided via expert consensus. The SCAT currently represents the most well-established and rigorously developed instrument available for sideline assessment. The addition of video review could potentially offer a promising approach to improve identification and evaluation of significant head impact events, and a multitime-based concussion evaluation process appears to

be important to detect delayed onset SRC. The KD test shows promise as a sideline screening test but requires adequately powered diagnostic accuracy studies which avoid a two-gate design with healthy controls, and enrolls a representative sample of athletes with suspected concussion. Collaboration between sporting codes to rationalise multimodal diagnostic sideline protocols may help facilitate more efficient application and monitoring. Current evidence does not support the use of impact sensor systems for real-time concussion screening.

Author affiliations

¹Section Sports Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

²Department of Emergency Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

³Centre for Urgent and Emergency Care Research, School of Health and Related Research, University of Sheffield, Sheffield, UK

⁴Department of Neurological Surgery, University of Washington, Seattle, Washington, USA

⁵Departments of Rehabilitation Medicine, University of Washington, Seattle, Washington, USA

⁶Orthopedics and Sports Medicine, University of Washington, Seattle, Washington, USA

⁷The Sports Neurology Clinic at the CORE Institute, Brighton, Michigan, USA

⁸Institute of Sport Exercise and Health, University College London, London, UK

⁹Florey Institute of Neuroscience and Mental Health, Austin Campus, Melbourne Brain Centre, Heidelberg, Victoria, Australia

¹⁰Olympic Park Sports Medicine Centre, Melbourne, Victoria, Australia

¹¹Department of Neurosurgery, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

¹²Director of Athletic Medicine, Princeton University, Princeton, New Jersey, USA

¹³Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Alberta, Canada

Competing interests JP received travel subsidies for conferences from South African Rugby and World Rugby. GWF is funded by the National Institute for Health Research and received travel funding from World Rugby. RE, SH, JSK, ML, MM, MP have no competing interests to declare. MM received travel and accommodation costs. KJS has received speaking honoraria for presentations at scientific meetings.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- McCrory P, Meeuwisse WH, Aubry M, *et al*. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Br J Sports Med* 2013;47:250–8.
- Helmy A, Agarwal M, Hutchinson PJ. Concussion and sport. *BMJ* 2013;347:f5748.

- 3 Herman D. C, Zaremski JL, Vincent HK, *et al.* Effect of neurocognition and concussion on musculoskeletal injury risk. *Curr Sports Med Rep* 2015;14:194–9.
- 4 Guskiewicz KM, Marshall SW, Bailes J, *et al.* Association between recurrent concussion and late-life cognitive impairment in retired professional football players. *Neurosurgery* 2005;57:719–26.
- 5 McKee AC, Cantu RC, Nowinski CJ, *et al.* Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury. *J Neuropathol Exp Neurol* 2009;68:709–35.
- 6 Higgins J, Green SP. *Cochrane handbook for systematic reviews of interventions*. Wiley-Blackwell: Oxford, 2008.
- 7 Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
- 8 Leeflang MM, Deeks JJ, Gatsonis C, *et al.* Cochrane Diagnostic Test Accuracy Working Group. Systematic reviews of diagnostic test accuracy. *Ann Intern Med* 2008;149:889–97.
- 9 Whiting PF, Rutjes AW, Westwood ME, *et al.* QUADAS-2 Group. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011;155:529–36.
- 10 Lewis S, Clarke M. Forest plots: trying to see the wood and the trees. *BMJ* 2001;322:1479–80.
- 11 Higgins JP, Thompson SG, Deeks JJ, *et al.* Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- 12 Schünemann HJ, Schünemann AH, Oxman AD, *et al.* GRADE Working Group. Grading quality of evidence and strength of recommendations for diagnostic tests and strategies. *Bmj* 2008;336:1106–10.
- 13 Rutjes AW, Reitsma JB, Vandenbroucke JP, *et al.* Case-control and two-gate designs in diagnostic accuracy studies. *Clin Chem* 2005;51:1335–41.
- 14 Lijmer JG, Mol BW, Heisterkamp S, *et al.* Empirical evidence of design-related bias in studies of diagnostic tests. *JAMA* 1999;282:1061–6.
- 15 Broglio SP, Schnebel B, Sosnoff JJ, *et al.* Biomechanical properties of concussions in high school football. *Med Sci Sports Exerc* 2010;42:2064–71.
- 16 Fuller CW, Fuller GW, Kemp SP, *et al.* Evaluation of World Rugby's concussion management process: results from Rugby World Cup 2015. *Br J Sports Med* 2017;51:64–9.
- 17 Greenwald RM, Gwin JT, Chu JJ, *et al.* Head impact severity measures for evaluating mild traumatic brain injury risk exposure. *Neurosurgery* 2008;62:789–98.
- 18 Guskiewicz KM, Mihalik JP, Shankar V, *et al.* Measurement of head impacts in collegiate football players: relationship between head impact biomechanics and acute clinical outcome after concussion. *Neurosurgery* 2007;61:1244–52.
- 19 Mihalik JP, Bell DR, Marshall SW, *et al.* Measurement of head impacts in collegiate football players: an investigation of positional and event-type differences. *Neurosurgery* 2007;61:1229–35.
- 20 Hodgson L, Patricios J. Clarifying concussion in youth rugby: recognise and remove. *Br J Sports Med* 2015;49:966–7.
- 21 Fuller GW, Kemp SP, Decq P. The International Rugby Board (IRB) Pitch side concussion assessment trial: a pilot test accuracy study. *Br J Sports Med* 2015;49:529–35.
- 22 Scorza KA, Raleigh MF, O'Connor FG. Current concepts in concussion: evaluation and management. *Am Fam Physician* 2012;85:123–32.
- 23 Doubilet PM, Cain KC. The superiority of sequential over simultaneous testing. *Med Decis Making* 1985;5:447–51.
- 24 Anon. SCAT3. *Br J Sports Med* 2013;47:259.
- 25 Akobeng AK. Understanding diagnostic tests 1: sensitivity, specificity and predictive values. *Acta Paediatr* 2007;96:338–41.
- 26 Eckner JT, Kutcher JS. Concussion symptom scales and sideline assessment tools: a critical literature update. *Curr Sports Med Rep* 2010;9:8–15.
- 27 Alla S, Sullivan SJ, Hale L, *et al.* Self-report scales/checklists for the measurement of concussion symptoms: a systematic review. *Br J Sports Med* 2009;43 (Suppl 1):i3–12.
- 28 Yengo-Kahn AM, Hale AT, Zalneraitis BH, *et al.* The Sport Concussion Assessment Tool: a systematic review. *Neurosurg Focus* 2016;40:E6.
- 29 Bell DR, Guskiewicz KM, Clark MA, *et al.* Systematic review of the balance error scoring system. *Sports Health* 2011;3:287–95.
- 30 Hunt AW, Mah K, Reed N, *et al.* Oculomotor-based vision assessment in mild traumatic brain injury: a systematic review. *J Head Trauma Rehabil* 2016;31:252–61.
- 31 Galetta KM, Liu M, Leong DF, *et al.* The King-Devick test of rapid number naming for concussion detection: meta-analysis and systematic review of the literature. *Concussion* 2015;15:1–15.
- 32 King D, Brughelli M, Hume P, *et al.* Assessment, management and knowledge of sport-related concussion: systematic review. *Sports Med* 2014;44:449–71.
- 33 Maddocks DL, Dicker GD, Saling MM. The assessment of orientation following concussion in athletes. *Clin J Sport Med* 1995;5:32–5.
- 34 McCrory PR, Ariens T, Berkovic SF. The nature and duration of acute concussive symptoms in Australian football. *Clin J Sport Med* 2000;10:235–8.
- 35 Barr WB, McCrea M. Sensitivity and specificity of standardized neurocognitive testing immediately following sports concussion. *J Int Neuropsychol Soc* 2001;7:693–702.
- 36 Erlanger D, Kaushik T, Cantu R, *et al.* Symptom-based assessment of the severity of a concussion. *J Neurosurg* 2003;98:477–84.
- 37 McCrea M. Standardized mental status testing on the sideline after sport-related concussion. *J Athl Train* 2001;36:274–9.
- 38 McCrea M, Kelly JP, Randolph C, *et al.* Immediate neurocognitive effects of concussion. *Neurosurgery* 2002;50:1032–40.
- 39 McCrea M, Barr WB, Guskiewicz K, *et al.* Standard regression-based methods for measuring recovery after sport-related concussion. *J Int Neuropsychol Soc* 2005;11:58–69.
- 40 Echlin PS, Tator CH, Cusimano MD, *et al.* Return to play after an initial or recurrent concussion in a prospective study of physician-observed junior ice hockey concussions: implications for return to play after a concussion. *Neurosurg Focus* 2010;29:E5.
- 41 Galetta KM, Barrett J, Allen M, *et al.* The King-Devick test as a determinant of head trauma and concussion in boxers and MMA fighters. *Neurology* 2011;76:1456–62.
- 42 Galetta KM, Brandes LE, Maki K, *et al.* The King-Devick test and sports-related concussion: study of a rapid visual screening tool in a collegiate cohort. *J Neurol Sci* 2011;309:34–9.
- 43 Barr WB, Pritchep LS, Chabot R, *et al.* Measuring brain electrical activity to track recovery from sport-related concussion. *Brain Inj* 2012;26:58–66.
- 44 King D, Clark T, Gissane C. Use of a rapid visual screening tool for the assessment of concussion in amateur rugby league: a pilot study. *J Neurol Sci* 2012;320:16–21.
- 45 Galetta MS, Galetta KM, McCrossin J, *et al.* Saccades and memory: baseline associations of the King-Devick and SCAT2 SAC tests in professional ice hockey players. *J Neurol Sci* 2013;328:28–31.
- 46 Dhawan PSA, Tapsell L, Adler J, *et al.* King-Devick test identifies symptomatic concussion in real-time and asymptomatic concussion over time. *Neurol Clin Pract* 2014;82:S11.
- 47 Leong DF, Balcer LJ, Galetta SL, *et al.* The King-Devick test as a concussion screening tool administered by sports parents. *J Sports Med Phys Fitness* 2014;54:70–7.
- 48 Galetta KM, Morganroth J, Moehringer N, *et al.* Adding vision to concussion testing: a prospective study of sideline testing in youth and collegiate athletes. *J Neuroophthalmol* 2015;35:235–41.
- 49 Leong DF, Balcer LJ, Galetta SL, *et al.* The King-Devick test for sideline concussion screening in collegiate football. *J Optom* 2015;8:131–9.
- 50 Marinides Z, Galetta KM, Andrews CN, *et al.* Vision testing is additive to the sideline assessment of sports-related concussion. *Neurology* 2015;5:25–34.
- 51 Putukian M, Echemendia R, Dettwiler-Danspeckgruber A, *et al.* Prospective clinical assessment using sideline Concussion Assessment Tool-2 testing in the evaluation of sport-related concussion in college athletes. *Clin J Sport Med* 2015;25:36–42.
- 52 Seidman DH, Burlingame J, Yousif LR, *et al.* Evaluation of the King-Devick test as a concussion screening tool in high school football players. *J Neurol Sci* 2015;356:97–101.