

Supplementary File 2. Summary of included trials (n=88) and comparisons (n=99) showing control group fall rate over follow-up period, sample size, length of follow-up, estimate of effect of intervention, and number and percentage of trials with each population, program, or study quality descriptor.

First author, year and program type if two intervention groups ^a	Control group falls/person year or % who fell in follow up period	Sample size at randomisation	Follow-up (weeks)	Estimate of fall rate ratio	High support residential care ^b	Population risk ^c	Average age over 75 years	Mod/high intensity strength training ^d	High challenge balance training ^e	Mod/high intensity endurance training ^f	Stretching program ^g	Walking program ^h	Ten or fewer exercise participants/instructor ⁱ	Exercises tailored in type or intensity ^j	Good adherence ^k	50+ total hours of exercise ^l	PEDro Score ^m
Arkkukangas, 2015 ^{a1}	1.23	45	12	0.72	N	2	Y	Y	Y	N	N	Y	N	Y	Y	N	6
Barnett, 2003 ²	0.95	163	52	0.60	N	2	Y	Y	Y	Y	Y	Y	Y	N	Y	Y	8
Beling, 2009 ³	1.13	23	12	0.18	N	2	Y	N	Y	N	N	Y	Y	N	N	N	4
Buchner, 1997 ⁴	0.81	105	72	0.61	N	2	Y	Y	N	Y	N	N	N	Y	Y	Y	7
Bunout, 2005 ⁵	0.18	298	52	1.22	N	3	Y	Y	N	Y	N	Y	Y	Y	Y	Y	5
Campbell, 1997 ⁶	1.35	233	50	0.68	N	2	Y	Y	Y	Y	Y	Y	N	Y	N	Y	8
Campbell, 1999 ⁷	0.97	93	44	0.87	N	1	N	Y	Y	Y	Y	Y	N	Y	Y	Y	6
Canning, 2015 ⁸	3.49	231	26	0.73	N	1	Y	Y	Y	N	N	N	N	Y	Y	Y	8
Carter, 2002 ⁹	0.52	93	20	0.88	N	3	N	N	N	N	Y	N	N	N	Y	N	5
Clegg, 2014 ¹⁰	27%	84	14	0.66	N	2	Y	N	N	N	N	N	N	Y	N	N	7
Clemson, 2010 ¹¹	0.12	34	26	0.21	N	2	N	N	Y	N	N	N	N	Y	N	Y	8
Clemson, 2012, ¹² LiFE - structured training ¹²	2.46	317	52	0.69	N	2	N	Y	Y	N	N	N	N	Y	Y	Y	7
				0.81	N	2	N	Y	Y	N	N	N	N	Y	Y	Y	7

Cornillon, 2002 ¹³	0.48	303	52	0.82	N	3	N	Y	N	N	N	N	N	N	Y	N	5
Dangour, 2011 ¹⁴	0.99	2002	104	0.95	N	3	Y	N	N	N	N	Y	Y	N	N	Y	7
Day, 2002 ¹⁵	64%	1090	18	0.79	N	3	Y	N	N	N	Y	N	N	N	Y	N	6
De Sure, 2013 ¹⁶	0.22	27	24	0.69	N	1	Y	N	N	N	N	N	Y	N	Y	N	3
Dean, 2012 ¹⁷	1.96	151	52	0.96	N	1	N	Y	Y	N	N	Y	Y	Y	Y	Y	8
Ebrahim, 1997 ¹⁸	0.29	165	104	1.29	N	3	N	N	N	Y	N	Y	N	N	Y	Y	6
El-Khoury, 2015 ¹⁹	4.44	706	104	0.88	N	2	Y	N	Y	N	Y	N	N	Y	N	Y	7
Faber, 2006, ²⁰ functional - Tai Chi	2.5	278	52	1.32	N	1	Y	N	Y	N	N	Y	Y	Y	Y	N	6
				0.96	N	1	Y	N	Y	N	Y	N	Y	Y	Y	N	6
Freiberger, 2012 ²¹	1.58	378	52	0.88	N	2	Y	N	N	N	Y	Y	N	N	Y	N	8
Gao, 2014 ²²	0.24	80	26	0.44	N	1	N	N	N	N	N	N	N	N	Y	N	6
Gianoudis, 2014 ²³	0.46	162	52	1.22	N	3	N	Y	Y	N	Y	N	Y	Y	Y	Y	6
Goodwin, 2011 ²⁴	24.3	130	20	0.68	N	1	N	N	Y	N	N	N	N	Y	Y	N	7
Grahn Kronhed 2009 ²⁵	0.8	65	52	0.75	N	3	N	Y	N	N	Y	N	Y	Y	Y	N	7
Haines, 2009 ²⁶	3.29	53	26	0.72	N	1	Y	N	Y	N	N	N	N	Y	N	N	7
Halvarsson, 2013 ²⁷	0.14	59	65	5.40	N	2	Y	N	N	N	N	N	Y	Y	Y	N	5
Hirase, 2015, ²⁸ foam surface - stable surface	2.93	93	16	0.53	N	2	Y	N	N	N	N	Y	Y	N	Y	N	6
				1.31	N	2	Y	N	N	N	N	Y	Y	N	Y	N	6
Huang, 2010 ²⁹	0.02	261	22	0.81	N	3	N	N	N	N	N	N	N	N	N	N	4
Huang, 2011 ³⁰	0.34	80	20	0.38	N	3	Y	N	N	N	N	N	N	N	N	N	8
Huang, 2015 ³¹	0.18	80	22	0.04	Y	1	Y	N	N	Y	N	N	Y	N	Y	N	8
Iliffe, 2015, ³² FAME - Otago Exercise Program	0.86	1256	76	0.74	N	3	N	Y	Y	N	Y	Y	N	Y	N	Y	6
				0.76	N	3	N	Y	Y	N	N	Y	N	Y	N	Y	6
Irez, 2011 ³³	5.6	60	12	0.28	N	1	N	N	N	N	N	N	N	Y	Y	N	6
Iwamoto, 2009 ³⁴	0.32	68	20	0.10	N	3	Y	N	N	N	N	N	Y	N	Y	N	6
Kamide, 2009 ³⁵	0.04	57	52	0.38	N	3	N	Y	N	N	Y	N	N	N	Y	N	6
Karinkanta, 2007, ³⁶ balance - resistance & balance	0.36	149	52	1.43	N	3	N	N	Y	N	N	N	N	N	Y	Y	6
				1.46	N	3	N	Y	Y	N	N	N	Y	Y	Y	Y	6
Kim, 2014 ³⁷	0.4	105	52	0.49	N	2	Y	Y	Y	N	N	N	N	Y	Y	Y	6
Korpelainen, 2006 ³⁸	0.6	160	130	0.79	N	3	N	Y	Y	Y	N	Y	Y	Y	Y	Y	8
Kovacs, 2013 ³⁹	0.97	86	52	0.71	Y	1	Y	Y	N	N	N	Y	Y	N	N	Y	8

Kruse, 2010 ⁴⁰	0.82	79	52	0.99	N	3	N	N	Y	N	N	Y	Y	Y	N	N	8
Latham, 2003 ⁴¹	2.7	243	26	1.08	N	1	Y	Y	N	N	N	N	N	Y	Y	N	8
Lehtola, 2000 ⁴²	8%	131	26	0.21	N	3	N	N	N	N	Y	Y	N	N	Y	Y	
Li, 2005 ⁴³	0.97	256	52	0.45	N	3	Y	N	Y	N	N	N	N	N	Y	Y	5
Lin, 2007 ⁴⁴	0.88	150	26	0.67	N	2	Y	N	Y	N	Y	Y	N	Y	N	N	5
Liu-Ambrose, 2004, ⁴⁵ agility	0.65	104	25	1.03	N	2	Y	N	Y	Y	N	N	Y	N	Y	N	6
- resistance				1.80	N	2	Y	Y	N	N	N	N	Y	Y	Y	N	6
Liu-Ambrose, 2008 ⁴⁶	67%	74	26	0.65	N	2	Y	Y	Y	N	N	Y	N	Y	Y	Y	7
Logghe, 2009 ⁴⁷	0.91	269	52	1.16	N	2	Y	N	Y	N	N	N	Y	N	Y	N	8
Lord, 1995 ⁴⁸	0.63	197	52	0.85	N	3	N	N	Y	Y	Y	Y	N	N	Y	Y	4
Lord, 2003 ⁴⁹	0.85	551	52	0.78	N	3	Y	N	Y	Y	Y	Y	N	N	N	Y	6
Luukinen, 2007 ⁵⁰	1.23	486	70	0.93	N	2	Y	N	N	N	Y	Y	N	Y	N	Y	6
Madureira, 2007 ⁵¹	0.9	66	52	0.48	N	3	N	N	Y	N	N	Y	N	N	Y	Y	6
Maduerira, 2010 ⁵²	0.4	66	52	2.25	N	3	N	N	N	N	Y	Y	N	N	Y	Y	6
McMurdo, 1997 ⁵³	0.33	118	104	0.53	N	3	N	N	N	N	N	N	N	N	Y	Y	4
Means, 2005 ⁵⁴	1.18	338	26	0.41	N	3	N	Y	N	Y	Y	Y	Y	Y	Y	N	5
Morris, 2015 ⁵⁵ mvt strat	15.47	210	52	0.39	N	1	N	N	N	N	N	Y	Y	Y	Y	Y	8
- resistance				0.15	N	1	N	Y	N	N	N	N	Y	Y	Y	Y	8
Mulrow, 1994 ⁵⁶	2.05	194	16	1.26	Y	1	Y	Y	N	N	N	Y	Y	Y	Y	N	6
Patil 2015 ⁵⁷	2.50	409	104	1.0	N	1	N	Y	N	N	N	N	N	Y	Y	Y	7
Pitkala, 2013 ⁵⁸	3.35	210	52	0.50	N	1	Y	N	Y	N	N	Y	Y	Y	Y	Y	8
Protas, 2006 ⁵⁹	37.6	18	2	0.62	N	1	N	N	Y	Y	N	Y	Y	Y	Y	N	5
Reinsch, 1992 ⁶⁰	40%	230	52	1.24	N	3	N	N	N	N	N	N	N	N	Y	Y	4
Resnick, 2002 ⁶¹	0.56	20	26	0.71	N	1	Y	N	N	Y	N	Y	Y	N	Y	N	4
Robertson, 2001 ⁶²	1.01	240	52	0.54	N	2	Y	Y	Y	Y	Y	Y	N	Y	Y	Y	5
Rolland, 2007 ⁶³	2.52	134	52	0.99	Y	1	Y	N	N	N	Y	Y	Y	Y	N	Y	8
Rosendahl, 2008 ⁶⁴	58%	191	26	0.82	Y	1	Y	Y	Y	N	N	Y	Y	Y	Y	Y	7
Rubenstein, 2000 ⁶⁵	2.25	59	12	0.90	N	2	Y	Y	N	Y	N	Y	Y	N	Y	Y	7
Sakamoto, 2006 ⁶⁶	1.14	553	26	0.82	Y	1	Y	N	N	N	N	N	Y	N	Y	N	4
Sakamoto, 2013 ⁶⁷	0.66	1365	26	0.85	N	2	Y	N	N	N	N	N	N	Y	N	N	4
Schnelle, 2003 ⁶⁸	0.99	190	32	0.62	Y	1	Y	N	N	N	N	Y	Y	Y	Y	Y	6

Schoenfelder, 2000 ⁶⁹	3.43	16	26	3.06	Y	1	Y	N	N	N	N	Y	Y	N	Y	N	4
Sherrington, 2014 ⁷⁰	0.73	340	52	1.43	N	1	Y	Y	Y	N	N	N	N	Y	N	Y	8
Sihvonen, 2004 ⁷¹	1.57	27	52	0.38	Y	1	Y	N	Y	N	N	N	Y	Y	Y	N	5
Skelton, 2005 ⁷²	3.21	81	86	0.69	N	2	N	N	Y	Y	Y	N	Y	Y	Y	Y	6
Smulders, 2010 ⁷³	1.21	96	52	0.61	N	2	N	N	N	N	N	Y	Y	N	Y	N	8
Steinberg, 2000 ⁷⁴	0.85	132	73	0.90	N	3	N	N	N	N	N	N	N	N	N	N	5
Suttanon, 2013 ⁷⁵	0.9	40	26	1.00	N	1	Y	Y	Y	N	N	Y	N	Y	Y	Y	6
Suzuki, 2004 ⁷⁶	55%	52	87	0.35	N	3	Y	N	Y	N	Y	N	N	N	Y	N	6
Swanenburg, 2007 ⁷⁷	0.6	24	52	0.17	N	3	N	Y	N	N	N	N	Y	Y	Y	N	6
Taylor Pilae, 2014 ⁷⁸ , Tai Chi	2.32	101	12	0.31	N	1	N	N	N	N	N	N	N	N	Y	N	8
- Silver Sneakers program				0.84	N	1	N	N	N	N	N	N	N	N	Y	N	8
Toulotte, 2003 ⁷⁹	1.95	20	16	0.08	Y	1	Y	N	Y	N	Y	Y	Y	N	Y	N	3
Trombeti, 2011 ⁸⁰	1.9	134	26	0.46	N	2	Y	N	N	N	N	N	N	N	Y	N	7
Uusi-Rasi, 2015 ⁸¹	1.18	205	104	1.07	N	2	N	Y	N	N	N	N	N	Y	Y	Y	7
Voukelatos, 2007 ⁸²	24%	702	24	0.67	N	3	N	N	Y	N	N	N	N	N	Y	N	7
Voukelatos, 2015 ⁸³	40%	386	48	0.88	N	3	N	N	N	N	N	Y	N	N	N	N	7
Weerdesteyn, 2006 ⁸⁴	32%	113	28	0.54	N	2	N	N	N	N	N	Y	Y	N	Y	N	6
Wesson, 2013 ⁸⁵	0.31	22	16	0.34	N	1	Y	Y	Y	N	N	N	N	Y	Y	N	8
Wolf, 2003 ⁸⁶	60%	311	48	0.75	N	2	Y	N	Y	N	N	N	N	N	Y	Y	7
Wolf, 1996, ⁸⁷ balance	3.18	200	87	0.98	N	3	Y	N	N	N	N	N	Y	Y	Y	N	6
- Tai Chi			87	0.51	N	3	Y	N	Y	N	N	N	N	N	Y	Y	6
Woo, 2007, ⁸⁸ resistance	0.53	180	52	0.78	N	3	N	Y	N	N	N	N	N	N	Y	Y	6
- Tai Chi				0.49	N	3	N	Y	N	N	N	N	N	N	Y	Y	6

^a additional data provided by the author; ^b mainly high support care facility (nursing home) residents; ^c presence of a particular falls risk factor was used as inclusion criteria to the study or the entire population was known to be at increased risk (1=aged care facility residents, aged over 75, recent hospitalisation, impaired cognition, psychotropic medication use; 2=impaired strength or balance, previous falls; 3=no specific risk factor); ^d

moderate intensity (40-60% of the 1Repetition Maximum (RM) i.e. a weight so heavy that it can only be lifted once) or high intensity (>60% 1RM);^e highly challenging = all 3 of the following criteria: movement of the centre of mass, narrowing of the base of support, and minimising upper limb support; ^f moderate intensity = 40-60% max heart rate, some increase in breathing or heart rate, or perceived exertion of 11 to 14 on the [Borg scale](#) or high intensity = above 60% of max heart rate or heart rate reserve, large increase in breathing or heart rate (conversation is difficult or broken) or perceived exertion of 15 or greater on the [Borg scale](#);^g short- or long-duration stretches was specifically mentioned; ^h walking program/practice was specifically mentioned ; ⁱ ten or fewer participants per instructor; ^j type or intensity of most exercises was designed for each individual based on an assessment; ^k ≥75% participants attended 50% or more sessions and/or >50% attendance rate; ^l ≥ 50 hours of exercise with instructor plus prescribed home exercise over study period; ^m Coded using PEDro rating scale; N = no; Y = yes.

References to all trials included in the review.

1. Arkkukangas M, Johnson ST, Hellstrom K, et al. A feasibility study of a randomised controlled trial comparing fall prevention using exercise with or without the support of motivational interviewing. *Prev Med Rep* 2015;2:134-40.
2. Barnett A, Smith B, Lord SR, et al. Community-based group exercise improves balance and reduces falls in at-risk older people: a randomised controlled trial. *Age Ageing* 2003;32(4):407-14.
3. Beling J, Roller M. Multifactorial intervention with balance training as a core component among fall-prone older adults. *J Geriatr Phys Ther* 2009;32(3):125-33.
4. Buchner DM, Cress ME, de Lateur BJ, et al. The effect of strength and endurance training on gait, balance, fall risk, and health services use in community-living older adults. *J Gerontol A Biol Sci Med Sci* 1997;52(4):M218-24.
5. Bunout D, Barrera G, Avendano M, et al. Results of a community-based weight-bearing resistance training programme for healthy Chilean elderly subjects. *Age Ageing* 2005;34(1):80-3.

6. Campbell AJ, Robertson MC, Gardner MM, et al. Randomised controlled trial of a general practice programme of home based exercise to prevent falls in elderly women. *BMJ* 1997;315(7115):1065-9.
7. Campbell AJ, Robertson MC, Gardner MM, et al. Psychotropic medication withdrawal and a home-based exercise program to prevent falls: a randomized, controlled trial. *J Am Geriatr Soc* 1999;47(7):850-3.
8. Canning CG, Sherrington C, Lord SR, et al. Exercise for falls prevention in Parkinson disease: a randomized controlled trial. *Neurology* 2015;84(3):304-12.
9. Carter ND, Khan KM, McKay HA, et al. Community-based exercise program reduces risk factors for falls in 65- to 75-year-old women with osteoporosis: randomized controlled trial. *CMAJ* 2002;167(9):997-1004.
10. Clegg A, Barber S, Young J, et al. The Home-based Older People's Exercise (HOPE) trial: a pilot randomised controlled trial of a home-based exercise intervention for older people with frailty. *Age Ageing* 2014;43(5):687-95.
11. Clemson L, Singh MF, Bundy A, et al. LiFE Pilot Study: A randomised trial of balance and strength training embedded in daily life activity to reduce falls in older adults. *Aust Occup Ther J* 2010;57(1):42-50.
12. Clemson L, Fiatarone Singh MA, Bundy A, et al. Integration of balance and strength training into daily life activity to reduce rate of falls in older people (the LiFE study): randomised parallel trial. *BMJ* 2012;345:e4547.
13. Cornillon E, Blanchon MA, Ramboatsisetraina P, et al. Impact d'un programme de prevention multidisciplinaire de la chute chez le sujet age autonome vivant a domicile, avec analyse avant-apres des performances physiques. *Annales de Readaptation et de Medecine Physique* 2002;45(9):493-504.
14. Dangour AD, Albala C, Allen E, et al. Effect of a nutrition supplement and physical activity program on pneumonia and walking capacity in Chilean older people: a factorial cluster randomized trial. *PLoS Med* 2011;8(4):e1001023.
15. Day L, Fildes B, Gordon I, et al. Randomised factorial trial of falls prevention among older people living in their own homes. *BMJ* 2002;325(7356):128.
16. DeSure AR, Peterson K, Gianan FV, et al. An exercise program to prevent falls in institutionalized elderly with cognitive deficits: a crossover pilot study. *Hawaii J Med Public Health* 2013;72(11):391-5.
17. Dean CM, Rissel C, Sherrington C, et al. Exercise to enhance mobility and prevent falls after stroke: the community stroke club randomized trial. *Neurorehabil Neural Repair* 2012;26(9):1046-57.
18. Ebrahim S, Thompson PW, Baskaran V, et al. Randomized placebo-controlled trial of brisk walking in the prevention of postmenopausal osteoporosis. *Age Ageing* 1997;26(4):253-60.
19. El-Khoury F, Cassou B, Latouche A, et al. Effectiveness of two year balance training programme on prevention of fall induced injuries in at risk women aged 75-85 living in community: Ossebo randomised controlled trial. *BMJ* 2015;351:h3830.

20. Faber MJ, Bosscher RJ, Chin APMJ, et al. Effects of exercise programs on falls and mobility in frail and pre-frail older adults: A multicenter randomized controlled trial. *Arch Phys Med Rehabil* 2006;87(7):885-96.
21. Freiberger E, Haberle L, Spirduso WW, et al. Long-term effects of three multicomponent exercise interventions on physical performance and fall-related psychological outcomes in community-dwelling older adults: a randomized controlled trial. *J Am Geriatr Soc* 2012;60(3):437-46.
22. Gao Q, Leung A, Yang Y, et al. Effects of Tai Chi on balance and fall prevention in Parkinson's disease: a randomized controlled trial. *Clinical Rehabil* 2014 Feb 11:Epub ahead of print 2014.
23. Gianoudis J, Bailey CA, Ebeling PR, et al. Effects of a targeted multimodal exercise program incorporating high-speed power training on falls and fracture risk factors in older adults: a community-based randomized controlled trial. *J Bone Miner Res* 2014 Jan;29(1):182-191 2014.
24. Goodwin VA, Richards SH, Henley W, et al. An exercise intervention to prevent falls in people with Parkinson's disease: a pragmatic randomised controlled trial. *J Neurol Neurosurg Psychiatry* 2011 Nov;82(11):1232-1238 2011.
25. Grahn Kronhed A, Hallberg I, Odqvist L, et al. Effect of training on health-related quality of life, pain and falls in osteoporotic women. *Advances in Physiotherapy* 2009;11:154-65.
26. Haines TP, Russell T, Brauer SG, et al. Effectiveness of a video-based exercise programme to reduce falls and improve health-related quality of life among older adults discharged from hospital: a pilot randomized controlled trial. *Clin Rehabil* 2009;23(11):973-85.
27. Halvarsson A, Franzén E, Farén E, et al. Long-term effects of new progressive group balance training for elderly people with increased risk of falling – a randomized controlled trial. *Clin Rehabil* 2013;27(5):450-58.
28. Hirase T, Inokuchi S, Matsusaka N, et al. Effects of a balance training program using a foam rubber pad in community-based older adults: a randomized controlled trial. *J Geriatr Phys Ther* 2015;38(2):62-70.
29. Huang HC, Liu CY, Huang YT, et al. Community-based interventions to reduce falls among older adults in Taiwan - long time follow-up randomised controlled study. *J Clin Nurs* 2010; 19(7-8).
30. Huang TT, Yang LH, Liu CY. Reducing the fear of falling among community-dwelling elderly adults through cognitive-behavioural strategies and intense Tai Chi exercise: a randomized controlled trial. *J Adv Nurs* 2011;67(5):961-71.
31. Huang TT, Chung ML, Chen FR, et al. Evaluation of a combined cognitive-behavioural and exercise intervention to manage fear of falling among elderly residents in nursing homes. *Aging Ment Health* 2016;20(1):2-12.
32. Iliffe S, Kendrick D, Morris R, et al. Promoting physical activity in older people in general practice: ProAct65+ cluster randomised controlled trial. *Br J Gen Pract* 2015;65(640):e731-8.
33. Irez GB, Ozdemir RA, Evin R, et al. Integrating pilates exercise into an exercise program for 65+ year-old women to reduce falls. *J Sports Sci Med* 2011;10(1):105-11.

34. Iwamoto J, Suzuki H, Tanaka K, et al. Preventative effect of exercise against falls in the elderly: a randomized controlled trial. *Osteoporos Int* 2009;20(7):1233-40.
35. Kamide N, Shiba Y, Shibata H. Effects on balance, falls, and bone mineral density of a home-based exercise program without home visits in community-dwelling elderly women: a randomized controlled trial. *J Physiol Anthropol* 2009;28(3):115-22.
36. Karinkanta S, Heinonen A, Sievanen H, et al. A multi-component exercise regimen to prevent functional decline and bone fragility in home-dwelling elderly women: randomized, controlled trial. *Osteoporos Int* 2007;18(4):453-62.
37. Kim H, Yoshida H, Suzuki T. Falls and fractures in participants and excluded non-participants of a fall prevention exercise program for elderly women with a history of falls: 1-year follow-up study. *Geriatr Gerontol Int* 2014; 14(2).
38. Korpelainen R, Keinänen-Kiukaanniemi S, Heikkinen J, et al. Effect of exercise on extraskeletal risk factors for hip fractures in elderly women with low BMD: a population-based randomized controlled trial. *J Bone Miner Res* 2006;21(5):772-9.
39. Kovacs E, Sztruhar Jonasne I, Karoczi CK, et al. Effects of a multimodal exercise program on balance, functional mobility and fall risk in older adults with cognitive impairment: a randomized controlled single-blind study. *European Journal of Physical and Rehabilitation Medicine* 2013 Oct;49(5):639-648 2013.
40. Kruse RL, Lemaster JW, Madsen RW. Fall and balance outcomes after an intervention to promote leg strength, balance, and walking in people with diabetic peripheral neuropathy: "feet first" randomized controlled trial. *Phys Ther* 2010; 90(11).
41. Latham NK, Anderson CS, Lee A, et al. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: the Frailty Interventions Trial in Elderly Subjects (FITNESS). *Journal of the American Geriatrics Society* 2003;51(3):291-9.
42. Lehtola S, Hanninen L, Paatalo M. Kaatumistapaturmien ilmaantuvuus 70- to 75-vuotiailla oululaisilla liikuntaintervention ja sen jalkeisen seurannan aikana (The incidence of falls during six-months exercise trial and four-months follow-up among home-dwelling persons aged 70 to 75 years) [Finnish]. *Liikunta and Tiede* 2000;37(6):41-46.
43. Li F, Harmer P, Fisher KJ, et al. Tai Chi and fall reductions in older adults: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 2005;60(2):187-94.
44. Lin MR, Wolf SL, Hwang HF, et al. A randomized, controlled trial of fall prevention programs and quality of life in older fallers. *J Am Geriatr Soc* 2007;55(4):499-506.
45. Liu-Ambrose T, Khan KM, Eng JJ, et al. Resistance and agility training reduce fall risk in women aged 75 to 85 with low bone mass: a 6-month randomized, controlled trial. *J Am Geriatr Soc* 2004;52(5):657-65.
46. Liu-Ambrose T, Donaldson MG, Ahamed Y, et al. Otago home-based strength and balance retraining improves executive functioning in older fallers: a randomized controlled trial. *J Am Geriatr Soc* 2008;56(10):1821-30.
47. Logghe IH, Zeeuwe PE, Verhagen AP, et al. Lack of effect of Tai Chi Chuan in preventing falls in elderly people living at home: a randomized clinical trial. *J Am Geriatr Soc* 2009;57(1):70-5.

48. Lord SR, Ward JA, Williams P, et al. The effect of a 12-month exercise trial on balance, strength, and falls in older women: a randomized controlled trial. *J Am Geriatr Soc* 1995;43(11):1198-206.
49. Lord SR, Castell S, Corcoran J, et al. The effect of group exercise on physical functioning and falls in frail older people living in retirement villages: a randomized, controlled trial. *J Am Geriatr Soc* 2003;51(12):1685-92.
50. Luukinen H, Lehtola S, Jokelainen J, et al. Pragmatic exercise-oriented prevention of falls among the elderly: a population-based, randomized, controlled trial. *Prev Med* 2007;44(3):265-71.
51. Madureira MM, Takayama L, Gallinaro AL, et al. Balance training program is highly effective in improving functional status and reducing the risk of falls in elderly women with osteoporosis: a randomized controlled trial. *Osteoporos Int* 2007;18(4):419-25.
52. Madureira MM, Bonfa E, Takayama L, et al. A 12-month randomized controlled trial of balance training in elderly women with osteoporosis: improvement of quality of life. *Maturitas* 2010;66(2):206-11.
53. McMurdo ME, Mole PA, Paterson CR. Controlled trial of weight bearing exercise in older women in relation to bone density and falls. *BMJ* 1997;314(7080):569.
54. Means KM, Rodell DE, O'Sullivan PS. Balance, mobility, and falls among community-dwelling elderly persons: effects of a rehabilitation exercise program. *Am J Phys Med Rehabil* 2005;84(4):238-50.
55. Morris ME, Menz HB, McGinley JL, et al. A Randomized Controlled Trial to Reduce Falls in People With Parkinson's Disease. *Neurorehabil Neural Repair* 2015;29(8):777-85.
56. Mulrow CD, Gerety MB, Kanten D, et al. A randomized trial of physical rehabilitation for very frail nursing home residents. *JAMA* 1994;271(7):519-24.
57. Patil R, Uusi-Rasi K, Tokola K, et al. Effects of a Multimodal Exercise Program on Physical Function, Falls, and Injuries in Older Women: A 2-Year Community-Based, Randomized Controlled Trial. *J Am Geriatr Soc* 2015;63(7):1306-13.
58. Pitkala KH, Poysti MM, Laakkonen ML, et al. Effects of the FINnish ALzheimer disease EXercise trial (FINALEX): a randomized controlled trial. *JAMA Internal Medicine* 2013 May 27;173(10):894-901 2013.
59. Protas EJ, Mitchell K, Williams A, et al. Gait and step training to reduce falls in Parkinson's disease. *NeuroRehabilitation* 2005;20(3):183-90.
60. Reinsch S, MacRae P, Lachenbruch PA, et al. Attempts to prevent falls and injury: a prospective community study. *Gerontologist* 1992;32(4):450-6.
61. Resnick B. Testing the effect of the WALC intervention on exercise adherence in older adults. *J Gerontol Nurs* 2002;28(6):40-9.
62. Robertson MC, Devlin N, Gardner MM, et al. Effectiveness and economic evaluation of a nurse delivered home exercise programme to prevent falls. 1: Randomised controlled trial. *BMJ* 2001;322(7288):697-701.

63. Rolland Y, Pillard F, Klapouszczak A, et al. Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial. *J Am Geriatr Soc* 2007;55(2):158-65.
64. Rosendahl E, Gustafson Y, Nordin E, et al. A randomized controlled trial of fall prevention by a high-intensity functional exercise program for older people living in residential care facilities. *Aging Clin Exp Res* 2008;20(1):67-75.
65. Rubenstein LZ, Josephson KR, Trueblood PR, et al. Effects of a group exercise program on strength, mobility, and falls among fall-prone elderly men. *J Gerontol A Biol Sci Med Sci* 2000;55(6):M317-21.
66. Sakamoto K, Nakamura T, Hagino H, et al. Effects of unipedal standing balance exercise on the prevention of falls and hip fracture among clinically defined high-risk elderly individuals: a randomized controlled trial. *J Orthop Sci* 2006;11(5):467-72.
67. Sakamoto K, Endo N, Harada A, et al. Why not use your own body weight to prevent falls? A randomized, controlled trial of balance therapy to prevent falls and fractures for elderly people who can stand on one leg for <15 s. *J Ortho Sci* 2013; 18(1).
68. Schnelle JF, Kapur K, Alessi C, et al. Does an exercise and incontinence intervention save healthcare costs in a nursing home population? *J Am Geriatr Soc* 2003;51(2):161-8.
69. Schoenfelder DP. A fall prevention program for elderly individuals. Exercise in long-term care settings. *J Gerontol Nurs* 2000;26(3):43-51.
70. Sherrington C, Lord SR, Vogler CM, et al. A post-hospital home exercise program improved mobility but increased falls in older people: a randomised controlled trial. *PloS One* 2014;9(9):e104412.
71. Sihvonen S, Sipila S, Taskinen S, et al. Fall incidence in frail older women after individualized visual feedback-based balance training. *Gerontology* 2004;50(6):411-6.
72. Skelton D, Dinan S, Campbell M, et al. Tailored group exercise (Falls Management Exercise -- FaME) reduces falls in community-dwelling older frequent fallers (an RCT). *Age Ageing* 2005;34(6):636-9.
73. Smulders E, Weerdesteyn V, Groen BE, et al. Efficacy of a short multidisciplinary falls prevention program for elderly persons with osteoporosis and a fall history: a randomized controlled trial. *Arch Phys Med Rehab* 2010; 91(11).
74. Steinberg M, Cartwright C, Peel N, et al. A sustainable programme to prevent falls and near falls in community dwelling older people: results of a randomised trial. *J Epidemiol Community Health* 2000;54(3):227-32.
75. Suttanon P, Hill KD, Said CM, et al. Feasibility, safety and preliminary evidence of the effectiveness of a home-based exercise programme for older people with Alzheimer's disease: a pilot randomized controlled trial. *Clin Rehabil* 2013 May;27(5):427-438 2013.
76. Suzuki T, Kim H, Yoshida H, et al. Randomized controlled trial of exercise intervention for the prevention of falls in community-dwelling elderly Japanese women. *J Bone Miner Metab* 2004;22(6):602-11.
77. Swanenburg J, de Bruin ED, Stauffacher M, et al. Effects of exercise and nutrition on postural balance and risk of falling in elderly people with decreased bone mineral density: randomized controlled trial pilot study. *Clin Rehabil* 2007;21(6):523-34.

78. Taylor-Piliae RE, Hoke TM, Hepworth JT, et al. The effect of Tai Chi on physical function, fall rates and quality of life among older stroke survivors. *Arch Phys Med Rehab* 2014 May;95(5):816-824 2014.
79. Toulotte C, Fabre C, Dangremont B, et al. Effects of physical training on the physical capacity of frail, demented patients with a history of falling: a randomised controlled trial. *Age Ageing* 2003;32(1):67-73.
80. Trombetti A, Hars M, Herrmann FR, et al. Effect of music-based multitask training on gait, balance, and fall risk in elderly people: a randomized controlled trial. *Arch Intern Med* 2011;171(6):525-33.
81. Uusi-Rasi K, Patil R, Karinkanta S, et al. Exercise and vitamin D in fall prevention among older women: a randomized clinical trial. *JAMA Intern Med* 2015;175(5):703-11.
82. Voukelatos A, Cumming RG, Lord SR, et al. A randomized, controlled trial of tai chi for the prevention of falls: the Central Sydney tai chi trial. *J Am Geriatr Soc* 2007;55(8):1185-91.
83. Voukelatos A, Merom D, Sherrington C, et al. The impact of a home-based walking programme on falls in older people: the Easy Steps randomised controlled trial. *Age Ageing* 2015;44(3):377-83.
84. Weerdesteyn V, Rijken H, Geurts AC, et al. A five-week exercise program can reduce falls and improve obstacle avoidance in the elderly. *Gerontology* 2006;52(3):131-41.
85. Wesson J, Clemson L, Brodaty H, et al. A feasibility study and pilot randomised trial of a tailored prevention program to reduce falls in older people with mild dementia. *BMC Geriatrics* 2013 Sep 3;13(89):Epub 2013.
86. Wolf SL, Sattin RW, Kutner M, et al. Intense tai chi exercise training and fall occurrences in older, transitionally frail adults: a randomized, controlled trial. *J Am Geriatr Soc* 2003;51(12):1693-701.
87. Wolf SL, Barnhart HX, Kutner NG, et al. Reducing frailty and falls in older persons: an investigation of Tai Chi and computerized balance training. Atlanta FICSIT Group. *Frailty and Injuries: Cooperative Studies of Intervention Techniques*. *J Am Geriatr Soc* 1996;44(5):489-97.
88. Woo J, Hong A, Lau E, et al. A randomised controlled trial of Tai Chi and resistance exercise on bone health, muscle strength and balance in community-living elderly people. *Age Ageing* 2007;36(3):262-8.