Wake up call for collegiate athlete sleep: narrative review and consensus recommendations from the NCAA Interassociation Task Force on Sleep and Wellness

Emily Kroshus,^{1,2} Jessica Wagner,³ David Wyrick,⁴ Amy Athey,⁵ Lydia Bell, Holly J Benjamin,⁶ Michael A Grandner,⁷ Christopher E Kline,⁸ Jessica M Mohler,⁹ J Roxanne Prichard,¹⁰ Nathaniel F Watson,¹¹ Brian Hainline³

¹Department of Pediatrics, University of Washington, Seattle, Washington, USA
²Department of Pediatrics, Seattle Children's Research Institute, Seattle, Washington, USA
³National Collegiate Athletic Association, Indianapolis, Indiana, USA
⁴Department of Public Health Education, University of North Carolina at Greensboro, Greensboro, North Carolina, USA
⁵University of Arizona, Tucson, Arizona, USA
⁶Orthopedic Surgery and Rehabilitation Medicine, University of Chicago, Chicago, Illinois, USA
⁷Department of Psychiatry, University of Arizona, Tucson, Arizona, USA
⁸Department of Health and Physical Activity, University of Pittsburgh, Pittsburgh, Pennsylvania, USA
⁹US Naval Institute, Annapolis, Maryland, USA
¹⁰Department of Psychology, University of St Thomas, St Thomas, Minnesota, USA
¹¹Department of Neurology, University of Washington, Seattle, Washington, USA

Correspondence to

Dr Emily Kroshus;
 ekroshus@u.washington.edu,
 ekroshus@gmail.com

Accepted 28 February 2019
 Published Online First
 8 May 2019

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Kroshus E, Wagner J, Wyrick D, et al. *Br J Sports Med* 2019;**53**:731–736.

ABSTRACT

Sleep is an important determinant of collegiate athlete health, well-being and performance. However, collegiate athlete social and physical environments are often not conducive to obtaining restorative sleep. Traditionally, sleep has not been a primary focus of collegiate athletic training and is neglected due to competing academic, athletic and social demands. Collegiate athletics departments are well positioned to facilitate better sleep culture for their athletes. Recognising the lack of evidence-based or consensus-based guidelines for sleep management and restorative sleep for collegiate athletes, the National Collegiate Athletic Association hosted a sleep summit in 2017. Members of the Interassociation Task Force on Sleep and Wellness reviewed current data related to collegiate athlete sleep and aimed to develop consensus recommendations on sleep management and restorative sleep using the Delphi method. In this paper, we provide a narrative review of four topics central to collegiate athlete sleep: (1) sleep patterns and disorders among collegiate athletes; (2) sleep and optimal functioning among athletes; (3) screening, tracking and assessment of athlete sleep; and (4) interventions to improve sleep. We also present five consensus recommendations for colleges to improve their athletes' sleep.

INTRODUCTION

Sleep is critical for overall health and well-being. The Healthy People 2020 objectives, a set of 10-year national objectives for the health of all Americans, prioritised sleep and set out objectives to increase sufficient sleep, increase diagnosis and treatment of obstructive sleep apnea and prevent accidents due to drowsy driving.¹ Poor sleep disproportionately negatively impacts those in late adolescence through their early 20s (emerging adults).² The relationship between sleep duration and cardiometabolic risk and neurocognitive dysfunction is especially pronounced among young adults.³ Although fewer studies have focused on collegiate athletes, the documented outcomes of insufficient sleep duration, poor sleep quality (restorative sleep) or both include many critically important domains, spanning physical and academic performance, cognitive function, recovery from athletic exertion and injury, mental health and cardiometabolic

What is already known

- ▶ Sleep is an important determinant of collegiate athlete health, well-being and performance.
- ▶ Collegiate athletes often neglect sleep because of academic, athletic and social demands.

What are the new findings

- ▶ College athletic departments can take steps to promote collegiate athlete well-being and performance through sleep by:
 - Conducting an annual time demands survey.
 - Incorporating sleep screening into the preparticipation exam.
 - Providing evidence-based sleep education to collegiate athletes and coaches.

health. Collegiate athletes' social and physical environments are often not conducive to obtaining restorative sleep. Furthermore, their academic and athletic schedules may vary due to travel, practice and game schedules, different meal schedules, mandatory team meetings or other factors, all of which can prevent consistent bed and rise times.

Good sleep hygiene practices are summarised in box 1.⁴ A 2015 survey of National Collegiate Athletic Association (NCAA) collegiate athletes found that they spend 27–41 hours per week on athletics.⁵ After allowing time for academics, other extracurricular activities and relaxation or socialising, few have sufficient time left for restorative sleep.⁵ In addition, collegiate athletes often practice poor sleep hygiene habits. Caffeine, alcohol, marijuana and other illicit drugs and non-medical use of prescription drugs such as stimulants all can interfere with neurochemical control of sleep.⁶ Evening use of electronic devices suppresses melatonin levels and increases alertness, both of which delay sleep onset and contribute to sleep deprivation.⁶ These behaviours are compounded by student housing environments that can be too loud, bright, or crowded for consistent restorative sleep.⁷

Elite athletics programmes are beginning to recognise that collegiate athlete health and performance can be improved through sleep optimisation.^{8,9} In

Box 1 Sleep hygiene guidelines

1. Maintain a regular sleep schedule as much as possible, including on weekends. If this is not possible due to situational demands, establish a bedtime routine that can be practised regularly.
2. Seek bright light during the day, especially the morning, and avoid bright light at night if possible. This is especially important later at night, when bright light can further delay sleep onset.
3. Keep the bedroom should be kept cool, dark and comfortable.
4. Avoid caffeine (eg, coffee, energy drinks and certain types of soda and tea) at least 6 hours before bedtime. Nicotine is also a stimulant and should be avoided. Alcohol may reduce time to fall asleep, but it can adversely impact sleep quality and should be avoided close to bedtime.
5. Avoid consuming excessive food and liquids at night as these may disrupt sleep through reflux and increased trips to the bathroom.
6. Avoid obsessive clock watching. Often, looking at the clock at night can increase mental activity rather than decrease it and make resuming sleep more difficult.
7. Individuals who have difficulty falling asleep should generally avoid naps. However, collegiate athletes may find that napping during the day may improve performance and functioning.
8. Use beds for sleep (and sex) only. Any other activities, especially if lying awake and unable to sleep, should be done out of the bed. Those who have difficulty falling asleep should get out of bed for a period of time and try again later. This practice (called stimulus control) is supported by strong evidence that it will help prevent insomnia and improve sleep quality over time.

May 2017, the NCAA convened the Interassociation Task Force on Sleep and Wellness, which consisted of sleep experts, coaches, collegiate athletes, athletic administrators and educators who met to address this issue (online supplementary table 1). This meeting indicated the potential for collegiate athletics departments to help improve their athletes' sleep.

In this study, we aimed to obtain consensus on useful and feasible sleep-supportive strategies for collegiate athletics departments. Guided by the Delphi method, we began this process with a narrative review of the following topics: epidemiology of sleep in sport; sleep and optimal functioning among athletes; screening, tracking and assessment of athlete sleep; and interventions to improve sleep. In reviewing literature across these topic areas, the goals were to provide an overview for individuals working with athletes in collegiate sport and other sport settings and to provide a foundation for consensus recommendations regarding sleep for athletics departments.

METHODS

The Delphi method provides a comprehensive, efficient and unbiased way of aggregating expert opinion that allows multiple stakeholders to have a voice.¹⁰ Consistent with best practice guidance for using the Delphi method, our process included exploration and evaluation phases,¹⁰ as described below.

Exploration phase

Content experts conducted a narrative review of: (1) sleep patterns and disorders among collegiate athletes; (2) sleep and

optimal functioning among collegiate athletes; (3) screening, tracking and assessment of collegiate athlete sleep; and (4) interventions to improve collegiate athlete sleep. PubMed, Scopus and Cochrane CENTRAL databases were searched using these subject headings and keywords: 'Sleep', 'Sports', 'Injuries', 'Adolescents', 'Collegiate', 'Exercise', 'Mental Health', 'Performance', 'Athlete', 'Screening', 'Technology', 'Assessment', 'Tracking', 'Consumer', 'Subjective', 'Objective', 'Questionnaires', 'Sleep-Disordered Breathing', 'Insomnia', 'Narcolepsy', 'Circadian', 'Disorders' and 'Review'. Pearling of articles was also performed to find related references. Articles were chosen if they focused on sleep and collegiate athletes. For some topics, little or no published literature was available for collegiate athletes, but relevant literature was available for college students, emerging adults and competitive athletes in general. In such instances, we broadened our selection criteria to include literature focusing on these related populations. Studies published before 1 June 2018 were included and no study design limits were applied in any searches.

Evaluation phase

Based on the results of this review, an internal working group prepared foundational action-oriented recommendations for collegiate athlete sleep to implement in collegiate sport settings. Content experts contributing to the narrative review provided initial feedback about these recommendations to ensure scientific appropriateness and clarity. Recommendations were modified, as needed, based on this feedback. Next, members of a multidisciplinary panel group who attended the NCAA's Interassociation Task Force on Sleep and Wellness were emailed the narrative review and a link to an anonymous online survey for scoring the recommendations. Members of the panel included content experts (sleep researchers and individuals who develop and implement sleep-related policy and programming) and stakeholders affected by the recommendations (eg, coaches, collegiate athletes and athletic administrators).

Drawing on the narrative review and their own expertise, panel members scored each recommendation on utility and feasibility.^{10 11} *Utility* was defined as whether the recommendation, if implemented, was likely to improve any aspect of collegiate athlete sleep. *Feasibility* was defined as the degree to which individuals or campuses would be able to implement the recommendation. For each dimension, scores were elicited on a nine-point scale, with higher scores indicating that the recommendation is more useful/feasible. Panel members were also invited to provide written comments about each recommendation. After panel members completed this process, mean scores for each recommendation were calculated separately for utility and feasibility. Any recommendations with a rounded mean score of 3 or less on either dimension were discarded, and any with a rounded mean score of 7 or more on both dimensions were included in the final product. Recommendations that received a rounded mean score >3 but <7 on either dimension were revisited.¹¹

Panel members' comments were reviewed and potential modifications to the statements (or potential additional support for implementation to address feasibility) were generated based on these comments. Content experts were subsequently invited to provide additional feedback on the proposed modifications and implementation support. The internal working group synthesised this feedback and final modifications were made to the recommendations. As with the first round of scoring, panel members were emailed a link to an anonymous online survey in which they scored each recommendation that was modified

Table 1 National College Health Assessment: self-reported sleep behaviour among full-time undergraduate students who self-identify as varsity athletes at NCAA member Institutions (n=14 134)^{12(p20)}

Sleep behaviours	Reported response option	Percent of varsity athletes
Sleep difficulties in the past 12 months.	Traumatic or very difficult to handle.	24
Falling asleep at night.	Extreme difficulty at least three nights in the last week.	24
Awaking too early and unable to go back to sleep.	Occurred at least three nights in the last week.	16
Getting enough sleep to feel rested.	Insufficient sleep more than 3 days out of past 7;	57
	Insufficient sleep 6–7 days out of past 7.	23
Daytime tiredness.	Felt tired, dragged out or sleepy at least 3 days during the last week;	61
	reported that daytime sleepiness has been a big problem in last 7 days.	17
Negative consequences of sleep difficulties.	Academic performance.	18

or did not achieve consensus during the prior round of scoring. This second round of data were analysed as in the first round. Recommendations that the panel determined to be useful and feasible were presented as the final product of this process. NCAA Research Review Board approval was obtained for the Delphi consensus process.

NARRATIVE REVIEW

Epidemiology

The American College Health Association's annual National College Health Assessment (NCHA) survey includes information about sleep behaviour and allows researchers to aggregate results by students who self-identify as varsity athletes.¹² Among 14 134 collegiate athletes at NCAA member institutions, 61% reported daytime fatigue at least three or more days in the past week, and others reported consistent or more serious sleep difficulties (table 1). These data are similar to those in non-athletes,¹² suggesting lack of sleep is endemic among all college students.

In a recent single-institution study of 628 collegiate athletes from 29 varsity teams, 42% experienced poor sleep quality (measured by the Pittsburgh Sleep Quality Index) and 51% reported high levels of excessive daytime sleepiness (assessed by the Epworth Sleepiness Scale).¹³ Similarly, in the NCAA Growth, Opportunities, Aspirations and Learning of Students survey (GOALS),⁵ less than 25% of collegiate athletes reported ≥ 8 hours of sleep on a typical night (online supplementary table 2). Poor sleep quality was also prevalent, as 19% of male and 23% of female respondents reported difficulty sleeping at least 8 of the past 30 days (see online supplementary table 3 for a breakdown by NCAA division). Moreover, 70% of male and 82% of female collegiate athletes reported a preference for more sleep. In sum, epidemiological data suggest that inadequate sleep quality and quantity could be targeted for improvement among collegiate athletes.

Sleep and optimal functioning

Athletic performance

Data are limited regarding the influence of sleep disturbance, sleep deprivation or circadian desynchrony on athletic performance.^{8,9,14–16} Poor sleep is associated with delayed reaction time, increased perceived effort, fatigue, mood disturbance and reduced time to both exhaustion and task failure; however, it does not affect physiological markers of anaerobic, aerobic or power output.^{8,14,17} Interestingly, sleep-deprived athletes may be able to execute or perform required sport-related tasks, but their task-related cognitive and psychological tolerance is diminished, thereby increasing perceived effort and somatic symptom reporting.¹⁸ However, data on the overall physiological effects of

sleep impairment and its dose–response relationship on endurance performance, anaerobic power, and sprint performance are conflicting. The data are derived from small samples with inconsistent methods to measure strength and performance.⁸

As described in greater detail in a subsequent section, few studies have evaluated the effects of sleep extension on athletic performance. However, some studies have shown that serving accuracy, shooting accuracy and reaction time significantly improve in collegiate tennis and basketball players, respectively, following objectively measured sleep extension beyond 8 hours per day for several weeks.^{19,20} Although regular exercise improves sleep quality, duration and efficiency,²¹ the relationship between training loads, sleep and performance is complex and poorly understood. Sport-specific tasks and individual responses to training and sleep are likely different enough that optimising performance in athletes will require individualised approaches and further research.

Injury and illness

Adolescent athletes who reported sleeping < 8 hours per night were 1.7 times more likely to have a musculoskeletal injury,²² while sleeping > 8 hours during weekday nights was associated with 61% lower odds of such an injury.²³ Similarly, a simultaneous increase in training load, training intensity and decreased hours of sleep resulted in a significantly higher risk of injury.²⁴ College students with poor quality sleep also report significantly more medical illness than those with near-ideal or ideal sleep (as determined by an aggregation of factors including sleep quality and duration).²⁵ One potential mechanism may be that increases in proinflammatory cytokines following sleep loss could promote immune system dysfunction.²⁶

Mental health

Sleep disturbances are clinically relevant for evaluation and treatment of mental health conditions in collegiate athletes.^{6,27,28} The relationship between mental health and sleep quality are bidirectional and interdependent: poor sleep exacerbates mental health disorders and negative mood, and stress and mental health disorders lessen sleep quality and quantity.²⁹ The most commonly reported barrier to sleep among college students is perceived stress.²⁵ Among individuals with clinically relevant manifestations of depression and anxiety (25% of all collegiate athletes³⁰), sleep disturbances include difficulty with sleep initiation and maintenance, circadian rhythm dysregulation and insomnia or hypersomnia.³¹

In the 2018 American College Health Association (ACHA)-NCHA dataset, collegiate athletes with self-identified sleep problems were more than twice as likely to report feeling

overwhelming anxiety, being so depressed it was difficult to function and considering suicide in the last 30 days compared with athletes who did not indicate sleep disturbance.¹² Prospective population studies demonstrate that disturbed sleep predicts development of anxiety and depression in adolescents.³² In adults, anxiety and depression at baseline predict new cases of insomnia, and insomnia at baseline predicts new diagnoses of anxiety and depression 1 year later.³³ Similarly, treating insomnia improves depression and anxiety symptoms, and treating anxiety/depression improves insomnia.³⁴ In both students with depression and anxiety and healthy college students, better sleep is associated with improved mental wellness.^{35 36}

Academic performance

Better sleep is associated with greater academic success perhaps in part because acute sleep deprivation impairs cognitive performance and protracted insufficient sleep exacerbates mood disorders and distorts emotion regulation.^{28 37–39} In general, students with better sleep (eg, >7 hours of night-time sleep, more consistent sleep schedules, lower levels of daytime sleepiness and less weekend 'catch-up' sleep) report higher grade point averages (GPAs).^{28 37–39} Multiple single-institution prospective and cross-sectional studies have shown that insufficient, inconsistent and/or poor quality (non-restorative) sleep independently predict college students' academic performance, even after accounting for other measures of achievement like standardised test scores.^{28 37} In a study of 1845 undergraduates, those screening positive for possible sleep disorders were significantly more at risk for academic failure (GPA <2.0).⁴⁰ Among respondents in the 2009 ACHA-NCHA survey, sleep disturbances (early awakening, insufficient sleep or difficulty falling asleep) independently predicted poor academic performance, on par with high-risk drinking in its association with GPA and course completion. On average, each additional day per week a student reported experiencing sleep problems raised the probability of dropping a course by 10%.⁴¹

Screening, assessment and tracking

Sleep measurement can be either subjective, assessed mostly through questionnaires, or objective, occurring in either a sleep laboratory or at home (see online supplementary table 5). Objective testing can be further subdivided into 'clinical grade' measurements that are Food and Drug Administration (FDA) approved to assess and diagnose sleep disorders, or 'consumer-based',^{42 43} which lack FDA approval but can measure sleep continuously in the subject's typical sleeping environment. The clinical sleep laboratory environment, where detailed objective testing such as polysomnography⁴⁴ and multiple sleep latency testing⁴⁵ occur, has the advantage of accuracy and detail, and the disadvantage of an atypical sleeping environment, which can influence the sleep it intends to measure. Furthermore, this testing only occurs during a discrete period of time. Objective sleep assessment can also occur in the home environment, which has the advantage of familiarity and longitudinal measurement but has the disadvantage of limiting the number of possible measures and compromised accuracy compared with sleep laboratory measures.

Tools for screening, assessment and tracking

In general, subjective screening questionnaires^{44–58} and objective clinical-grade measurements^{44 45 57 59 60} focus on the diagnosis and treatment of sleep disorders such as sleep-disordered breathing (eg, obstructive sleep apnoea), insomnia, narcolepsy and circadian rhythm sleep disorders. The presence of these and

Box 2 Key issues to address in conducting a sleep history

1. Current sleep schedule on weekdays and weekends:
2. Bedtimes.
3. Wake times.
4. How long to fall asleep.
5. Napping (number, timing and duration).
6. Sleep quality:
 - a. Number of nightly awakenings and reason.
 - b. Presence of sleep onset or sleep maintenance insomnia (early morning awakening with trouble falling back asleep).
 - c. Wake with alarm or spontaneously.
 - d. Waking feeling rested.
 - e. Presence and extent of daytime sleepiness.
7. Sleep disorders:
 - a. Presence of hypnagogic or hypnopompic hallucinations, sleep paralysis or cataplexy.
 - b. Presence of restless leg symptoms (night-time leg discomfort relieved by movement).
 - c. Sleep walking, confusional arousals or night terrors.
 - d. Snoring, witnessed apnoeas, nocturnal choking and gasping or shortness of breath.
8. Medication and drug use:
 - a. Use of stimulants.
 - b. Use of sedative hypnotic medications.
 - c. Recreational drug and alcohol use.
9. Mental health status.
10. Environmental conditions:
 - a. Exposure to light.
 - b. Electronic device use.

other sleep disorders may be tested in collegiate athletes during their preparticipation physical exam or in those seeking diagnosis and treatment for sleep problems. Sports medicine team members can also use subjective measures (see online supplementary table 5) to refer collegiate athletes to team physicians, licenced mental health providers and board-certified sleep medicine professionals. These specialists can diagnose sleep disorders using clinical-grade objective measurements and implement treatment plans.

When screening athletes for disordered sleep, it is important to obtain a thorough sleep history. Key issues to address are summarised in box 2.⁶¹ Current sleep habits describe bedtimes; awakening history; napping; and exposure to light, dark and electronic device use. Sleep quality includes a subjective assessment of 'average sleep', feeling rested and presence of daytime fatigue or somnolence. Family medical history should screen for a history of insomnia, narcolepsy, sleep apnoea, depression, anxiety, attention deficit hyperactivity disorder or other psychiatric disorders (eg, somatoform disorders, substance abuse and personality disorders). A sleep diary can be a valuable tool for establishing baseline sleep habits and identifying changes in hygiene practices.

The emergence of consumer sleep technologies holds great promise for helping athletics departments better understand, and thus improve, collegiate athlete sleep.^{42 43} Their ability to measure sleep in the collegiate athletes' home environment over time provides a heretofore unattainable ecologically valid measure of sleep. Consumer sleep technologies can also identify problematic sleep that may indicate a sleep disorder. Despite the value of these technologies to coaches, team physicians and

collegiate athletes, their use assisting medical providers in diagnosing sleep disorders is evolving and not yet mature. However, they can currently help collegiate athletes and their support team to understand their sleep environment, duration and timing and can provide a baseline on which these collegiate athletes can improve.

Implementation of screening, assessment and tracking

One challenge in achieving collegiate athlete sleep health is determining the party responsible for screening, assessment and tracking of sleep beyond the athlete themselves. This group includes coaches, strength and conditioning specialists, team physicians, athletic trainers and psychologists. In some settings, strength and conditioning specialists are responsible for collecting subjective and objective collegiate athlete sleep data. They then relay the information and communicate with other clinicians/practitioners such as athletic trainers, team physicians and psychologists. Together, these professionals can provide the necessary support and expertise to the collegiate athlete to optimise his or her sleep health.

When considering the use of consumer sleep technologies, ethical issues of privacy, including who has access to the sleep data, and informed consent are hurdles that need to be cleared. Any sleep-monitoring programmes should be voluntary for collegiate athletes and include sleep health and performance education. Coaches and medical personnel should thoroughly investigate data supporting the accuracy and functionality of any consumer sleep technologies before collegiate athletes use them.

Interventions to improve sleep

Evidence-based approaches to improving sleep include behavioural, social and pharmacological interventions. Here we review how they apply to collegiate athletes, noting that in many instances the evidence base is primarily among college students more generally. Nearly three-quarters of college students report receiving no health education from their universities about how to manage sleep difficulties.¹² Interventions are particularly important for collegiate athletes who struggle to practice good sleep hygiene due to heightened time demands and for whom poor sleep impairs both academic and athletic performance. Unique elements of the collegiate athlete experience may present important opportunities for population-specific tailoring of educational messaging. For example, less than one-quarter of NCAA Division I collegiate athletes at a single institution regarded sleep as a way to recover from a game.⁶² Knowledge translation about sleep's positive impact on athletic performance is an important starting point for population-tailored educational interventions.

The most common non-pharmacological sleep interventions studied in college-aged populations^{63–65} include cognitive-behavioural therapy for insomnia (CBT-I), sleep hygiene and education and relaxation/mindfulness. In general, CBT-I has shown large effect sizes across numerous studies.⁶⁴ While CBT-I is generally focused on insomnia complaints, sleep hygiene and education interventions have been disseminated widely to college students. These interventions have generally resulted in small to moderate improvements in sleep quality⁶⁴ and sleep knowledge.⁶³ In addition, when used as stand-alone interventions, relaxation and mindfulness generally lead to moderate sleep improvements.^{64–66} Overall, sleep interventions in college students are associated with a full range of effect sizes, with increases in sleep duration and sleep efficiency and decreases in sleep onset latency and wake after sleep onset.⁶⁴

Among collegiate athletes in particular evidence is more limited. Two studies prescribed increased sleep duration to small samples of basketball players¹⁹ and tennis players.²⁰ One week²⁰ or 5–7 weeks¹⁹ of sleep extension (ie, increasing daily sleep duration to >9–10 hours) were associated with significant increases in sleep duration, decreases in sleepiness and improvements in sport-specific performance indices (eg, tennis serve accuracy and sprint times, basketball shooting accuracy and faster reaction times). However, these studies did not include long-term follow-up, so the persistence of improvements is unknown. Two other studies reported on interventions focused on sleep hygiene and health for collegiate athletes. A 1-month sleep health intervention (delivered via pamphlet) improved self-reported sleep and improved perceived soccer performance among soccer players.⁶⁷ In addition, preliminary analyses of a pilot 10-week sleep health intervention indicated significantly better sleep, energy levels and collegiate athlete mental health.⁶⁸ Several studies have examined the effects of sleep interventions in trained athletes, primarily young adults.⁶⁹ Short-term (ranging from a single night to 1 week) sleep hygiene interventions have modestly improved sleep duration but not other sleep parameters or markers of athletic performance.^{70–72} Improvement in perceived recovery was found in one study⁷¹ but not another.⁷⁰ Biofeedback did not improve sleep quality in a sample of soccer and track athletes,⁷³ and progressive muscle relaxation led to minimal improvements in sleep among dancers.⁷⁴ Short-term (48 hours) or long-term (4 weeks) restrictions on use of late-night electronic devices did not improve sleep^{75–76} or various markers of athletic and cognitive performance,⁷⁶ and five nights of sleep extension led to minimal improvements in measures of sleep or performance in adolescent pistol shooters.⁷⁷ In contrast, interventions involving comprehensive sleep education, feedback and tailored approaches to behaviour change over several weeks significantly improved sleep.^{78–79} Finally, in a sample of golfers with sleep apnoea, treatment with continuous positive airway pressure therapy significantly improved sleep quality, decreased reported sleepiness and lowered golf handicaps.⁸⁰

Although some collegiate athletics departments have modified their facilities or added programmes to improve sleep (eg, creation of napping rooms and comprehensive sleep monitoring systems), it remains unclear whether these strategies have improved collegiate athlete sleep. Overall, in addition to minimal research on non-pharmacological sleep interventions among collegiate athletes, the literature is greatly limited due to: small sample sizes, lack of control groups, insufficient sleep measures and minimal consideration for other relevant domains (eg, cognition, mental health, academic performance and sport performance).

Three per cent of NCAA collegiate athletes report non-prescription sleep aid use, while 18.7% of NCAA collegiate athletes report prescription sleep aid use.⁸¹ However, there is a dearth of literature on pharmacological treatment of insomnia and other sleep disorders in collegiate athletes; furthermore, hypnotic medication use among college students is broadly discouraged due to cognitive side effects and potential for adverse events when used with other drugs (eg, alcohol).⁶⁵ For these reasons, pharmacological recommendations as sleep aids were considered beyond the scope of the Interassociation Task Force on Sleep, and the reader is referred to the International Olympic Committee consensus statement on mental health in elite athletes for a detailed discussion of this matter.⁸² Given the broad concerns of medication aids for sleep in collegiate athletes, insomnia-related complaints in this population should be managed behaviourally, since CBT-I produces similar short-term treatment outcomes and

superior long-term treatment outcomes compared with pharmacotherapy.⁸³ Nevertheless, given the seemingly high prevalence of sedative-hypnotic use among collegiate athletes, more research on sleep medication use among collegiate athletes is needed to understand its sleep, health and performance-related implications.⁸⁴

CONSENSUS PROCESS

Based on the narrative review summarised above, the internal working group developed an initial list of 16 recommendations for collegiate athletics departments regarding collegiate athlete sleep (online supplementary table 6). Following the first round of voting, there was consensus on the utility of all 16 recommendations (scores of ≥ 7); however, six did not achieve consensus regarding feasibility. Descriptive statistics for these 16 initial recommendations are included in online supplementary table 5. Next, open-ended responses were reviewed for each recommendation, and key themes were identified. The most consistent theme related to feasibility was concern regarding lack of support for implementation. For example, an environmental assessment related to sleep barriers would be challenging without a readily available valid and reliable environmental assessment tool for institutions to use. Similarly, for on-campus implementation to occur, institutions would need a valid and reliable time demands survey (to assess quantitative and qualitative information on collegiate athlete commitments) and evidence-based educational/informational resources. A second emergent theme related to feasibility was how to provide institution-specific guidance when resources vary markedly across institutions (eg, personnel, funding and organisational structures). The internal working group reviewed these themes and revised the recommendations accordingly to improve feasibility. Implementation guidance for each recommendation was added, reviewed by the internal working group and content experts who contributed to the narrative review.

These revised recommendations and implementation guidance were returned to the panel for a second round of voting on utility and feasibility for the revised items. Open-ended feedback was elicited related to the implementation guidance for each recommendation, efficiently aggregating expert feedback about additional implementation resources and guidance accompanying each statement. All results of the second round of voting are provided in online supplementary table 7.

There was consensus that each revised recommendation was useful (rounded scores of ≥ 7). However, only 5 of the 10 second round recommendations achieved consensus on their feasibility in college sports settings. The final set of five recommendations is provided in **box 3** and include: (1) conducting an annual time demands survey, (2) incorporating sleep screening into the preparticipation exam, (3) providing evidence-based sleep education related to collegiate athletes and (4) coaches, and (5) ensuring that consumer sleep technology, if used, is compliant with Health Insurance Portability and Accountability Act and Family Educational Rights and Privacy Act laws.

CONCLUSION

Improving sleep in collegiate athletes has significant implications for health, athletic performance and academic achievement. However, athletes frequently fail to obtain restorative sleep. Factors related to their academic pursuits, training and competition can alter sleep patterns. Here we provide five recommendations related to collegiate athlete sleep that a panel of diverse experts considered useful and feasible for implementation in

Box 3 Recommendations with consensus utility and feasibility

- Recommendation 1: Conduct a collegiate athlete time demands survey annually.
- Recommendation 2: Ensure that consumer sleep technology, if used, is compliant with HIPAA and FERPA laws.
- Recommendation 3: Incorporate sleep screening into the preparticipation exam.
- Recommendation 4: Provide collegiate athletes with evidence-based sleep education that includes: (1) information on sleep best practices; (2) information about the role of sleep in optimising athletic and academic performance and overall well-being; and (3) strategies for addressing sleep barriers.
- Recommendation 5: Provide coaches with evidence-based sleep education that includes: (1) information on sleep best practices; (2) information about the role of sleep in optimising athletic and academic performance and overall well-being; and (3) strategies to help optimise collegiate athlete sleep.

FERPA, Family Educational Rights and Privacy Act; HIPAA, Health Insurance Portability and Accountability Act.

collegiate athletics departments. The Delphi process underscored the need for continued focus on developing and evaluating valid and reliable tools for sleep assessment and effective behavioural interventions among athletes and to support implementation of such tools and interventions in diverse sport environments. Such ongoing work can help contribute to the health and athletic performance of collegiate athletes and may have utility in other sport populations.

Acknowledgements We would like to thank all participants in the National Collegiate Athletic Association Sleep Task Force for their contributions at the Task Force meeting and to the Delphi voting process.

Collaborators National Collegiate Athletic Association Interassociation Task Force on Sleep and Wellness.

Contributors All authors have made substantial contributions to either conception and design or analysis and interpretation of data, drafting or revision of the article and have seen and given final approval of the submission.

Funding NCAA, Sport Science Institute.

Competing interests NFW reports consulting fees from SleepScore Labs, Jazz Pharmaceuticals and Harmony Biosciences. MG reports grants from Kemin Foods, grants from Nexalin Technology, personal fees from Fitbit, personal fees from Natrol LC, personal fees from Curaegis Technologies, personal fees from Thrive Global, outside the submitted work.

Patient consent for publication Not required.

Ethics approval National Collegiate Athletic Association Research Review Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data are available from the corresponding author on request.

REFERENCES

- 1 Sleep Health | Healthy People. 2020 <https://www.healthypeople.gov/2020/topics-objectives/topic/sleep-health> (Accessed 11 December, 2018).
- 2 Arnett JJ. Emerging adulthood: A theory of development from the late teens through the twenties. *Am Psychol* 2000;55:469–80.
- 3 Patrick Y, Lee A, Raha O, et al. Effects of sleep deprivation on cognitive and physical performance in university students. *Sleep Biol Rhythms* 2017;15:217–25.
- 4 Irish LA, Kline CE, Gunn HE, et al. The role of sleep hygiene in promoting public health: A review of empirical evidence. *Sleep Med Rev* 2015;22:23–36.
- 5 National Collegiate Athletic Association. *NCAA GOALS Study*. Indianapolis, IN: National Collegiate Athletic Association, 2016. <http://www.ncaa.org/about/resources/research/ncaa-goals-study>. (Accessed 11 December, 2018).

- 6 Hershner S, Chervin R. Causes and consequences of sleepiness among college students. *Nat Sci Sleep* 2014;6:73–84.
- 7 Prichard J, Cunningham B, Kiehlblock C, et al. The College Sleep Environmental Scan: Development and Initial Outcomes. Poster Presentation presented at the: Annual Meeting for Sleep Research Society; June 2016, Denver, CO. http://www.sleepmeeting.org/docs/default-source/default-document-library/sleep-39-as_final.pdf?sfvrsn=2
- 8 Fullagar HHK, Skorski S, Duffield R, et al. Sleep and Athletic Performance: The Effects of Sleep Loss on Exercise Performance, and Physiological and Cognitive Responses to Exercise. *Sports Medicine* 2015;45:161–86.
- 9 Gupta L, Morgan K, Gilchrist S. Does Elite Sport Degrade Sleep Quality? A Systematic Review. *Sports Medicine* 2017;47:1317–33.
- 10 Adler M, Ziglio E. Gazing Into the Oracle: The Delphi Method and Its Application to Social Policy and Public Health. *Jessica Kingsley Publishers* 1996.
- 11 Zumsteg JM, Cooper JS, Noon MS. Systematic Review Checklist. *J Ind Ecol* 2012;16(5):S12–S21.
- 12 American College Health Association. American College Health Association-National College Health Assessment, Fall 2015, Spring 2016, Fall 2016, Spring 2017, Fall 2017 [data file]. Hanover, MD: American College Health Association. [producer and distributor] 2018-11-15.
- 13 Mah CD, Kezirian EJ, Marcello BM, et al. Poor sleep quality and insufficient sleep of a collegiate student-athlete population. *Sleep Health* 2018;4:251–7.
- 14 Watson AM. Sleep and Athletic Performance. *Curr Sports Med Rep* 2017;16:413–8.
- 15 Thun E, Bjorvatn B, Flo E, et al. Sleep, circadian rhythms, and athletic performance. *Sleep Med Rev* 2015;23:1–9.
- 16 Simpson NS, Gibbs EL, Matheson GO. Optimizing sleep to maximize performance: implications and recommendations for elite athletes. *Scand J Med Sci Sports* 2017;27:266–74.
- 17 Taheri M, Arabameri E. The Effect of Sleep Deprivation on Choice Reaction Time and Anaerobic Power of College Student Athletes. *Asian J Sports Med* 2012;3:15–20.
- 18 Blumert PA, Crum AJ, Ernsting M, et al. The Acute Effects of Twenty-Four Hours of Sleep Loss on the Performance of National-Caliber Male Collegiate Weightlifters. *The Journal of Strength and Conditioning Research* 2007;21:1146–54.
- 19 Mah CD, Mah KE, Kezirian EJ, et al. The Effects of Sleep Extension on the Athletic Performance of Collegiate Basketball Players. *Sleep* 2011;34:943–50.
- 20 Schwartz J, Simon RD. Sleep extension improves serving accuracy: A study with college varsity tennis players. *Physiol Behav* 2015;151:541–4.
- 21 Kredlow MA, Capozzoli MC, Hearon BA, et al. The effects of physical activity on sleep: a meta-analytic review. *J Behav Med* 2015;38:427–49.
- 22 Milewski MD, Skaggs DL, Bishop GA, et al. Chronic Lack of Sleep is Associated With Increased Sports Injuries in Adolescent Athletes. *Journal of Pediatric Orthopaedics* 2014;34:129–33.
- 23 von Rosen P, Frohm A, Kottorp A, et al. Too little sleep and an unhealthy diet could increase the risk of sustaining a new injury in adolescent elite athletes. *Scand J Med Sci Sports* 2017;27:1364–71.
- 24 von Rosen P, Frohm A, Kottorp A, et al. Multiple factors explain injury risk in adolescent elite athletes: Applying a biopsychosocial perspective. *Scand J Med Sci Sports* 2017;27:2059–69.
- 25 Lund HG, Reider BD, Whiting AB, et al. Sleep Patterns and Predictors of Disturbed Sleep in a Large Population of College Students. *Journal of Adolescent Health* 2010;46:124–32.
- 26 Faraut B, Boudjeltia KZ, Vanhamme L, et al. Immune, inflammatory and cardiovascular consequences of sleep restriction and recovery. *Sleep Med Rev* 2012;16:137–49.
- 27 Gaultney JF. The Prevalence of Sleep Disorders in College Students: Impact on Academic Performance. *Journal of American College Health* 2010;59:91–7.
- 28 Taylor DJ, Bramoweth AD, Grieser EA, et al. Epidemiology of Insomnia in College Students: Relationship With Mental Health, Quality of Life, and Substance Use Difficulties. *Behav Ther* 2013;44:339–48.
- 29 Neckelmann D, Mykletun A, Dahl AA. Chronic Insomnia as a Risk Factor for Developing Anxiety and Depression. *Sleep* 2007;30:873–80.
- 30 Wolanin A, Hong E, Marks D, et al. Prevalence of clinically elevated depressive symptoms in college athletes and differences by gender and sport. *Br J Sports Med* 2016;50:167–71.
- 31 Lovato N, Grdisar M. A meta-analysis and model of the relationship between sleep and depression in adolescents: Recommendations for future research and clinical practice. *Sleep Med Rev* 2014;18:521–9.
- 32 Johnson EO, Roth T, Breslau N. The association of insomnia with anxiety disorders and depression: Exploration of the direction of risk. *J Psychiatr Res* 2006;40:700–8.
- 33 Jansson-Fröjmark M, Lindblom K. A bidirectional relationship between anxiety and depression, and insomnia? A prospective study in the general population. *J Psychosom Res* 2008;64:443–9.
- 34 Mason EC, Harvey AG. Insomnia before and after treatment for anxiety and depression. *J Affect Disord* 2014;168:415–21.
- 35 Boehm MA, Lei QM, Lloyd RM, et al. Depression, anxiety, and tobacco use: Overlapping impediments to sleep in a national sample of college students. *Journal of American College Health* 2016;64:565–74.
- 36 Milojevic HM, Lukowski AF. Sleep and Mental Health in Undergraduate Students with Generally Healthy Sleep Habits. *PLoS One* 2016;11:e0156372.
- 37 Singleton RA, Wolfson AR. Alcohol Consumption, Sleep, and Academic Performance Among College Students. *J Stud Alcohol Drugs* 2009;70:355–63.
- 38 Hartmann M, Prichard JR. Calculating the contribution of sleep problems to undergraduates' academic success and long-term economic outcomes. *Sleep* 2014.
- 39 Phillips AJK, Clerx WM, O'Brien CS, et al. Irregular sleep/wake patterns are associated with poorer academic performance and delayed circadian and sleep/wake timing. *Sci Rep* 2017;7.
- 40 Gaultney JF. Risk for Sleep Disorder Measured During Students' First College Semester May Predict Institutional Retention and Grade Point Average Over a 3-Year Period, With Indirect Effects Through Self-Efficacy. *Journal of College Student Retention: Research, Theory & Practice* 2016;18:333–59.
- 41 Hartmann M, Prichard JR. Calculating the contribution of sleep problems to undergraduates' academic success and long-term economic outcomes. *Sleep Health* 2018;4:463–71.
- 42 Khosla S, Deak MC, Gault D, et al. American Academy of Sleep Medicine Board of Directors. Consumer sleep technology: an American academy of sleep medicine position statement. *J Clin Sleep Med* 2018;14:877–80.
- 43 Ko PR, Kientz JA, Choe EK, et al. Consumer sleep technologies: A review of the landscape. *J Clin Sleep Med* 2015;11:1455–61.
- 44 Vaughn BV, Giallanza P. Technical Review of Polysomnography. *Chest* 2008;134:1310–9.
- 45 Littner MR, Kushida C, Wise M, et al. Practice Parameters for Clinical Use of the Multiple Sleep Latency Test and the Maintenance of Wakefulness Test. *Sleep* 2005;28:113–21.
- 46 Buysse DJ, Reynolds CF, Monk TH, et al. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193–213.
- 47 Parrott AC, Hindmarch I. Factor analysis of a sleep evaluation questionnaire. *Psychol Med* 1978;8:325–9.
- 48 Bastien C, Vallières A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med* 2001;2:297–307.
- 49 Chung F, Subramanyam R, Liao P, et al. High STOP-Bang score indicates a high probability of obstructive sleep apnoea. *Br J Anaesth* 2012;108:768–75.
- 50 Hoddes E, Zarcone V, Smythe H, et al. Quantification of Sleepiness: A New Approach. *Psychophysiology* 1973;10:431–6.
- 51 Kaida K, Takahashi M, Åkerstedt T, et al. Validation of the Karolinska sleepiness scale against performance and EEG variables. *Clinical Neurophysiology* 2006;117:1574–81.
- 52 Johns MW. A New Method for Measuring Daytime Sleepiness: The Epworth Sleepiness Scale. *Sleep* 1991;14:540–5.
- 53 Dittner AJ, Wessely SC, Brown RG. The assessment of fatigue: a practical guide for clinicians and researchers. *J Psychosom Res* 2004;56:157–70.
- 54 Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol* 1976;4:97–110.
- 55 Kallus W, Kellmann M. The Recovery-Stress Questionnaires: User Manual. *Champaign, IL, US: Human Kinetics* 2001;2001.
- 56 Montgomery-Downs HE, Insana SP, Bond JA. Movement toward a novel activity monitoring device. *Sleep and Breathing* 2012;16:913–7.
- 57 Morgenthaler T, Alessi C, Friedman L, et al. Practice Parameters for the Use of Actigraphy in the Assessment of Sleep and Sleep Disorders: An Update for 2007. *Sleep* 2007;30:519–29.
- 58 Monk TH, Reynolds CF, Kupfer DJ, et al. The Pittsburgh Sleep Diary. *J Sleep Res* 1994;3:111–20.
- 59 Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med* 2017;13:479–504.
- 60 Collop NA, Tracy SL, Kapur V, et al. Obstructive sleep apnea devices for out-of-center (OOC) testing: technology evaluation. *J Clin Sleep Med* 2011;7:531–48.
- 61 Kapur VK. Approach to the Patient with a Sleep Complaint. In: *Clinician's Guide to Sleep Disorders*. New York, NY, USA: Taylor and Francis, 2006.
- 62 Murray A, Fullagar H, Turner AP, et al. Recovery practices in Division 1 collegiate athletes in North America. *Phys Ther Sport* 2018;32:67–73.
- 63 Dietrich SK, Francis-Jimenez CM, Knibbs MD, et al. Effectiveness of sleep education programs to improve sleep hygiene and/or sleep quality in college students: a systematic review. *JBI Database System Rev Implement Rep* 2016;14:108–34.
- 64 Friedrich A, Schlarb AA. Let's talk about sleep: a systematic review of psychological interventions to improve sleep in college students. *J Sleep Res* 2018;27:4–22.
- 65 Kloss JD, Nash CO, Horsey SE, et al. The Delivery of Behavioral Sleep Medicine to College Students. *Journal of Adolescent Health* 2011;48:553–61.
- 66 Caldwell K, Emery L, Harrison M, et al. Changes in Mindfulness, Well-Being, and Sleep Quality in College Students Through Taijiquan Courses: A Cohort Control Study. *The Journal of Alternative and Complementary Medicine* 2011;17:931–8.
- 67 Harada T, Wada K, Tsuji F, et al. Intervention study using a leaflet entitled "three benefits of "go to bed early! get up early! and intake nutritionally rich breakfast!" a message for athletes" to improve the soccer performance of university soccer team. *Sleep Biol Rhythms* 2016;14:65–74.
- 68 Grandner MA, Athey A, Killgore WD, et al. 0793 Preliminary results of a sleep health intervention in student athletes: Changes in sleep, energy level, and mental well-being, and body weight. *Sleep* 2017;40:A294.

- 69 Bonnar D, Bartel K, Kakoschke N, *et al.* Sleep Interventions Designed to Improve Athletic Performance and Recovery: A Systematic Review of Current Approaches. *Sports Medicine* 2018;48:683–703.
- 70 Fullagar H, Skorski S, Duffield R, *et al.* The effect of an acute sleep hygiene strategy following a late-night soccer match on recovery of players. *Chronobiol Int* 2016;33:490–505.
- 71 Duffield R, Murphy A, Kellett A, *et al.* Recovery from Repeated On-Court Tennis Sessions: Combining Cold-Water Immersion, Compression, and Sleep Interventions. *Int J Sports Physiol Perform* 2014;9:273–82.
- 72 O'Donnell S, Driller M. Sleep-hygiene education improves sleep indices in elite female athletes. *International Journal of Exercise Science* 2017;10:522–30.
- 73 Rijken NH, Soer R, de Maar E, de M, *et al.* Increasing Performance of Professional Soccer Players and Elite Track and Field Athletes with Peak Performance Training and Biofeedback: A Pilot Study. *Appl Psychophysiol Biofeedback* 2016;41:421–30.
- 74 McCloughan LJ, Hanrahan SJ, Anderson R, *et al.* Psychological recovery: Progressive muscle relaxation (PMR), anxiety, and sleep in dancers. *Perform Enhanc Health* 2016;4:12–17.
- 75 Dunican IC, Martin DT, Halson SL, *et al.* The Effects of the Removal of Electronic Devices for 48 Hours on Sleep in Elite Judo Athletes. *J Strength Cond Res* 2017;31:2832–9.
- 76 Harris A, Gundersen H, Mørk-Andreassen P, *et al.* Restricted use of electronic media, sleep, performance, and mood in high school athletes—a randomized trial. *Sleep Health* 2015;1:314–21.
- 77 Suppiah HT, Low CY, Choong GCW, *et al.* Restricted and unrestricted sleep schedules of Asian adolescent, high-level student athletes: effects on sleep durations, marksmanship and cognitive performance. *Biol Rhythm Res* 2016;47:505–18.
- 78 Van Ryswyk E, Weeks R, Bandick L, *et al.* A novel sleep optimisation programme to improve athletes' well-being and performance. *Eur J Sport Sci* 2017;17:144–51.
- 79 Tuomilehto H, Vuorinen V-P, Penttälä E, *et al.* Sleep of professional athletes: Underexploited potential to improve health and performance. *J Sports Sci* 2017;35:704–10.
- 80 Benton ML, Friedman NS. Treatment of obstructive sleep apnea syndrome with nasal positive airway pressure improves golf performance. *J Clin Sleep Med* 2013;9:1237–42.
- 81 NCAA. *National Study of Substance Use Habits of College Student Athletes*, 2018.
- 82 Reardon CL, Hainline B, Aron CM, *et al.* International olympic committee consensus statement on mental health in elite athletes. *Br J Sports Med* 2019.
- 83 Smith MT, Perlis ML, Park A, *et al.* Comparative meta-analysis of pharmacotherapy and behavior therapy for persistent insomnia. *Am J Psychiatry* 2002;159:5–11.
- 84 Taylor L, Christmas BCR, Dascombe B, *et al.* Sleep medication and athletic performance—the evidence for practitioners and future research directions. *Front Physiol* 2016;7:83.
- 85 Monk TH, Reynolds CF, Kupfer DJ, *et al.* The pittsburgh sleep diary. *J Sleep Res* 1994;3:111–20.
- 86 Valko PO, Bassetti CL, Bloch KE, *et al.* Validation of the fatigue severity scale in a Swiss cohort. *Sleep* 2008;31:1601–7.
- 87 Chung F, Abdullah HR, Liao P. STOP-bang questionnaire: a practical approach to screen for obstructive sleep apnea. *Chest* 2016;149:631–8.
- 88 Kapur VK, Auckley DH, Chowdhuri S, *et al.* Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea: an American academy of sleep medicine clinical practice guideline. *J Clin Sleep Med* 2017;13:479–504.
- 89 S+ sleep monitoring device | ResMed.com. <https://www.resmed.com/us/en/consumer/s-plus.html> (Accessed 11 December, 2018).
- 90 Beattie Z, Oyang Y, Statan A, *et al.* Estimation of sleep stages in a healthy adult population from optical plethysmography and accelerometer signals. *Physiol Meas* 2017;38:1968–79.