Which criteria are used to clear patients to return to sport after primary ACL reconstruction? A scoping review

Ciara R Burgi,¹ Scott Peters,² Clare L Ardern,^{3,4} John R Magill,¹ Christina D Gomez,⁵ Jonathan Sylvain,⁶ Michael P Reiman⁷

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2018-099982>).

¹Department of Physical Therapy and Occupational Therapy, Duke University Health System, Durham, North Carolina, USA

²Physical Therapy, Toronto Blue Jays Major League Baseball Club, Toronto, Ontario, Canada

³Division of Physiotherapy, Linköping University, Linköping, Sweden

⁴School of Allied Health, La Trobe University, Melbourne, Victoria, Australia

⁵Department of Orthopedics and Sports Medicine, Houston Methodist Sugar Land Hospital, Sugar Land, Texas, USA

⁶Hartford Healthcare Rehabilitation Network, Farmington, Connecticut, USA

⁷Division of Physical Therapy, Duke University Medical Center, Durham, North Carolina, USA

Correspondence to

Dr Ciara R Burgi, Duke Sports Medicine Physical Therapy, Durham, NC 27710, USA; ciara.burgi@gmail.com

Accepted 5 January 2019

Published Online First
2 February 2019

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Burgi CR, Peters S, Ardern CL, et al. *Br J Sports Med* 2019;**53**:1154–1161.

ABSTRACT

Objective To describe the criteria used to clear athletes to return to sport (RTS) following primary ACL reconstruction.

Design Scoping review.

Data sources MEDLINE, Embase, CINAHL and SPORTDiscus electronic databases were searched using keywords related to ACL and RTS.

Eligibility criteria Prospective or retrospective studies reporting at least one RTS criterion for athletes who had primary ACL reconstruction with an autograft.

Results In total, 209 studies fulfilled the inclusion criteria. RTS criteria were categorised into six domains: time, strength, hop testing, clinical examination, patient-report and performance-based criteria. From the 209 included studies, time was used in 178 studies (85%), and in 88 studies (42%) was the sole RTS criterion. Strength tests were reported in 86 studies (41%). Sixteen different hop tests were used in 31 studies (15%). Clinical examination was used in 54 studies (26%), patient report in 26 studies (12%) and performance-based criteria in 41 studies (20%).

Summary Time and impairment-based measures dominated RTS criteria, despite sport being a complex physical and biopsychosocial activity with demands across all aspects of function. Time was included as a criterion in 85% of studies, and over 80% of studies allowed RTS before 9 months. Whether RTS tests are valid—do they predict successful RTS?—is largely unknown.

INTRODUCTION

When should an athlete return to sport (RTS) after an ACL reconstruction?^{1–2} Evidence-based guidelines^{2–4} advocate for a test battery to evaluate quality and quantity of movement with a limb symmetry index (LSI) of at least 90%. However, limb symmetry does not indicate the athlete has reached the preinjury level of function, the preinjury function may be insufficient for safe sports participation and performance, and qualitative asymmetries can exist despite quantitative symmetry.^{5–6} Inconsistent and contradictory recommendations, which can arise when using a time-based as opposed to a criterion-based protocol, challenge practitioners and athletes when they are working together to plan RTS. It is unclear whether RTS testing practices reflect recommendations for a multifactorial, multidisciplinary approach to RTS decision making.^{7–9} The first step in establishing guidance on RTS testing is to understand how such testing is approached in the published literature.

A previous review¹⁰ describing the criteria used to determine unrestricted participation in sport was limited in at least four ways: (1) only one electronic database (MEDLINE) was searched; (2) only studies published between April 2001 and April 2011 were eligible for inclusion; (3) the population of interest was not clearly defined; and (4) the International Classification of Functioning, Disability and Health (ICF) criteria were not addressed.¹¹ The ICF framework reflects an athlete-centred approach to understanding the current state of RTS testing.

The primary aim of this scoping review was to describe the criteria used in published research to clear athletes to RTS following primary ACL reconstruction. The secondary aim was to summarise the historical trends of RTS criteria and the current status of RTS testing.

METHODS

A scoping review design and methodology was used due to the exploratory nature of the research question. Scoping reviews are recommended when the aim is to collate and comprehensively summarise the literature on topics of a substantial and varied nature.^{12–13} Due to the descriptive and exploratory nature of scoping reviews, study quality and risk of bias assessments are not appropriate and do not influence scoping review outcomes.^{12–13}

Identification and selection of studies

We conducted librarian-assisted computer searches of the MEDLINE, CINAHL, Embase and SPORTDiscus electronic databases for studies on discharge criteria for RTS following primary ACL reconstruction. The search was performed on 12 September 2017 with no date limit. The goal was to maximise the sensitivity of our search strategy,^{14–15} increasing the likelihood that all appropriate studies were identified. The search was limited to humans and English language, peer-reviewed publications. The search strategies for MEDLINE, CINAHL, Embase and SPORTDiscus are listed in online appendix I. The reference lists of relevant systematic reviews identified during the title and abstract screen were also hand searched to identify any potentially eligible articles that may have been missed in the electronic database search.¹⁶

Inclusion in this review was based on:

- **Study design:** prospective or retrospective intervention or observational studies with a population greater than 10 athletes (to avoid small case series and case reports that may not

Systematic review

best represent RTS after ACL reconstruction) published in English language. Systematic reviews, conference abstracts, case studies, narrative reviews and non peer-reviewed studies were excluded.

- **Participants:** athletes (as defined by each study) of any age or competition level who had undergone primary ACL reconstruction with an autograft. At least 80% of participants had to be an athlete for the study to be included for review. We set a minimum of 80% athletes to maximise the chance that participants desired to RTS. We defined non-athletes as those who did not participate in sport. If studies did not define the population of interest as athletes but stated that patients returned to sport, we assumed they were athletes. Athletes with concomitant surgical procedures (eg, meniscus resection or repair, collateral ligament repair, posterior cruciate ligament reconstruction or repair) performed in combination with ACLR were included provided the ACL reconstruction was the primary procedure (80% of surgical procedures performed in the study).
- **Intervention:** at least 80% of athletes in the study underwent ACL reconstruction with an autograft. Studies where allografts, ACL repair or other graft sources were used in a primary surgical procedure, and studies where ACL reconstruction was not the primary procedure performed were excluded.
- **Outcomes:** any criterion used for clearance to RTS.
- **Time:** all postsurgical time frames reporting RTS criteria were included. There were no limits to year of study publication.

All references were downloaded to the Rayyan reference management platform (rayyan.qcri.org), and duplicates were removed prior to screening. Two authors independently screened titles, abstracts and reviewed articles in full text. During title and abstract screening, if at least one reviewer concluded that a study met the selection criteria or if it was unclear whether the study should be included or excluded, the study was included. Discrepancies between reviewers after full-text screening were resolved via consensus or discussion with a third reviewer for final inclusion.

Data extraction

We extracted the criteria used for clearance to RTS and general demographic and surgical data (table 1). Data elements were extracted to a custom Excel spreadsheet by one reviewer and verified by a second independent reviewer. Disagreements were resolved via consensus or discussion with a third reviewer. For the majority of studies, RTS criteria were not the main focus of the study. Therefore, RTS criteria were most commonly reported in the methods section. In studies where RTS criteria were the main focus of the study, criteria were reported in the results section.

Table 1 Variables extracted for this scoping review

Demographic and study information	Subjective and objective measures
Year	Anthropometry
Country of study origin	Range of motion
Sample size	Strength
Demographics (age and sex)	Ligament testing
Sport at time of injury	Patient-reported outcome measures
Level of sport participation	Functional testing

Table 2 Return to sport criteria categorisation and associated International Classification of Functioning, Disability and Health (ICF)¹⁷ framework classification

Return to sport criteria categorisation	Corresponding ICF ¹⁷ framework classification
Time	N/A
Strength	Impairments
Clinical examination	Impairments
Hop testing	Activity (limitations)
Performance-based criteria	Participation (restrictions)
Patient report	Contextual factors

Data management

Demographic data were summarised as frequencies, or arithmetic mean and SD or median and IQR, as appropriate (Microsoft Excel, Microsoft Corporation, Santa Rosa, California). After data extraction, we used consensus to identify six main RTS criteria categories based on consensus of the logical grouping of criteria: time, strength, hop testing, patient-report, clinical examination and performance-based criteria (table 2). We further classified these categories (and associated RTS criteria)—reflecting the ICF¹⁷—as impairment, activity, participation, contextual and time criteria (table 2). RTS criteria were summarised as frequencies and presented in summary tables. To summarise time trends in RTS criteria, criteria were binned by year of study publication 1986 to 2017 and presented as a relative proportion of the total criteria reported in each year.

RESULTS

The electronic database search yielded 4930 records. We included 209 for qualitative review after title, abstract and full-text review (figure 1). Studies that were excluded after full-text review are reported in online appendix II.

Participants

This scoping review summarises the results of RTS testing of over 22 000 participants with primary ACL reconstruction. Approximately two-thirds of participants were men, and the mean time to RTS was 7.6 months (table 3).

RTS criteria after ACL reconstruction

The previous review of RTS criteria¹¹ included 159 studies reporting RTS criteria, and due to minor differences in selection criteria, 113 were eligible for inclusion in our scoping review. There were 96 articles published since the 2011 review¹¹ that were eligible for inclusion in our scoping review.

Time

Time to RTS was reported as a criterion in 178 of the 209 included studies (85%) (table 4). Time was the sole criterion used to clear athletes to RTS in 88 studies (42%). Thirty-one studies (15%) did not use time as a RTS criterion. Eighteen studies (8%) reported a time frame for RTS clearance; seven studies^{18–24} (3%) reported a range (eg, RTS at 6–12 months) and 11 studies^{24–34} (5%) reported different RTS times based on type of sport (eg, non-contact at 4 months and contact at 6 months).

Strength tests (impairment measures)

Of the 209 included studies, 86 (41%) included strength as a RTS criterion (table 5). Two studies^{35 36} (1%) used isokinetic and isometric testing measures. Forty per cent of the 50 studies that

Figure 1 Study search flow diagram. RTS, return to sport.

Table 3 Demographic characteristics of included studies

		Studies reporting (% of 209)
Patients (n)	22 009	
Sex (proportion of men)	65%	94
Athlete level (n=studies reporting)		57
Amateur	97	
Professional	3	
Mixed	18	
Not reported	91	
Preinjury sport participation		
Soccer		23
Basketball		14
Volleyball		7
American football		5
All other sports		21
Mean age (SD), years	25.7 (5.3)	96
Median (IQR) time from injury to surgery, months	10.2 (4.1–17.9)	45
Median (IQR) postsurgical final follow-up time, months	29.5 (22.1–43.9)	56
Mean (SD) time to commence running, months	3.0 (1.0)	55
Mean (SD) time to RTS, months	7.6 (1.6)	16

used isokinetic strength as part of RTS criteria required a LSI of at least 85% for clearance to RTS.

Hop tests (activity measures)

From the 209 included studies, 30 (14%) reported using at least one hop test as a RTS criterion (table 6). Of these 30 studies, 22 (73%) required a LSI of at least 85% for clearance to RTS.

Table 4 Time as a RTS criterion

	Studies reporting, n (%)
Time criterion	178 (85 of 209)
<6 months from ACLR	43 (24 of 178)*
≥6 months to <9 months	129 (72)*
≥9 months to <12 months	18 (10)*
≥12 months	6 (3)*
Time as a single criterion	88 (49 of 178)
Time as part of a RTS test battery	90 (51 of 178)
Studies not using time in RTS test battery	31 (15 of 209)

*Numbers sum to >178 because 18 studies reported a RTS time frame and were counted in the each of the categories the time frame spanned. References in online appendix III.

ACLR, ACL reconstruction; RTS, return to sport.

Systematic review

Table 5 Strength as a RTS criterion

	Studies reporting, n (%)
Strength tests used as RTS criteria	86 (41 of 209)
Isokinetic testing	50 (58 of 86)
LSI $\geq 90\%$	11 (22 of 50)
Quadriceps and hamstring strength at 60°/s	1 (2 of 50)
Quadriceps and hamstring strength at 180°/s	1 (2)
Quadriceps and hamstring strength at 60°/s and 180°/s	2 (4)
Quadriceps and hamstring strength at 90°/s and 180°/s	1 (2)
Quadriceps and hamstring strength at 180°/s and 300°/s	1 (2)
Quadriceps and hamstring strength at 60°/s, 180°/s and 240°/s	1 (2)
Quadriceps and hamstring strength at 60°/s, 180°/s and 300°/s	2 (4)
Strength (parameters not stated)	2 (4)
LSI $\geq 85\%$	9 (18 of 50)
Quadriceps and hamstring strength at 60°/s	1 (2 of 50)
Quadriceps and hamstring strength at 120°/s	1 (2)
Quadriceps and hamstring strength at 60°/s and 180°/s	2 (4)
Quadriceps and hamstring strength at 90°/s and 240°/s	1 (2)
Quadriceps and hamstring strength (parameters not stated)	4 (8)
LSI $\geq 80\%$	11 (22 of 50)
Quadriceps and hamstring strength at 60°/s	3 (6 of 50)
Quadriceps and hamstring strength at 90°/s	2 (4)
Quadriceps and hamstring strength at 60°/s and 180°/s	1 (2)
Quadriceps and hamstring strength at 80°/s and 160°/s	1 (2)
Quadriceps strength at 60°/s, 180°/s and 240°/s	2 (4)
Quadriceps and hamstring strength (parameters not stated)	2 (4)
LSI $\geq 65\%$	1 (2 of 50)
Quadriceps strength	1 (2 of 50)
Multiple LSI criteria	6 (12 of 50)
Quadriceps LSI $\geq 85\%$; hamstring LSI $\geq 90\%$; H/Q ratio $\geq 70\%$ at 60°/s, 180°/s and 300°/s	1 (2 of 50)
Quadriceps and hamstring LSI $\geq 90\%$ at 60°/s, 180°/s and 300°/s; normalised isokinetic quadriceps strength ≥ 3 Nm/kg at 60°/s; H/Q ratio $\geq 55\%$ for females and $\geq 62.5\%$ for males at 300°/s	1 (2)
Quadriceps and hamstring LSI $\geq 90\%$ at 60°/s, 180°/s and 300°/s; H/Q ratio $\geq 55\%$ for females and $\geq 62.5\%$ for males at 300°/s	1 (2)
Quadriceps strength of LSI $\geq 90\%$ and hamstring strength of LSI=100% at 60°/s and 300°/s	1 (2)
'Good muscular... recovery' of quadriceps and hamstring strength; H/Q ratio between 40%–80% (parameters not stated)	1 (2)
Eccentric or concentric hamstring strength LSI $\geq 80\%$ and quadriceps strength of LSI=100%	1 (2)
Criteria not clearly stated	12 (24 of 50)
Quadriceps and hamstring strength at 60°/s	3 (2 of 50)
Quadriceps and hamstring strength at 180°/s	1 (2)
Quadriceps and hamstring strength at 60°/s and 180°/s	1 (2)
Quadriceps and hamstring strength at 60°/s and 240°/s	2 (4)
Quadriceps and hamstring strength at 90°/s and 230°/s	2 (4)
Quadriceps and hamstring strength at 180°/s and 230°/s	1 (2)
Quadriceps and hamstring strength at 60°/s, 120°/s and 180°/s	1 (2)
Quadriceps and hamstring strength at 60°/s, 180°/s, 300°/s; internal rotation and external rotation strength at 60°/s, 120°/s and 180°/s	1 (2)
Isometric testing	8 (9 of 86)

Continued

Table 5 Continued

	Studies reporting, n (%)
LSI $\geq 90\%$	5 (63 of 8)
Quadriceps strength at 60° of knee flexion	1 (13 of 8)
Quadriceps strength at 90° of knee flexion	1 (13)
Quadriceps burst superimposition strength	2 (25)
Isometric leg press testing	1 (13)
LSI $\geq 85\%$	1 (13 of 8)
Quadriceps and hamstring strength at 30° of knee flexion	1 (13 of 8)
LSI $\geq 80\%$	2 (25 of 8)
Quadriceps strength at 90° of knee flexion	1 (13 of 8)
Hamstring strength at 70° and 90° of knee flexion in prone and supine positions	1 (13)
Method of testing not clearly stated	25 (29 of 86)
Full strength	1 (1 of 86)
Close to full strength	2 (2)
Quadriceps strength of LSI $\geq 90\%$	2 (2)
Muscle strength of LSI $\geq 90\%$	2 (2)
Quadriceps and hamstring strength of LSI $\geq 90\%$	2 (2)
Quadriceps and hamstring strength of LSI $\geq 85\%$	1 (1)
Quadriceps strength of LSI $\geq 85\%$	2 (2)
Thigh strength of LSI $\geq 85\%$	1 (1)
Quadriceps strength of LSI $\geq 80\%$	1 (1)
Quadriceps strength of LSI $\geq 80\%$ and normal hamstring strength	1 (1)
Adequate quadriceps and hamstring strength	1 (1)
Good quadriceps function	1 (1)
Adequate strength	2 (2)
Functional strength	1 (1)
Proper muscular conditioning	1 (1)
Complete muscular recovery	1 (1)
Full functional stability in terms of strength	1 (1)
Regained strength	1 (1)
Leg press test demonstrating comparable strength to uninvolved lower extremity	1 (1)
Predetermined benchmark	5 (6 of 86)
Leg press for >15 reps at 100% BW of affected limb	5 (6 of 86)

References in online appendix III.

BW, body weight; H/Q ratio, hamstring/quadriceps ratio; LSI, limb symmetry index ; BW, body weight; H/Q ratio, hamstring/quadriceps ratio; RTS, return to sport.

Patient-report (contextual factors)

Patient-report criteria were used in 26 (12%) of the 209 studies and included the subcategories patient-reported outcome measures (PROMs), subjective statements and reports of pain (table 7). Six studies^{37–42} (3%) used one outcome measure, and two studies^{43–44} (1%) used two measures. One study⁴⁵ reported pain and subjective reports as RTS criteria and one study⁴⁰ used PROMs and subjective statements as RTS criteria.

Clinical examination (impairment measures)

Fifty-four studies (26%) reported completing a clinical examination as part of RTS testing (table 7). Tests of ligament stability, range of motion, effusion and thigh circumference were included as part of different clinical examinations.

Thigh circumference

Thigh circumference was reported in three studies (1%) with defined criteria of ≤ 1 cm^{46–47} or ≤ 2 cm.⁴⁸

Table 6 Hop performance as a RTS criterion

	Studies reporting, n (%)
Hop tests used as RTS criteria	30 (14 of 209)
Single leg hop	28 (93 of 30)
LSI $\geq 90\%$	17 (60 of 28)
LSI $\geq 85\%$	5 (18)
LSI $\geq 80\%$	2 (7)
LSI not stated	4 (14)
Triple hop	18 (60 of 30)
LSI $\geq 90\%$	14 (78 of 18)
LSI $\geq 85\%$	2 (11)
LSI not stated	2 (11)
Crossover hop	11 (37 of 30)
LSI $\geq 90\%$	9 (82 of 11)
LSI $\geq 85\%$	2 (18)
Timed 6 m hop	11 (37 of 30)
LSI $\geq 90\%$	8 (72 of 11)
LSI $\geq 85\%$	2 (18)
LSI $\geq 80\%$	1 (9)
Other hop tests	11 (37 of 30)
Side hop test	2 (18 of 11)
Single leg vertical jump	4 (36)
Timed lateral hop	1 (9)
Single leg 10 m speed jump test	1 (9)
Square hop	1 (9)
Side hop from 30 cm box	1 (9)
Forward hop from 30 cm box	1 (9)
Two-leg and one-leg stability tests, two-leg and one-leg countermovement jump, speedy jumps, plyometric jumps and quick feet test	2 (18)

Two studies used seven different tests, 11 studies used four hop tests, 7 studies used three tests, 3 studies used two hop tests and 7 studies used one hop test. References in online appendix III.

LSI, limb symmetry index; RTS, return to sport.

Ligamentous stability

Twenty-five studies (6%) used at least one laxity test (table 7). Two studies^{48 49} defined a criterion of <3 mm side-to-side difference for clearance to RTS.

Range of motion (ROM)

Thirty studies (14%) used knee ROM as a RTS criterion (table 7). Nine studies^{48 50–57} required full range of flexion and extension, and four studies^{31 32 58 59} required only full extension. 'Full ROM' was not consistently defined. One study⁵² examined active and passive range of motion, and two studies^{57 60} examined passive range of motion only.

Effusion

Twenty studies (10%) used effusion as a RTS criterion (table 7). One study used the stroke test,⁵⁸ and one study used a measurement of suprapatellar circumference.⁶¹

Performance-based criteria (impairment, activity and participation measures)

From the 209 included studies, 41 (20%) used performance-based criteria (tests of agility, proprioception, aerobic endurance, quality of motion, reported completion of a specific programme and non-specific performance-based criteria) to clear athletes to RTS (table 7). The 28 studies (13%) requiring completion of

Table 7 Additional measures used as criteria for clearance to return to sport

	Studies reporting, n (%)	ICF domain
Patient-report	26 (12 of 209)	
Patient-reported outcome measures	8 (31 of 26)	Contextual
IKDC	5 (63 of 8)	
Within 15% of gender aged-matched controls	3 (38)	
Composite score of 90%	1 (13)	
Question #10: ≥ 9 points	1 (13)	
KOS-ADL: score $\geq 90\%$	3 (38)	
ACL-RSI: score $\geq 56\%$	1 (13)	
Global rating of knee function: score $\geq 90\%$	1 (13)	
Subjective statements	9 (35 of 26)	
'Full subjective functional stability'	4 (44 of 9)	
When athlete felt comfortable	2 (22)	
Full subjective confidence in knee	2 (22)	
'Desire to return to previous activity'	1 (11)	
Pain	11 (42 of 26)	
'No pain'	7 (64 of 11)	
'No problematic symptoms'	3 (27)	
'<3/10 pain'	1 (9)	
Clinical examination	54 (26 of 209)	Impairments
Ligament stability	35 (65 of 54)	
KT-1000	8 (23 of 35)	
Lachman test	6 (17)	
Pivot shift	3 (9)	
Anterior drawer	1 (3)	
Stability testing method not stated	22 (63)	
Range of motion (ROM)	30 (56 of 54)	
Full/complete ROM	25 (83 of 30)	
Less than full ROM	2 (7)	
ROM equal to contralateral leg	2 (7)	
Adequate ROM	1 (3)	
Effusion	20 (37 of 54)	
No effusion	18 (90 of 20)	
Minimal effusion	2 (10)	
Thigh circumference	3 (6 of 54)	
Atrophy ≤ 1 cm	2 (67 of 3)	
≤ 2 cm difference in thigh circumference	1 (33)	
'Satisfactory clinical exam'	2 (4 of 54)	
Performance-based criteria	41 (20 of 209)	
Agility	4 (10 of 41)	Activities
T-test	2 (50 of 4)	
'Good agility'	2 (50)	
Proprioception	6 (15 of 41)	Impairments
Equal to contralateral	2 (33 of 6)	
Y-balance test within 80% uninjured	1 (17)	
Star excursion balance test symmetry	1 (17)	
Demonstrated balance/coordination	1 (17)	
Prek Kinesthetic Action Training System 100%	1 (17)	
Aerobic endurance	2 (5 of 41)	Activities
Lactate aerobic–anaerobic threshold test	1 (50 of 2)	
Treadmill run for 10 min at 8 km/hour	1 (50)	
Quality of motion	3 (7 of 41)	Impairments
Landing Error Scoring System <5	2 (67 of 3)	
Limb alignment control following fatiguing task: single-leg land (QASLS score 0–1)	1 (33)	

Continued

Table 7 Continued

	Studies reporting, n (%)	ICF domain
Completion of a specific programme	27 (66 of 41)	
Running programme	6 (22 of 27)	Activities
On-field/sport-specific/functional programme	9 (33)	Participation
Rehabilitation programme	7 (26)	Activities
Agility programme	4 (15)	Activities
Proprioceptive programme	1 (4)	Activities
Non-specific performance-based criteria	6 (15 of 41)	Activities
Rehabilitation goals met	2 (33 of 6)	
Regained agility, strength and coordination	1 (17)	
Full functional stability in terms of strength, coordination and balance	1 (17)	
Able to perform sport-related movements safely per physical therapist and surgeon	1 (17)	
No functional complaints, confidence when running cutting and jumping at full speed	1 (17)	

References in online appendix III.

ACL-RSI, ACL – Return to Sport After Injury; IKDC, International Knee Documentation Committee; KOS-ADL, Knee Outcome Survey-Activities of Daily Living; QASLS, single-leg loading qualitative assessment tool.

a specific programme as part of RTS criteria did not provide specific details regarding what the programme entailed.

Time trends in RTS criteria

Time was the sole criterion used to clear athletes to RTS in studies published between 1986 and 1989 (figure 2). Since 1990, time has represented approximately 50% of the total RTS criteria reported in published research (figure 2). Measures of impairments (ie, strength and clinical examination) represented approximately 30% of the total RTS criteria reported in published research (figure 2). Measures of performance-based criteria represented approximately 10% of the total RTS criteria reported in published research (figure 2).

DISCUSSION

This review provides a compendium of tests that might be a helpful starting point for the practitioner and athlete planning RTS. The 209 included studies reported criteria that covered each of the ICF domains, although time and impairment measures were the most frequently reported criteria (figure 3).

Figure 2 Relative proportion of return to sport criteria reported in a given year.

Figure 3 Return to sport criteria mapped to the ICF domains. The relative size of the footballs represents how frequently different classifications of RTS criteria are reported in published studies.

Measures of participation were rarely reported (figure 3). For the researcher, the results of our scoping review highlight the limited assessment of the participation and contextual elements of the ICF (figure 3). These parameters are relevant for at least two reasons: (1) participation is what athletes typically care most about, and (2) the call for RTS testing to focus on replicating the key physical, physiological and psychological demands of the athlete's sport.⁹

Finding a balance between respecting biology and increasing load

Time was the most frequently reported criterion for RTS—85% of the studies included in our scoping review reported time as a RTS criterion. In two out of every five studies in our scoping review, athletes were cleared to RTS solely on the basis of time since ACL reconstruction. In over 90% of studies that used time as a RTS criterion, athletes were cleared to RTS within nine post-operative months.

Current guidelines for rehabilitation and RTS following ACL reconstruction include time among other criteria.³ The days of rehabilitation and RTS progressions solely governed by time (eg, at 3 months the athlete starts running, at 5 months the athlete starts training and at 6 months the athlete starts unrestricted sport) should be long behind us.⁶² However, tempering aggressive rehabilitation and RTS progression^{40 63} to account for biology—healing of the ACL graft and recovery of neuromuscular function—is important because of the link between time since surgery and knee reinjury risk.⁶⁴ Criterion-based rehabilitation and RTS progressions should be evaluated with appropriate tests of impairments, activities, participation and contextual factors and combined with a minimum 9-month time criterion.^{3 64}

Greater focus on the athlete and the sport

The dominance of measures of impairment (in comparison with measures of activity, participation and contextual factors) might reflect that impairment-based measures (eg, static knee laxity and strength) often require limited (or relatively easily accessible) equipment and space. They may also be easier to standardise than other types of tests. In contrast, assessing movement quality or other performance-based tests may require more complex equipment (eg, high definition video set-up), large amounts of space (eg, a football pitch) and can be more difficult to standardise.⁶⁵ However, there is a problem: if we only test impairments, we lack information about the athlete's capacity to cope with all of the physical and mental demands of playing sport. Measures of impairments are also poorly related to participation.⁶⁶ The validity and responsiveness of higher level assessments, such as performance-based measures, also has limitations and requires further clarification.⁶⁵

Over 75% of studies examining strength and/or hop testing used an LSI cut-off to determine pass or fail. Recently, there has been a call for caution when interpreting LSI,^{5 6 67 68} especially in the absence of an accurate baseline measure. Presurgical strength measurements of the uninvolved limb have been advocated⁵ as a basis for a more accurate LSI due to deterioration in the strength of the uninvolved limb following surgery.⁶⁹ Biomechanical injury risk factors may also be present, despite limb symmetry.^{6 70} Therefore, the relevance and applicability of LSI to RTS testing requires further clarification. The practitioner working in the team environment may have the luxury of access to preinjury or presurgery screening data, which can help overcome the need to rely on LSI. For the practitioner without access to a trustworthy baseline evaluation of the uninjured limb, LSI remains an acceptable measure, provided you are cognisant of its limitations.⁶⁸

Helping practitioners contribute effectively to shared RTS decisions

Contributing effectively to shared RTS decisions relies on the practitioner's knowledge of the health of the athlete's injured tissue or body part, the stress that the injured tissue or body part needs to be able to withstand and the context of sports participation.⁷⁻⁹ Tests of body structure (ie, impairments, including effusion and quadriceps strength) can provide information about tissue health. Tests of activities that resemble the basic demands of sports performance (eg, hopping and agility) can provide information about the capacity of the injured tissue or body part to withstand stress. Tests of participation (eg, pitch-based programme) provide more relevant, sport-specific information regarding tissue stress.

Understanding the demands of the sport is also a key source of information regarding how much stress is exerted on the body during sport.^{7 8} Understanding the demands of the athlete's sport is vital, and an area that coaches—who are the experts in the technical and tactical demands of the sport—might help with.

Understanding the context of sports participation (including the importance of the match and coach pressure) provides information to guide the practitioner and athlete when deciding on their shared risk tolerance. There is always some risk associated with returning to and participating in sport. Effective RTS planning involves managing risk—balancing the sometimes competing tensions of performance and reinjury prevention.⁷¹ Tests of psychological readiness to return to sport, reinjury anxiety or performance anxiety might provide insight into the context of sports participation from the athlete's perspective.

Only one study included in our scoping review used a validated measure to assess psychological readiness.⁴³ Given that psychological factors can influence RTS,⁷² these might be important factors to assess with respect to RTS.

Is research guiding practice in RTS?

How do the RTS tests reported in research stack up against the sort of information that practitioners and athletes need to help them make high-quality RTS decisions?⁷⁻⁹ In our scoping review, impairment-based tests of body structure were the most frequently reported group of tests (half of all studies). Twenty-six per cent of studies used activity-based tests (hop tests were the most common). Ten per cent of studies assessed personal or contextual factors including confidence and self-reported knee function. Fewer than 5% of studies used participation-based tests. These results suggest that often researchers may not be providing athletes and practitioners with sufficient information to make an informed RTS decision in practice.⁷³

How researchers can help practitioners

Validity (how sure we are that the test is measuring what it is supposed to be measuring) is an important aspect to consider when measuring outcomes, both in research and in practice. Take movement quality as an example—validity governs the extent to which we can be sure that our observation of knee position in the frontal plane (ie, the knee varus/valgus position) is an accurate measure of neuromuscular function.

The RTS predictive validity (ie, does the result on the test predict the outcome) of many RTS tests is unknown. There is preliminary evidence of RTS predictive validity for psychological readiness to return to sport, low fear of reinjury, symmetrical hop performance and 'normal' knee function.^{74 75} This means that for most RTS tests, we do not know whether passing the test means the athlete will RTS. Therefore, establishing the RTS predictive validity of RTS tests should be a priority for future research.

Research is especially needed to underpin the development of accurate, sport-specific, practically relevant and cost-effective tests of participation for athletes after ACL injury. Capitalising on new technologies such as augmented or virtual reality might also be a worthwhile avenue for future research as a way of realistically replicating some of the context of sports participation without excessive space or personnel requirements.

How practitioners can help athletes

The multifactorial nature of RTS suggests that a group of tests measuring athlete-centred constructs, rather than a single test in isolation, can provide an important basis for practitioners and athletes when planning RTS. Clearly, one test will not provide enough information and, for this reason, should not be the sole deciding factor in an RTS risk assessment.⁷ The 2016 consensus on RTS⁹ outlines five recommendations to guide the choice of RTS tests:

1. Use a group of tests (test battery).
2. Choose open tasks (less controlled) over closed tasks (more controlled) where possible.
3. Include tests with reactive decision-making elements.
4. Assess psychological readiness to return to sport.
5. Monitor internal and external workload.

Our scoping review indicates that published research on RTS often does not mirror the 2016 consensus statement. One explanation is that 95% of the studies in our scoping review were published prior to the most recent consensus on RTS, which outlines these

RETURN TO
PARTICIPATIONRETURN TO
SPORTRETURN TO
PERFORMANCE

Figure 4 The three elements of the RTS continuum.

recommendations. Therefore, in future research, we may begin to see a shift in the RTS criteria reported. Appropriate RTS tests of impairments, activities, participation and contextual factors, combined with a minimum 9-month time criterion,^{3 64} are worthy of further investigation. In at least one recent study, a movement quality test (Landing Error Scoring System) was used as part of a RTS test battery along with the International Knee Documentation Committee subjective knee form and the ACL-Return to Sport after Injury scale.⁴²

All of the studies included in our scoping review were focused on the RTS element of the RTS continuum (figure 4).⁹ However, for many athletes, return to performance—the third element of the RTS continuum—is their primary focus. The RTS continuum illustrates that there are at least two key transitions for athletes following injury: return to playing their sport and return to performing at the desired level in their sport. Research on athlete performance following ACL reconstruction is limited and has focused on match statistics.⁷⁶ However, match statistics may not provide adequate information on load, physiological capacity or psychological aspects of performance.

Limitations

There are limitations to our scoping review, with implications for athletes, practitioners and researchers:

- We reported tests that have been used in published research. Other innovative and potentially useful tests may be used in practice but not reflected here.⁷⁷
- It is unclear whether passing RTS tests predicts whether an athlete will return to sport participation or the previous level of sport participation after ACL reconstruction.

What is already known

- In the decade leading up to 2011, few objective functional criteria were used to clear athletes to return to sport (RTS) after ACL reconstruction.
- Sport has physical and psychological demands that must be considered when planning RTS.

What are the new findings

- Time continues to be the primary criterion used to clear athletes to RTS after ACL reconstruction.
- Strength (41% of studies), physical performance-based criteria (20% of studies) and patient-report criteria (12% of studies) were infrequently used as RTS after ACL reconstruction.
- Despite the initiation of the International Classification of Functioning, Disability and Health model in 2002 with the goals of assessment at the individual, social and policy level, RTS testing after ACL reconstruction was primarily limited to assessment of time and impairments.

- It was beyond the scope of this review to assess the predictive validity of RTS tests for safe RTS following ACL reconstruction. For many of these tests, we cannot draw conclusions regarding whether the RTS tests were associated with a reduced risk of new knee injury on resuming participation in sport.

To be included in our review, studies had to be published in English and peer reviewed, representing a possible publication bias, as well as the possibility that these limitations may have excluded other relevant tests and data. We excluded studies where participants received an allograft for their ACL reconstruction to avoid graft type as a confounding variable and studies of participants with revision surgery. Therefore, the results of our scoping review may not be generalisable to athletes with allograft or revision ACL reconstruction. We accepted the definition of ‘athlete’ as it was used in an individual study. Therefore, it is possible that different studies defined athlete in different ways, and different RTS tests may be used in different athlete populations. We did not extract data in duplicate. However, a second person independently verified the extracted data. In most studies, it was unclear how the RTS tests were conducted (ie, which practitioner(s) administered the tests and made judgements about clearance to RTS). It is possible that practitioners with different education and training backgrounds might prioritise different RTS criteria and contribute different information to the shared RTS decision-making process.

CONCLUSION

Eighty-five per cent of studies used time as a RTS criterion after ACL reconstruction. Measures of participation and contextual factors were uncommonly reported as RTS criteria after ACL reconstruction. Strength (41% of studies), physical performance-based criteria (20% of studies) and patient-report criteria (12% of studies) were infrequently used as RTS criteria.

Acknowledgements The authors would like to thank Leila Ledbetter, MLIS, for assisting with the literature search for this study.

Contributors CRB contributed to initial search, data extraction, initial drafting, review and editing of the manuscript. SP contributed to initial search, data extraction, initial drafting, review and editing of the manuscript. CLA contributed to initial drafting, review and editing of manuscript. JRM contributed to initial search, data extraction and review of manuscript. CDG and JS contributed to data extraction and review of manuscript. MPR contributed the idea of the manuscript, initial search, review of data, initial drafting, review and editing of the manuscript. All authors approved of final version of manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data available upon request.

REFERENCES

- 1 Greenberg EM, Greenberg ET, Albaugh J, *et al*. Rehabilitation practice patterns following anterior cruciate ligament reconstruction: a survey of physical therapists. *J Orthop Sports Phys Ther* 2018;48:801–11.

- 2 Dingenen B, Gokeler A. Optimization of the return-to-sport paradigm after anterior cruciate ligament reconstruction: a critical step back to move forward. *Sports Med* 2017;47:1487–500.
- 3 van Melick N, van Cingel RE, Brooijmans F, *et al.* Evidence-based clinical practice update: practice guidelines for anterior cruciate ligament rehabilitation based on a systematic review and multidisciplinary consensus. *Br J Sports Med* 2016;50:1506–15.
- 4 Grindem H, Arundale AJ, Ardern CL. Alarming underutilisation of rehabilitation in athletes with anterior cruciate ligament reconstruction: four ways to change the game. *Br J Sports Med* 2018;52:1162–3.
- 5 Wellsandt E, Failla MJ, Snyder-Mackler L. Limb symmetry indexes can overestimate knee function after anterior cruciate ligament injury. *J Orthop Sports Phys Ther* 2017;47:334–8.
- 6 Wren TAL, Mueske NM, Brophy CH, *et al.* Hop Distance Symmetry Does Not Indicate Normal Landing Biomechanics in Adolescent Athletes With Recent Anterior Cruciate Ligament Reconstruction. *J Orthop Sports Phys Ther* 2018;1–23.
- 7 Shrier I. Strategic Assessment of Risk and Risk Tolerance (StARRT) framework for return-to-play decision-making. *Br J Sports Med* 2015;49:1311–5.
- 8 Creighton DW, Shrier I, Shultz R, *et al.* Return-to-play in sport: a decision-based model. *Clin J Sport Med* 2010;20:379–85.
- 9 Ardern CL, Glasgow P, Schneiders A, *et al.* Consensus statement on return to sport from the First World Congress in Sports Physical Therapy, Bern. *Br J Sports Med* 2016;2016:853–64.
- 10 Barber-Westin SD, Noyes FR. Factors used to determine return to unrestricted sports activities after anterior cruciate ligament reconstruction. *Arthroscopy* 2011;27:1697–705.
- 11 Barber-Westin SD, Noyes FR. Objective criteria for return to athletics after anterior cruciate ligament reconstruction and subsequent reinjury rates: a systematic review. *Phys Sportsmed* 2011;39:100–10.
- 12 Murray A, Daines L, Archibald D, *et al.* The relationship and effects of golf on physical and mental health: a scoping review protocol. *Br J Sports Med* 2016;50:647–50.
- 13 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8:19–32.
- 14 Wilczynski NL, Haynes RB, Hedges Team. Developing optimal search strategies for detecting clinically sound prognostic studies in MEDLINE: an analytic survey. *BMC Med* 2004;2:23.
- 15 Wilczynski NL, Haynes RB. Optimal search strategies for detecting clinically sound prognostic studies in EMBASE: an analytic survey. *J Am Med Inform Assoc* 2005;12:481–5.
- 16 Greenhalgh T, Peacock R. Effectiveness and efficiency of search methods in systematic reviews of complex evidence: audit of primary sources. *BMJ* 2005;331:1064–5.
- 17 World Health Organization. Towards a common language for functioning, disability and health, ICF. 2002 <http://www.who.int/classifications/icf/training/icfbeginnersguide.pdf> (Accessed 28 May 2014).
- 18 Baltaci G, Yilmaz G, Atay AO. The outcomes of anterior cruciate ligament reconstructed and rehabilitated knees versus healthy knees: a functional comparison. *Acta Orthop Traumatol Turc* 2012;46:186–95.
- 19 Goradia VK, Grana WA. A comparison of outcomes at 2 to 6 years after acute and chronic anterior cruciate ligament reconstructions using hamstring tendon grafts. *Arthroscopy* 2001;17:383–92.
- 20 Harilainen A, Linko E, Sandelin J. Randomized prospective study of ACL reconstruction with interference screw fixation in patellar tendon autografts versus femoral metal plate suspension and tibial post fixation in hamstring tendon autografts: 5-year clinical and radiological follow-up results. *Knee Surg Sports Traumatol Arthrosc* 2006;14:517–28.
- 21 Jansson KA, Linko E, Sandelin J, *et al.* A prospective randomized study of patellar versus hamstring tendon autografts for anterior cruciate ligament reconstruction. *Am J Sports Med* 2003;31:12–18.
- 22 Matsumoto A, Yoshiya S, Muratsu H, *et al.* A comparison of bone-patellar tendon-bone and bone-hamstring tendon-bone autografts for anterior cruciate ligament reconstruction. *Am J Sports Med* 2006;34:213–9.
- 23 Wierer G, Runer A, Hoser C, *et al.* Acute ACL reconstruction in patients over 40 years of age. *Knee Surg Sports Traumatol Arthrosc* 2017;25:1528–34.
- 24 Zhu W, Wang D, Han Y, *et al.* Anterior cruciate ligament (ACL) autograft reconstruction with hamstring tendons: clinical research among three rehabilitation procedures. *Eur J Orthop Surg Traumatol* 2013;23:939–43.
- 25 Aglietti P, Giron F, Buzzi R, *et al.* Anterior cruciate ligament reconstruction: bone-patellar tendon-bone compared with double semitendinosus and gracilis tendon grafts. A prospective, randomized clinical trial. *J Bone Joint Surg Am* 2004;86-A:2143–55.
- 26 Buda R, Verni E, Ferruzzi A, *et al.* Anterior cruciate ligament replacement with distally inserted doubled hamstring graft: Prospective clinical and instrumental evaluation. *Med Sport* 2005;58:303–11.
- 27 Larsen JB, Farup J, Lind M, *et al.* Muscle strength and functional performance is markedly impaired at the recommended time point for sport return after anterior cruciate ligament reconstruction in recreational athletes. *Hum Mov Sci* 2015;39:73–87.
- 28 Ventura A, Iori S, Legnani C, *et al.* Single-bundle versus double-bundle anterior cruciate ligament reconstruction: assessment with vertical jump test. *Arthroscopy* 2013;29:1201–10.
- 29 Glasgow SG, Gabriel JP, Sapega AA, *et al.* The effect of early versus late return to vigorous activities on the outcome of anterior cruciate ligament reconstruction. *Am J Sports Med* 1993;21:243–8.
- 30 Noyes FR, Berrios-Torres S, Barber-Westin SD, *et al.* Prevention of permanent arthrofibrosis after anterior cruciate ligament reconstruction alone or combined with associated procedures: a prospective study in 443 knees. *Knee Surg Sports Traumatol Arthrosc* 2000;8:196–206.
- 31 Shelbourne KD, Nitz P. Accelerated rehabilitation after anterior cruciate ligament reconstruction. *Am J Sports Med* 1990;18:292–9.
- 32 Shelbourne KD, Nitz P. Accelerated rehabilitation after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 1992;15:256–64.
- 33 Sonnerly-Cottet B, Saithna A, Cavalier M, *et al.* Anterolateral ligament reconstruction is associated with significantly reduced acl graft rupture rates at a minimum follow-up of 2 years: a prospective comparative study of 502 patients from the SANTI study group. *Am J Sports Med* 2017;45:1547–57.
- 34 Sonnerly-Cottet B, Thauinat M, Freychet B, *et al.* Outcome of a Combined Anterior Cruciate Ligament and Anterolateral Ligament Reconstruction Technique With a Minimum 2-Year Follow-up. *Am J Sports Med* 2015;43:1598–605.
- 35 Tashiro T, Kurosawa H, Kawakami A, *et al.* Influence of medial hamstring tendon harvest on knee flexor strength after anterior cruciate ligament reconstruction. A detailed evaluation with comparison of single- and double-tendon harvest. *Am J Sports Med* 2003;31:522–9.
- 36 Hohmann E, Tetsworth K, Bryant A. Physiotherapy-guided versus home-based, unsupervised rehabilitation in isolated anterior cruciate injuries following surgical reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1158–67.
- 37 Capin JJ, Khandha A, Zarzycki R, *et al.* Gait mechanics and second ACL rupture: Implications for delaying return-to-sport. *J Orthop Res* 2017;35:1894–901.
- 38 Failla MJ, Logerstedt DS, Grindem H, *et al.* Does Extended Preoperative Rehabilitation Influence Outcomes 2 Years After ACL Reconstruction? A Comparative Effectiveness Study Between the MOON and Delaware-Oslo ACL Cohorts. *Am J Sports Med* 2016;44:2608–14.
- 39 Gokeler A, Welling W, Zaffagnini S, *et al.* Development of a test battery to enhance safe return to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2017;25:192–9.
- 40 Shelbourne KD, Sullivan AN, Bohard K, *et al.* Return to basketball and soccer after anterior cruciate ligament reconstruction in competitive school-aged athletes. *Sports Health* 2009;1:236–41.
- 41 Toole AR, Ithurburn MP, Rau MJ, *et al.* Young athletes cleared for sports participation after anterior cruciate ligament reconstruction: how many actually meet recommended return-to-sport criterion cutoffs? *J Orthop Sports Phys Ther* 2017;47:825–33.
- 42 Welling W, Benjaminse A, Seil R, *et al.* Low rates of patients meeting return to sport criteria 9 months after anterior cruciate ligament reconstruction: a prospective longitudinal study. *Knee Surg Sports Traumatol Arthrosc* 2018;26:3636–44.
- 43 Langford JL, Webster KE, Feller JA. A prospective longitudinal study to assess psychological changes following anterior cruciate ligament reconstruction surgery. *Br J Sports Med* 2009;43:377–8.
- 44 Goodstadt NM, Hunter-Giordano A, Axe MJ, *et al.* Functional testing to determine readiness to discontinue brace use, one year after acl reconstruction. *Int J Sports Phys Ther* 2013;8:91–6.
- 45 Ferrari JD, Bach BR, Bush-Joseph CA, *et al.* Anterior cruciate ligament reconstruction in men and women: An outcome analysis comparing gender. *Arthroscopy* 2001;17:588–96.
- 46 Zaffagnini S, Bruni D, Marcheggiani Muccioli GM, *et al.* Single-bundle patellar tendon versus non-anatomical double-bundle hamstrings ACL reconstruction: a prospective randomized study at 8-year minimum follow-up. *Knee Surg Sports Traumatol Arthrosc* 2011;19:390–7.
- 47 Zaffagnini S, Marcacci M, Lo Presti M, *et al.* Prospective and randomized evaluation of ACL reconstruction with three techniques: a clinical and radiographic evaluation at 5 years follow-up. *Knee Surg Sports Traumatol Arthrosc* 2006;14:1060–9.
- 48 Fibiger W. Developing an objective criteria for recommendations to athletes prior to a return to full training following knee acl reconstruction. *Medicina Sportiva* 2014;18:147–52.
- 49 Noyes FR, Barber-Westin SD. Arthroscopic repair of meniscus tears extending into the avascular zone with or without anterior cruciate ligament reconstruction in patients 40 years of age and older. *Arthroscopy* 2000;16:822–9.
- 50 Ardern CL, Taylor NF, Feller JA, *et al.* Psychological responses matter in returning to preinjury level of sport after anterior cruciate ligament reconstruction surgery. *Am J Sports Med* 2013;41:1549–58.
- 51 Ardern CL, Webster KE, Taylor NF, *et al.* Return to the preinjury level of competitive sport after anterior cruciate ligament reconstruction surgery: two-thirds of patients have not returned by 12 months after surgery. *Am J Sports Med* 2011;39:538–43.
- 52 Beynon BD, Johnson RJ, Fleming BC, *et al.* Anterior cruciate ligament replacement: comparison of bone-patellar tendon-bone grafts with two-strand hamstring grafts. A prospective, randomized study. *J Bone Joint Surg Am* 2002;84-A:1503–13.

- 53 MacDonald PB, Hedden D, Pacin O, *et al.* Effects of an accelerated rehabilitation program after anterior cruciate ligament reconstruction with combined semitendinosus-gracilis autograft and a ligament augmentation device. *Am J Sports Med* 1995;23:588–92.
- 54 Mendias CL, Lynch EB, Davis ME, *et al.* Changes in circulating biomarkers of muscle atrophy, inflammation, and cartilage turnover in patients undergoing anterior cruciate ligament reconstruction and rehabilitation. *Am J Sports Med* 2013;41:1819–26.
- 55 Palmer W, Crichlow A, Mansingh A. Return to sports and functional outcome after primary anterior cruciate ligament reconstruction in Jamaica. *Sports Nutr Ther* 2016;1:2.
- 56 Palmieri-Smith RM, Lepley LK. Quadriceps strength asymmetry after anterior cruciate ligament reconstruction alters knee joint biomechanics and functional performance at time of return to activity. *Am J Sports Med* 2015;43:1662–9.
- 57 Siebold R, Webster KE, Feller JA, *et al.* Anterior cruciate ligament reconstruction in females: a comparison of hamstring tendon and patellar tendon autografts. *Knee Surg Sports Traumatol Arthrosc* 2006;14:1070–6.
- 58 Della Villa S, Boldrini L, Ricci M, *et al.* Clinical outcomes and return-to-sports participation of 50 soccer players after anterior cruciate ligament reconstruction through a sport-specific rehabilitation protocol. *Sports Health* 2012;4:17–24.
- 59 Webster KE, Feller JA, Hameister KA. Bone tunnel enlargement following anterior cruciate ligament reconstruction: a randomised comparison of hamstring and patellar tendon grafts with 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2001;9:86–91.
- 60 Müller U, Krüger-Franke M, Schmidt M, *et al.* Predictive parameters for return to pre-injury level of sport 6 months following anterior cruciate ligament reconstruction surgery. *Knee Surg Sports Traumatol Arthrosc* 2015;23:3623–31.
- 61 Beynnon BD, Johnson RJ, Naud S, *et al.* Accelerated versus nonaccelerated rehabilitation after anterior cruciate ligament reconstruction: a prospective, randomized, double-blind investigation evaluating knee joint laxity using roentgen stereophotogrammetric analysis. *Am J Sports Med* 2011;39:2536–48.
- 62 Rambaud AJM, Ardern CL, Thoreux P, *et al.* Criteria for return to running after anterior cruciate ligament reconstruction: a scoping review. *Br J Sports Med* 2018;52:1437–44.
- 63 Roi GS, Creta D, Nanni G, *et al.* Return to official Italian First Division soccer games within 90 days after anterior cruciate ligament reconstruction: a case report. *J Orthop Sports Phys Ther* 2005;35:52–66.
- 64 Grindem H, Snyder-Mackler L, Moksnes H, *et al.* Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. *Br J Sports Med* 2016;50:804–8.
- 65 Hegedus EJ, McDonough S, Bleakley C, *et al.* Clinician-friendly lower extremity physical performance measures in athletes: a systematic review of measurement properties and correlation with injury, part 1. The tests for knee function including the hop tests. *Br J Sports Med* 2015;49:642–8.
- 66 Ross MD, Irrgang JJ, Denegar CR, *et al.* The relationship between participation restrictions and selected clinical measures following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2002;10:10–19.
- 67 Benjamin A, Holden S, Myer GD. ACL rupture is a single leg injury but a double leg problem: too much focus on 'symmetry' alone and that's not enough!. *Br J Sports Med* 2018;52:1029–30.
- 68 Trulsson A. Additional perspectives on 'ACL rupture is a single leg injury but a double leg problem...'. *Br J Sports Med* 2018;bjsports-2017-098974.
- 69 Thomeé R, Neeter C, Gustavsson A, *et al.* Variability in leg muscle power and hop performance after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2012;20:1143–51.
- 70 Xergia SA, Pappas E, Georgoulis AD. Association of the single-limb hop test with isokinetic, kinematic, and kinetic asymmetries in patients after anterior cruciate ligament reconstruction. *Sports Health* 2015;7:217–23.
- 71 McCall A, Lewin C, O'Driscoll G, *et al.* Return to play: the challenge of balancing research and practice. *Br J Sports Med* 2017;51:702–3.
- 72 Ardern CL, Taylor NF, Feller JA, *et al.* A systematic review of the psychological factors associated with returning to sport following injury. *Br J Sports Med* 2013;47:1120–6.
- 73 Ardern CL, Kvist J, Webster KE. Psychological aspects of anterior cruciate ligament injuries. *Oper Tech Sports Med* 2016;24:77–83.
- 74 Ardern CL, Taylor NF, Feller JA, *et al.* Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med* 2014;48:1543–52.
- 75 Nawasreh Z, Logerstedt D, Cummer K, *et al.* Functional performance 6 months after ACL reconstruction can predict return to participation in the same preinjury activity level 12 and 24 months after surgery. *Br J Sports Med* 2018;52:bjsports-2016-097095.
- 76 Lai CCH, Ardern CL, Feller JA, *et al.* Eighty-three per cent of elite athletes return to preinjury sport after anterior cruciate ligament reconstruction: a systematic review with meta-analysis of return to sport rates, graft rupture rates and performance outcomes. *Br J Sports Med* 2018;52:128–38.
- 77 Ardern CL, Dupont G, Impellizzeri FM, *et al.* Unravelling confusion in sports medicine and sports science practice: a systematic approach to using the best of research and practice-based evidence to make a quality decision. *Br J Sports Med* 2019;53.