

Appendix II. Full text exclusion reasons

No return to sport criteria reported n = 189
Ageberg, E., et al. Knee extension and flexion muscle power after anterior cruciate ligament reconstruction with patellar tendon graft or hamstring tendons graft: a cross-sectional comparison 3 years post surgery. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2009;17(2):162-169.
Ageberg, E., et al. Muscle strength and functional performance in patients with anterior cruciate ligament injury treated with training and surgical reconstruction or training only: a two to five-year followup. <i>Arthritis and Rheumatism</i> 2008;59(12): 1773-1779.
Ageberg, E. and E. M. Roos The association between knee confidence and muscle power, hop performance, and postural orientation in people with anterior cruciate ligament injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2016;46(6): 477-482.
Ahlden, M., et al. The Swedish National Anterior Cruciate Ligament Register: a report on baseline variables and outcomes of surgery for almost 18,000 patients. <i>American Journal of Sports Medicine</i> 2012;40(10): 2230-2235.
Akelman, M. R., et al. Effect of matching or overconstraining knee laxity during anterior cruciate ligament reconstruction on knee osteoarthritis and clinical outcomes: a randomized controlled trial with 84-month follow-up. <i>American Journal of Sports Medicine</i> 2016;44(7): 1660-1670.
Alentorn-Geli, E., et al. Anteromedial portal versus transtibial drilling techniques in ACL reconstruction: a blinded cross-sectional study at two- to five-year follow-up." <i>International Orthopaedics</i> 2010;34(5): 747-754.
Ali, S. D., et al. Functional outcome of ACL reconstruction using patellar bone tendon bone graft. <i>Journal of the Pakistan Medical Association</i> 2014;64(12): S79-82.
Arciero, R. A., et al. Single versus two-incision arthroscopic anterior cruciate ligament reconstruction. <i>Arthroscopy</i> 1996;12(4): 462-469.
Ardern, C. L., et al. The impact of psychological readiness to return to sport and recreational activities after anterior cruciate ligament reconstruction. <i>British Journal of Sports Medicine</i> 2014;48(22): 1613-1619.
Barber-Westin, S. D., et al. The effect of exercise and rehabilitation on anterior-posterior knee displacements after anterior cruciate ligament autograft reconstruction. <i>American Journal of Sports Medicine</i> 1999;27(1): 84-93.
Barenius, B., et al. Functional recovery after anterior cruciate ligament reconstruction, a study of health-related quality of life based on the Swedish National Knee Ligament Register. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2013;21(4): 914-927.
Bell, D. R., et al. Jump-landing mechanics after anterior cruciate ligament reconstruction: a landing error scoring system study. <i>J Athl Train</i> 2014;49(4): 435-441.
Billotti, J. D., et al. A prospective, clinical study evaluating arthroscopic ACL reconstruction using the semitendinosus and iliotibial band: 2- to 5-year follow up. <i>Orthopedics</i> 1997;20(2): 125-131.
Birmingham, T. B., et al. A randomized controlled trial comparing the effectiveness of functional knee brace and neoprene sleeve use after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 2008;36(4): 648-655.
Blonna, D., et al. Validity and reliability of the SPORTS score. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2012;20(2): 356-360.
Boden, B. P., et al. Arthroscopically-assisted anterior cruciate ligament reconstruction: a follow-up study. <i>Contemporary Orthopaedics</i> 1990;20(2): 187-194.
Bonnard, C., et al. Physseal-sparing reconstruction of anterior cruciate ligament tears in children: results of 57 cases using patellar tendon. <i>Journal of Bone and Joint Surgery (British Volume)</i> 2011;93(4): 542-547.
Bordes, P., et al. No beneficial effect of bracing after anterior cruciate ligament reconstruction in a cohort of 969 athletes followed in rehabilitation. <i>Annals of Physical and Rehabilitation Medicine</i> 2017;60(4): 230-236.
Boyle, M. J., et al. Functional movement competency and dynamic balance after anterior cruciate ligament reconstruction in adolescent patients. <i>Journal of Pediatric Orthopedics</i> 2016;36(1): 36-41.
Brewer, B. W., et al. Age-related differences in predictors of adherence to rehabilitation after anterior cruciate ligament reconstruction. <i>J Athl Train</i> 2003;38(2): 158-162.
Brewer, B. W., et al. Psychological factors, rehabilitation adherence, and rehabilitation outcome after

anterior cruciate ligament reconstruction. <i>Rehabilitation Psychology</i> 2000;45(1): 20-37.
Brewer, B. W. and A. E. Cornelius. Self-protective changes in athletic identity following anterior cruciate ligament reconstruction. <i>Psychology of Sport and Exercise</i> 2010;11(1): 1-5.
Brophy, R. H., et al. Return to play and future ACL injury risk after ACL reconstruction in soccer athletes from the Multicenter Orthopaedic Outcomes Network (MOON) group. <i>American Journal of Sports Medicine</i> 2012;40(11): 2517-2522.
Chaudhary, D., et al. Arthroscopic reconstruction of the anterior cruciate ligament using bone-patellar tendon-bone autograft: experience of the first 100 cases. <i>Journal of Orthopaedic Surgery (Hong Kong)</i> 2005;13(2): 147-152.
Chaves, S., et al. Neuromuscular efficiency of the vastus medialis obliquus in soccer athletes following an anterior cruciate ligament reconstruction. <i>Physical Therapy in Sport</i> 2012;13(3): e2-e2.
Chmielewski, T. L., et al. Low- versus high-intensity plyometric exercise during rehabilitation after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 2016;44(3): 609-617.
Clagg, S., et al. Performance on the modified star excursion balance test at the time of return to sport following anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2015;45(6): 444-452.
Cohen, M., et al. Transphyseal anterior cruciate ligament reconstruction in patients with open physes. <i>Arthroscopy</i> 2009;25(8): 831-838.
Colombet, P., et al. Anterior cruciate ligament reconstruction using four-strand semitendinosus and gracilis tendon grafts and metal interference screw fixation. <i>Arthroscopy</i> 2002;18(3): 232-237.
Cordasco, F. A., et al. All-inside, all-epiphyseal ACL reconstruction in skeletally immature athletes: Incidence of second surgery and two-year clinical outcomes. <i>Orthop J Sports Med</i> 2015;3(7).
Cox, C. L., et al. Are articular cartilage lesions and meniscus tears predictive of IKDC, KOOS, and Marx Activity Level outcomes after ACL reconstruction? A 6-year moon cohort study. <i>Osteoarthritis and Cartilage</i> 2013;21: S221.
Culvenor, A. G., et al. Accelerated return to sport after ACL reconstruction and early knee osteoarthritis features at 1 year: an exploratory study. <i>Pm R.</i> 2018;10(4):349-356.
Cvjetkovic, D. D., et al. Isokinetic Testing in Evaluation Rehabilitation Outcome After ACL Reconstruction. <i>Med Arch</i> 2015;69(1): 21-23.
Czamara, A., et al. Single- vs. double-bundle anterior cruciate ligament reconstruction: a new aspect of knee assessment during activities involving dynamic knee rotation. <i>Journal of Strength and Conditioning Research</i> 2015;29(2): 489-499.
Dauty, M., et al. Effects of running retraining after knee anterior cruciate ligament reconstruction. <i>Annals of Physical and Rehabilitation Medicine</i> 2010;53(3): 150-161.
Deehan, D. J., et al. Endoscopic reconstruction of the anterior cruciate ligament with an ipsilateral patellar tendon autograft. <i>Journal of Bone and Joint Surgery - Series B</i> 2000;82(7): 984-991.
Dei Giudici, L., et al. Arthroscopic transphyseal anterior cruciate ligament reconstruction in adolescent athletes. <i>Journal of Orthopaedic Surgery (Hong Kong)</i> 2016;24(3): 307-311.
Dekker, T. J., et al. Return to sport after pediatric anterior cruciate ligament reconstruction and its effect on subsequent anterior cruciate ligament injury. <i>Journal of Bone and Joint Surgery (American Volume)</i> 2017;99(11): 897-904.
Demange, M. K. and G. L. Camanho. Nonanatomic anterior cruciate ligament reconstruction with double-stranded semitendinosus grafts in children with open physes: minimum 15-year follow-up. <i>American Journal of Sports Medicine</i> 2014;42(12): 2926-2932.
Dingenen, B., et al. Lower extremity muscle activation onset times during the transition from double-leg stance to single-leg stance in anterior cruciate ligament reconstructed subjects. <i>Clinical Biomechanics (Bristol, Avon)</i> 2016;35: 116-123.
do Carmo Almeida, T. C., et al. Evaluation of functional rehabilitation physiotherapy protocol in the postoperative patients with anterior cruciate ligament reconstruction through clinical prognosis: an observational prospective study. <i>BMC Research Notes</i> 2016;9(1): 449.
Domzalski, M., et al. Anterior Cruciate Ligament Reconstruction Using the transphyseal technique in prepubescent athletes: midterm, prospective evaluation of results. <i>Arthroscopy</i> 2016;32(6): 1141-1146.
Draper, V. and C. Ladd. Subjective evaluation of function following moderately accelerated rehabilitation of anterior cruciate ligament reconstructed knees. <i>J Athl Train</i> 1993;28(1): 38-41.
Ebert, J. R., et al. Strength and functional symmetry is associated with post-operative rehabilitation in

patients following anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> . 2018;26(8):2353-2361.
Edwards, P. H. and W. A. Grana. Anterior cruciate ligament reconstruction in the immature athlete: long-term results of intra-articular reconstruction. <i>American Journal of Knee Surgery</i> 2001;14(4): 232-237.
Eisenstein, E. D., et al. Variables affecting return to play after anterior cruciate ligament injury in the National Football League. <i>Orthop J Sports Med</i> 2016;4(10): 2325967116670117.
Ellera Gomes, J. L. and L. R. Marczyk. Anterior cruciate ligament reconstruction with a loop or double thickness of semitendinosus tendon. <i>American Journal of Sports Medicine</i> 1984;12(3): 199-203.
Engelen-van Melick, N., et al. Functional performance 2-9 years after ACL reconstruction: cross-sectional comparison between athletes with bone-patellar tendon-bone, semitendinosus/gracilis and healthy controls. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2017;25(5): 1412-1423.
Erickson, B. J., et al. Performance and return to sport after anterior cruciate ligament reconstruction in male Major League Soccer players. <i>Orthop J Sports Med</i> 2013;1(2): 2325967113497189.
Erickson, B. J., et al. Performance and return to sport after anterior cruciate ligament reconstruction in X-Games skiers and snowboarders. <i>Orthop J Sports Med</i> 2013;1(6): 2325967113511196.
Erickson, B. J., et al. Performance and return to sport after anterior cruciate ligament reconstruction in National Hockey League pPlayers. <i>Orthop J Sports Med</i> 2014;2(9): 2325967114548831.
Erickson, B. J., et al. Performance and return-to-sport after ACL reconstruction in NFL quarterbacks. <i>Orthopedics</i> 2014;37(8): e728-734.
Fabbriciani, C., et al. Anterior cruciate ligament reconstruction with doubled semitendinosus and gracilis tendon graft in rugby players. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2005;13(1): 2-7.
Fabricant, P. D., et al. Return to play after anterior cruciate ligament reconstruction in major league baseball athletes. <i>Arthroscopy</i> 2015;31(5): 896-900.
Faltstrom, A., et al. Factors associated with playing football after anterior cruciate ligament reconstruction in female football players. <i>Scandinavian Journal of Medicine and Science in Sports</i> 2016;26(11): 1343-1352.
Faltstrom, A., et al. Functional performance among active female soccer players after unilateral primary anterior cruciate ligament reconstruction compared with knee-healthy controls. <i>American Journal of Sports Medicine</i> 2017;45(2): 377-385.
Fernandes, T. L., et al. The influence of femoral tunnel position in single-bundle ACL reconstruction on functional outcomes and return to sports. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2014;22(1): 97-103.
Fernandes, T. L., et al. Isokinetic muscle strength and knee function associated with double femoral pin fixation and fixation with interference screw in anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2012;20(2): 275-280.
Fischer, D. A., et al. Home based rehabilitation for anterior cruciate ligament reconstruction. <i>Clinical Orthopaedics and Related Research</i> 1998;347:194-199.
Flanagan, E. P., et al. Force production and reactive strength capabilities after anterior cruciate ligament reconstruction. <i>J Athl Train</i> 2008;43(3): 249-257.
Flanigan, D. C., et al. Fear of reinjury (kinesiophobia) and persistent knee symptoms are common factors for lack of return to sport after anterior cruciate ligament reconstruction. <i>Arthroscopy</i> 2013;29(8): 1322-1329.
Flosadottir, V., et al. Muscle function is associated with future patient-reported outcomes in young adults with ACL injury. <i>BMJ Open Sport Exerc Med</i> 2016;2(1): e000154.
Gadea, F., et al. Knee pain after anterior cruciate ligament reconstruction: evaluation of a rehabilitation protocol. <i>European Journal of Orthopaedic Surgery & Traumatology</i> 2014;24(5): 789-795.
Gebhard, F., et al. Multicenter-study of operative treatment of intraligamentous tears of the anterior cruciate ligament in children and adolescents: comparison of four different techniques. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2006;14(9): 797-803.
Gobbi, A., et al. Comparison of anterior cruciate ligament reconstruction in male and female athletes using the patellar tendon and hamstring autografts. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2004;12(6): 534-539.
Gobbi, A. and R. Francisco. Factors affecting return to sports after anterior cruciate ligament reconstruction with patellar tendon and hamstring graft: a prospective clinical investigation. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2006;14(10): 1021-1028.

Gobbi, A., et al. Patellar tendon anterior cruciate ligament reconstruction with conical press-fit femoral fixation: 5-year results in athletes population. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2002;10(2): 73-79.
Gobbi, A., et al. Patellar tendon versus quadrupled bone-semitendinosus anterior cruciate ligament reconstruction: a prospective clinical investigation in athletes. <i>Arthroscopy</i> 2003;19(6): 592-601.
Gokeler, A., et al. A critical analysis of limb symmetry indices of hop tests in athletes after anterior cruciate ligament reconstruction: A case control study. <i>Orthopaedics & Traumatology, Surgery & Research</i> 2017;103(6): 947-951.
Grant, J. A., et al. Comparison of home versus physical therapy-supervised rehabilitation programs after anterior cruciate ligament reconstruction: a randomized clinical trial. <i>American Journal of Sports Medicine</i> 2005;33(9): 1288-1297.
Grindem, H., et al. How does a combined preoperative and postoperative rehabilitation programme influence the outcome of ACL reconstruction 2 years after surgery? A comparison between patients in the Delaware-Oslo ACL Cohort and the Norwegian National Knee Ligament Registry. <i>British Journal of Sports Medicine</i> 2015;49(6): 385-389.
Guimarães, M. V., et al. Reconstruction of the anterior cruciate ligament with the central third of the quadriceps muscle tendon: analysis of 10-year results. <i>Revista Brasileira de Ortopedia (English Edition)</i> 2009;44(4): 306-312.
Hamrin Senorski, E., et al. Return to knee-strenuous sport after anterior cruciate ligament reconstruction: a report from a rehabilitation outcome registry of patient characteristics. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2017;25(5): 1364-1374.
Harner, C. D., et al. Anterior cruciate ligament reconstruction: endoscopic versus two-incision technique. <i>Arthroscopy</i> 1994;10(5): 502-512.
Harpur, G., et al. Relation of self-reported knee function and physical performance with psychological responses in anterior cruciate ligament reconstructed individuals. <i>Fizyoterapi Rehabilitasyon</i> 2016;27(3): 102-107.
Harpur, G., et al. External supports improve knee performance in anterior cruciate ligament reconstructed individuals with higher kinesiophobia levels. <i>Knee</i> 2016;23(5): 807-812.
Harris, J. D., et al. Return-to-sport and performance after anterior cruciate ligament reconstruction in National Basketball Association players. <i>Sports Health</i> 2013;5(6): 562-568.
Hartigan, E., et al. Biomechanical profiles when towing a sled and wearing a weighted vest once cleared for sports post-ACL reconstruction. <i>Sports Health</i> 2016;8(5): 456-464.
Hegde, A. S., et al. A Comparison of functional outcomes after metallic and bioabsorbable interference screw fixations in arthroscopic ACL reconstructions. <i>J Clin Diagn Res</i> 2014;8(4): Lc01-03.
Holwein, C., et al. Functional outcome after transphyseal anterior cruciate ligament reconstruction in young patients with open growth plates. <i>Knee</i> 2016;23(6): 1121-1132.
Hopper, D. M., et al. Functional recovery after anterior cruciate ligament reconstruction: a longitudinal perspective. <i>Archives of Physical Medicine and Rehabilitation</i> 2008;89(8): 1535-1541.
Imbert, P., et al. Midterm results of combined intra- and extra-articular ACL reconstruction compared to historical ACL reconstruction data Multicenter study of the Francophone Society of Arthroscopy. <i>Orthopaedics & Traumatology, Surgery & Research</i> 2017;103(8S):S215-S221.
Ithurnburn, M. P., et al. Young athletes after anterior cruciate ligament reconstruction with single-leg landing asymmetries at the time of return to sport demonstrate decreased knee function 2 years later. <i>American Journal of Sports Medicine</i> 2017;45(11): 2604-2613.
Ithurnburn, M. P., et al. Young athletes with quadriceps femoris strength asymmetry at return to sport after anterior cruciate ligament reconstruction demonstrate asymmetric single-leg drop-landing mechanics. <i>American Journal of Sports Medicine</i> 2015;43(11): 2727-2737.
Ithurnburn, M. P., et al. Young athletes after ACL reconstruction with quadriceps strength asymmetry at the time of return-to-sport demonstrate decreased knee function 1 year later. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> . 2018;26(2)426-433.
Jacopetti, M., et al. Evaluation of strength muscle recovery with isokinetic, squat jump and stiffness tests in athletes with ACL reconstruction: a case control study. <i>Acta Bio-Medica: Atenei Parmensis</i> 2016;87(1): 76-80.
Kadija, M., et al. The effect of anterior cruciate ligament reconstruction on hamstring and quadriceps muscle function outcome ratios in male athletes. <i>Srpski Arhiv za Celokupno Lekarstvo</i> 2016;144(3-4): 151-

157.
Khalid Ur Rehman, M., et al. Return to sports following anterior cruciate ligament reconstruction - Short term follow up. <i>Medical Forum Monthly</i> 2015;26(8): 54-57.
Kilinc, B. E., et al. Transtibial vs anatomical single bundle technique for anterior cruciate ligament reconstruction: A Retrospective Cohort Study. <i>International Journal of Surgery (London, England)</i> 2016;29: 62-69.
Kline, P. W., et al. Clinical predictors of knee mechanics at return to sport after ACL reconstruction. <i>Medicine and Science in Sports and Exercise</i> 2016;48(5): 790-795.
Kline, P. W., et al. Hip external rotation strength predicts hop performance after anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> .2018;26(4):1137-1144.
Knezevic, O. M., et al. Alternating consecutive maximum contraction as a test of muscle function in athletes following ACL reconstruction. <i>J Hum Kinet</i> 2012;35: 5-13.
Knezevic, O. M., et al. Evaluation of isokinetic and isometric strength measures for monitoring muscle function recovery after anterior cruciate ligament reconstruction. <i>Journal of Strength and Conditioning Research</i> 2014;28(6): 1722-1731.
Kong, D. H., et al. Validation of functional performance tests after anterior cruciate ligament reconstruction. <i>Knee Surg Relat Res</i> 2012;24(1): 40-45.
Koutras, G., et al. Comparison of knee flexion isokinetic deficits between seated and prone positions after ACL reconstruction with hamstrings graft: Implications for rehabilitation and return to sports decisions. <i>Journal of Science and Medicine in Sport</i> 2016;19(7): 559-562.
Kuenze, C. M., et al. Persistent neuromuscular and corticomotor quadriceps asymmetry after anterior cruciate ligament reconstruction. <i>J Athl Train</i> 2015;50(3): 303-312.
Kuenze, C., et al. Clinical thresholds for quadriceps assessment after anterior cruciate ligament reconstruction. <i>J Sport Rehabil</i> 2015;24(1): 36-46.
Labanca, L., et al. Asymmetrical lower extremity loading early after anterior cruciate ligament reconstruction is a significant predictor of asymmetrical loading at the time of return to sport. <i>American Journal of Physical Medicine and Rehabilitation</i> 2016;95(4): 248-255.
Laboute, E., et al. Analysis of return to competition and repeat rupture for 298 anterior cruciate ligament reconstructions with patellar or hamstring tendon autograft in sportspeople. <i>Annals of Physical and Rehabilitation Medicine</i> 2010;53(10): 598-614.
Lebel, B., et al. Arthroscopic reconstruction of the anterior cruciate ligament using bone-patellar tendon-bone autograft: a minimum 10-year follow-up. <i>American Journal of Sports Medicine</i> 2008;36(7): 1275-1282.
Lee, M., et al. Enhanced knee joint function due to accelerated rehabilitation exercise after anterior cruciate ligament reconstruction surgery in Korean male high school soccer players. <i>J Exerc Rehabil</i> 2016;2(1): 29-36.
Lepley, L. K., et al. Combination of eccentric exercise and neuromuscular electrical stimulation to improve quadriceps function post-ACL reconstruction. <i>Knee</i> 2015;22(3): 270-277.
Letchford, R., et al. A novel clinical approach for assessing hop landing strategies: a 2D telescopic inverted pendulum (TIP) model. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2016;24(1): 279-286.
Mae, T., et al. Anatomic double-bundle anterior cruciate ligament reconstruction using hamstring tendons with minimally required initial tension. <i>Arthroscopy</i> 2010;26(10): 1289-1295.
Maletis, G. B., et al. A prospective randomized study of anterior cruciate ligament reconstruction: A comparison of patellar tendon and quadruple-strand semitendinosus/gracilis tendons fixed with bioabsorbable interference screws. <i>American Journal of Sports Medicine</i> 2007;35(3): 384-394.
Mardani-Kivi, M., et al. Antero-medial portal vs. transtibial techniques for drilling femoral tunnel in ACL reconstruction using 4-strand hamstring tendon: a cross-sectional study with 1-year follow-up. <i>Medical Science Monitor</i> 2012;18(11): Cr674-679.
Mattacola, C. G., et al. Strength, functional outcome, and postural stability after anterior cruciate ligament reconstruction. <i>J Athl Train</i> 2002;37(3): 262-268.
Mayer, S. W., et al. Functional Testing Differences in Anterior Cruciate Ligament Reconstruction Patients Released Versus Not Released to Return to Sport. <i>American Journal of Sports Medicine</i> 2015;43(7): 1648-1655.
Memeo, A., et al. Anterior cruciate ligament reconstruction with bone-patellar tendon-bone autograft in Tanner 3 stage patients with open physes. <i>Journal of Pediatric Orthopaedics. Part B</i> 2012;21(5): 415-420.

Mikkelsen, C., et al. Closed kinetic chain alone compared to combined open and closed kinetic chain exercises for quadriceps strengthening after anterior cruciate ligament reconstruction with respect to return to sports: a prospective matched follow-up study. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2000;8(6): 337-342.
Muaidi, Q. I., et al. Effect of anterior cruciate ligament injury and reconstruction on proprioceptive acuity of knee rotation in the transverse plane. <i>American Journal of Sports Medicine</i> 2009;37(8): 1618-1626.
Mulcahey, M. K., et al. Transtibial versus anteromedial portal anterior cruciate ligament reconstruction using soft-tissue graft and expandable fixation. <i>Arthroscopy</i> 2014;30(11): 1461-1467.
Muneta, T., et al. Effects of different initial bundle tensioning strategies on the outcome of double-bundle ACL reconstruction: A cohort study. <i>Sports Medicine, Arthroscopy, Rehabilitation, Therapy and Technology</i> 2011;3(1).
Myer, G. D., et al. No association of time from surgery with functional deficits in athletes after anterior cruciate ligament reconstruction: evidence for objective return-to-sport criteria. <i>American Journal of Sports Medicine</i> 2012;40(10): 2256-2263.
Myer, G. D., et al. Utilization of modified NFL combine testing to identify functional deficits in athletes following ACL reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2011;41(6): 377-387.
Nakayama, Y., et al. Knee functions and a return to sports activity in competitive athletes following anterior cruciate ligament reconstruction. <i>J Nippon Med Sch</i> 2000;67(3): 172-176.
Neeter, C., et al. Development of a strength test battery for evaluating leg muscle power after anterior cruciate ligament injury and reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2006;14(6): 571-580.
Notarnicola, A., et al. Returning to sport after anterior cruciate ligament reconstruction in amateur sports men: a retrospective study. <i>Muscles Ligaments Tendons J</i> 2016;6(4): 486-491.
Oberoi, I. P. S., et al. Anatomical double bundle ACL reconstruction using hamstring tendon graft-clinical evaluation. <i>Journal of Clinical Orthopaedics and Trauma</i> 2010;1(1): 26-32.
Olivier, N., et al. The effect of a one-leg cycling aerobic training program during the rehabilitation period in soccer players with anterior cruciate ligament reconstruction. <i>Clinical Journal of Sport Medicine</i> 2010;20(1): 28-33.
Osteras, H., et al. Isokinetic muscle strength after anterior cruciate ligament reconstruction. <i>Scandinavian Journal of Medicine and Science in Sports</i> 1998;8(5): 279-282.
Pantano, K. J., et al. A pilot study on the relationship between physical impairment and activity restriction in persons with anterior cruciate ligament reconstruction at long-term follow-up. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2001;9(6): 369-378.
Papalia, R., et al. Anterior cruciate ligament reconstruction and return to sport activity: postural control as the key to success. <i>International Orthopaedics</i> 2015;39(3): 527-534.
Paterno, M. V., et al. Clinical Factors That Predict a Second ACL Injury After ACL Reconstruction and Return to Sport: Preliminary Development of a Clinical Decision Algorithm. <i>Orthop J Sports Med</i> 2017;5(12): 2325967117745279.
Paterno, M. V., et al. Altered postural sway persists after anterior cruciate ligament reconstruction and return to sport. <i>Gait and Posture</i> 2013;38(1): 136-140.
Paterno, M. V., et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. <i>American Journal of Sports Medicine</i> 2010;38(10): 1968-1978.
Paterno, M. V., et al. Effects of sex on compensatory landing strategies upon return to sport after anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2011;41(8): 553-559.
Paterno, M. V., et al. Incidence of contralateral and ipsilateral anterior cruciate ligament (ACL) injury after primary ACL reconstruction and return to sport. <i>Clinical Journal of Sport Medicine</i> 2012;22(2): 116-121.
Paterno, M. V., et al. Incidence of Second ACL Injuries 2 Years After Primary ACL Reconstruction and Return to Sport. <i>American Journal of Sports Medicine</i> 2014;42(7): 1567-1573.
Paterno, M. V., et al. Limb asymmetries in landing and jumping 2 years following anterior cruciate ligament reconstruction. <i>Clinical Journal of Sport Medicine</i> 2007;17(4): 258-262.
Patterson, M. R. and E. Delahunt. A diagonal landing task to assess dynamic postural stability in ACL reconstructed females. <i>Knee</i> 2013;20(6): 532-536.
Pollard, C. D., et al. Altered lower extremity movement variability in female soccer players during side-

step cutting after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 2015;43(2): 460-465.
Pratt, K. A. and S. M. Sigward. Knee Loading Deficits During Dynamic Tasks in Individuals Following Anterior Cruciate Ligament Reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2017;47(6): 411-419.
Pua, Y. H., et al. Associations of isokinetic and isotonic knee strength with knee function and activity level after anterior cruciate ligament reconstruction: A prospective cohort study. <i>Knee</i> 2017;24(5):1067-1074.
Rebeyrotte-Boulègue, I., et al. Isokinetic evaluation of anterior cruciate ligament reconstruction using a free fascia lata graft strengthened by gracilis tendon. <i>Isokinetics and Exercise Science</i> 2005;13(1): 20-24.
Roi, G. S., et al. Time to return to professional soccer matches after ACL reconstruction. <i>Sport Sciences for Health</i> 2006;1(4): 142-145.
Rudroff, T. Functional capability is enhanced with semitendinosus than patellar tendon acl repair. / La capacite physique est amelieoree par la restauration chirurgicale du ligament croise anterieur avec une greffe demi-tendineuse ou rotulienne. <i>Medicine and Science in Sports and Exercise</i> 2003;35(9): 1486-1492.
Sandon, A., et al. Factors associated with returning to football after anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2015;23(9): 2514-2521.
Schmitt, L. C., et al. Strength Asymmetry and Landing Mechanics at Return to Sport after Anterior Cruciate Ligament Reconstruction. <i>Medicine and Science in Sports and Exercise</i> 2015;47(7): 1426-1434.
Schmitt, L. C., et al. The impact of quadriceps femoris strength asymmetry on functional performance at return to sport following anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2012;42(9): 750-759.
Schneider, D. K., et al. A Novel Mass-Spring-Damper Model Analysis to Identify Landing Deficits in Athletes Returning to Sport After Anterior Cruciate Ligament Reconstruction. <i>Journal of Strength and Conditioning Research</i> 2017;31(9): 2590-2598.
Seijas, R., et al. Return to prelesional Tegner level after anatomic anterior cruciate ligament reconstruction. <i>Archives of Orthopaedic and Trauma Surgery</i> 2016;136(12): 1695-1699.
Seon, J. K., et al. A clinical comparison of screw and suture fixation of anterior cruciate ligament tibial avulsion fractures. <i>American Journal of Sports Medicine</i> 2009;37(12): 2334-2339.
Setuain, I., et al. Differential Effects of Two Rehabilitation Programs Following Anterior Cruciate Ligament Reconstruction. <i>J Sport Rehabil</i> 2017;26(6):544-555.
Shabani, B., et al. Gait knee kinematics after ACL reconstruction: 3D assessment. <i>International Orthopaedics</i> 2015;39(6): 1187-1193.
Shah, V. M., et al. Return to play after anterior cruciate ligament reconstruction in National Football League athletes. <i>American Journal of Sports Medicine</i> 2010;38(11): 2233-2239.
Shanbehzadeh, S., et al. Relative muscle contribution of lower extremity muscles during isokinetic single leg squat in patients following reconstruction of the anterior cruciate ligament. <i>Isokinetics and Exercise Science</i> 2014;22(4): 343-349.
Shaw, T., et al. Do early quadriceps exercises affect the outcome of ACL reconstruction? A randomised controlled trial. <i>Australian Journal of Physiotherapy</i> 2005;51(1): 9-17.
Shelbourne, K. D., et al. Arthrofibrosis in acute anterior cruciate ligament reconstruction. The effect of timing of reconstruction and rehabilitation. <i>American Journal of Sports Medicine</i> 1991;19(4): 332-336.
Shelbourne, K. D. and D. A. Foulk. Timing of surgery in acute anterior cruciate ligament tears on the return of quadriceps muscle strength after reconstruction using an autogenous patellar tendon graft. <i>American Journal of Sports Medicine</i> 1995;23(6): 686-689.
Shelbourne, K. D. and T. J. Davis Evaluation of knee stability before and after participation in a functional sports agility program during rehabilitation after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 1999;27(2): 156-161.
Shelbourne, K. D., et al. Ligament stability two to six years after anterior cruciate ligament reconstruction with autogenous patellar tendon graft and participation in accelerated rehabilitation program. <i>American Journal of Sports Medicine</i> 1995;23(5): 575-579.
Shelbourne, K. D. and S. E. Urch. Primary anterior cruciate ligament reconstruction using the contralateral autogenous patellar tendon. <i>American Journal of Sports Medicine</i> 2000;28(5): 651-658.
Sisk, T. D., et al. Effect of electrical stimulation on quadriceps strength after reconstructive surgery of the anterior cruciate ligament. <i>American Journal of Sports Medicine</i> 1987;15(3): 215-220.
Smith, F. W., et al. Subjective functional assessments and the return to competitive sport after anterior

cruciate ligament reconstruction. <i>British Journal of Sports Medicine</i> 2004;38(3): 279-284.
Sonesson, S., et al. Psychological factors are important to return to pre-injury sport activity after anterior cruciate ligament reconstruction: expect and motivate to satisfy. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2017;25(5): 1375-1384.
Sugimoto, D., et al. Single-leg postural stability deficits following anterior cruciate ligament reconstruction in pediatric and adolescent athletes. <i>Journal of Pediatric Orthopaedics. Part B</i> 2016;25(4): 338-342.
Sukur, E., et al. Comparing Transtibial and Anteromedial Drilling Techniques for Single-bundle Anterior Cruciate Ligament Reconstruction. <i>Open Orthopaedics Journal</i> 2016;10: 481-489.
Tambe, A. D., et al. Anterior cruciate ligament insufficiency: does delay in index surgery affect outcome in recreational athletes. <i>International Orthopaedics</i> 30(2): 104-109.
Taradaj, J., et al. The effect of neuromuscular electrical stimulation on quadriceps strength and knee function in professional soccer players: return to sport after ACL reconstruction. <i>Biomed Res Int</i> 2013: 802534.
Tate, J., et al. The Associations Between Hip Strength and Hip Kinematics During a Single Leg Hop in Recreational Athletes Post ACL Reconstruction Compared to Healthy Controls. <i>International Journal of Sports Physical Therapy</i> 2017;12(3): 341-351.
Thomee, R., et al. Variability in leg muscle power and hop performance after anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2012;20(6): 1143-1151.
Thomee, P., et al. Self-efficacy of knee function as a pre-operative predictor of outcome 1 year after anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2008;16(2): 118-127.
Thomeé, R., et al. Return to sports after Anterior Cruciate Ligament reconstruction in women. <i>Sport-Orthopädie - Sport-Traumatologie</i> 2013;29(1): 22-28.
Tibone, J. E. and T. J. Antich A biomechanical analysis of anterior cruciate ligament reconstruction with the patellar tendon. A two year followup. <i>American Journal of Sports Medicine</i> 1988;16(4): 332-335.
Timmins, R. G., et al. Biceps Femoris Architecture and Strength in Athletes with a Previous Anterior Cruciate Ligament Reconstruction. <i>Medicine and Science in Sports and Exercise</i> 2016;48(3): 337-345.
Tjong, V. K., et al. A qualitative investigation of the decision to return to sport after anterior cruciate ligament reconstruction: to play or not to play. <i>American Journal of Sports Medicine</i> 2014;42(2): 336-342.
Trigsted, S. M., et al. Landing mechanics during single hop for distance in females following anterior cruciate ligament reconstruction compared to healthy controls. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2017;25(5): 1395-1402.
Tripp, D. A., et al. Fear of reinjury, negative affect, and catastrophizing predicting return to sport in recreational athletes with anterior cruciate ligament injuries at 1 year postsurgery. <i>Rehabilitation Psychology</i> 2007;52(1): 74-81.
Trulsson, A., et al. Postural orientation in subjects with anterior cruciate ligament injury: development and first evaluation of a new observational test battery. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2010;18(6): 814-823.
Tsaklis, P. and G. Abatzides. ACL rehabilitation program using a combined isokinetic and isotonic strengthening protocol. <i>Isokinetics and Exercise Science</i> 2002;10(4): 211-219.
Tudisco, C., et al. Knee stability, athletic performance and sport-specific tasks in non-professional soccer players after ACL reconstruction: comparing trans-tibial and antero-medial portal techniques. <i>Muscles Ligaments Tendons J</i> 2015;5(3): 175-180.
Tunay, V. B., et al. Quadriceps femoris strength and knee functions in soccer players after anterior cruciate ligament reconstruction: six month follow-up. <i>Turkish Journal of Physiotherapy Rehabilitation</i> 2008;19(1): 10-14.
Ueda, Y., et al. Factors affecting quadriceps strength recovery after anterior cruciate ligament reconstruction with hamstring autografts in athletes. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2017; 25(10):3213-3219.
Vairo, G. L., et al. Neuromuscular and biomechanical landing performance subsequent to ipsilateral semitendinosus and gracilis autograft anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2008;16(1): 2-14.
Walden, M., et al. ACL injuries in men's professional football: a 15-year prospective study on time trends and return-to-play rates reveals only 65% of players still play at the top level 3 years after ACL rupture. <i>British Journal of Sports Medicine</i> 2016;50(12): 744-750.

Webster, K. A. and P. A. Gribble. Time to stabilization of anterior cruciate ligament-reconstructed versus healthy knees in National Collegiate Athletic Association Division I female athletes. <i>J Athl Train</i> 2010;45(6): 580-585.
Webster, K. E. and J. A. Feller. Tibial rotation in anterior cruciate ligament reconstructed knees during single limb hop and drop landings. <i>Clinical Biomechanics (Bristol, Avon)</i> 2012;27(5): 475-479.
Wilk, K. E., et al. The relationship between subjective knee scores, isokinetic testing, and functional testing in the ACL-reconstructed knee. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 1994;20(2): 60-73.
Wilson, W. J. and P. E. Scranton, Jr. Combined reconstruction of the anterior cruciate ligament in competitive athletes. <i>Journal of Bone and Joint Surgery (American Volume)</i> 1990;72(5): 742-748.
Xergia, S. A., et al. Association of the Single-Limb Hop Test With Isokinetic, Kinematic, and Kinetic Asymmetries in Patients After Anterior Cruciate Ligament Reconstruction. <i>Sports Health</i> 2015;7(3): 217-223.
Yamazaki, J., et al. The kinematic analysis of female subjects after double-bundle anterior cruciate ligament reconstruction during single-leg squatting. <i>Journal of Orthopaedic Science</i> 2013;18(2): 284-289.
Yosmaoglu, H. B., et al. Effects of additional gracilis tendon harvest on muscle torque, motor coordination, and knee laxity in ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2011;19(8): 1287-1292.
Zaffagnini, S., et al. Does chronic medial collateral ligament laxity influence the outcome of anterior cruciate ligament reconstruction?: a prospective evaluation with a minimum three-year follow-up. <i>Journal of Bone and Joint Surgery (British Volume)</i> 2011;93(8): 1060-1064.
Zaffagnini, S., et al. Return to sport after anterior cruciate ligament reconstruction in professional soccer players. <i>Knee</i> 2014;21(3): 731-735.
Zouita Ben Moussa, A., et al. Single-leg assessment of postural stability and knee functional outcome two years after anterior cruciate ligament reconstruction. <i>Annals of Physical and Rehabilitation Medicine</i> 2009;52(6): 475-484.
Zwolski, C., et al. The influence of quadriceps strength asymmetry on patient-reported function at time of return to sport after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 2015;43(9): 2242-2249.
Zwolski, C., et al. The Utility of Limb Symmetry Indices in Return-to-Sport Assessment in Patients With Bilateral Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> 2016;44(8): 2030-2038.

Study design n = 79

Albright, J., et al. Anterior Cruciate Ligament Reconstruction in Pediatric and Adolescent Patients Using Quadriceps Tendon Autograft. <i>Sports Medicine and Arthroscopy Review</i> 2016;24(4): 159-169.
Ardern, C. L., et al. Sports participation 2 years after anterior cruciate ligament reconstruction in athletes who had not returned to sport at 1 year: a prospective follow-up of physical function and psychological factors in 122 athletes. <i>American Journal of Sports Medicine</i> 2015;43(4): 848-856.
Baker, C. L., Jr. and J. Graham Intraarticular ACL reconstruction using the patellar tendon: arthroscopic technique. <i>Orthopedics</i> 1993;16(4): 437-441.
Barber, F. A. and S. D. Click. Meniscus repair rehabilitation with concurrent anterior cruciate reconstruction. <i>Arthroscopy</i> 1997;13(4): 433-437.
Barrett, G., et al. Anterior cruciate ligament reconstruction in patients older than 40 years: allograft versus autograft patellar tendon. <i>American Journal of Sports Medicine</i> 2005;33(10): 1505-1512.
Bisciotti, G. N., et al. Return to sports after ACL reconstruction: a new functional test protocol. <i>Muscles Ligaments Tendons</i> 2016;6(4): 499-509.
Bjorklund, K., et al. Validity and responsiveness of the test of athletes with knee injuries: the new criterion based functional performance test instrument. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2009;17(5): 435-445.
Bottoni, C. R., et al. Postoperative range of motion following anterior cruciate ligament reconstruction using autograft hamstrings: A prospective, randomized clinical trial of early versus delayed reconstructions. <i>American Journal of Sports Medicine</i> 2008;36(4): 656-662.
Bynum, E. B., et al. Open versus closed chain kinetic exercises after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 1995;23(4): 401-406.

Chang, E., et al. Repeated bouts of exercise in patients with anterior cruciate ligament reconstruction. <i>Medicine and Science in Sports and Exercise</i> 2014;46(4): 769-775.
Chen, C. H., et al. Arthroscopic anterior cruciate ligament reconstruction with quadriceps tendon-patellar bone autograft. <i>Journal of Trauma</i> 1999;46(4): 678-682.
Chen, C. H., et al. Arthroscopic anterior cruciate ligament reconstruction with quadriceps tendon autograft: clinical outcome in 4-7 years. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2006;14(11): 1077-1085.
Chen, C. H., et al. Arthroscopic anterior cruciate ligament reconstruction with quadruple semitendinosus and gracilis tendon autograft: Surgical technique and clinical results. <i>Journal of Orthopaedic Surgery</i> 1998;6(2): 61-70.
Chen, C. H., et al. Arthroscopic single-bundle anterior cruciate ligament reconstruction with periosteum-enveloping hamstring tendon graft: clinical outcome at 2 to 7 years." <i>Arthroscopy</i> 2010;26(7): 907-917.
Christino, M. A., et al. Psychological Factors Associated With Anterior Cruciate Ligament Reconstruction Recovery. <i>Orthop J Sports Med</i> 2016;4(3): 2325967116638341.
Cimino, F., et al. Anterior cruciate ligament injury: diagnosis, management, and prevention. <i>American Family Physician</i> 2010;82(8): 917-922.
Dai, B., et al. Anterior cruciate ligament reconstruction in adolescent patients: limb asymmetry and functional knee bracing. <i>American Journal of Sports Medicine</i> 2012;40(12): 2756-2763.
Dandy, D. J. Some clinical aspects of reconstruction for chronic anterior cruciate ligament deficiency. <i>Annals of the Royal College of Surgeons of England</i> 1995;77(4): 290-298.
Daruwalla, J. H., et al. Rates and Determinants of Return to Play After Anterior Cruciate Ligament Reconstruction in NCAA Division I College Football Athletes: A Study of the ACC, SEC, and PAC-12 Conferences. <i>Orthop J Sports Med</i> 2014;2(8): 2325967114543901.
Davies, G. J. Individualizing the Return to Sports After Anterior Cruciate Ligament Reconstruction. <i>Operative Techniques in Orthopaedics</i> 2017;27(1): 70-78.
Di Stasi, S. L., et al. Gait patterns differ between ACL-reconstructed athletes who pass return-to-sport criteria and those who fail. <i>American Journal of Sports Medicine</i> 2013;41(6): 1310-1318.
Drogset, J. O., et al. Autologous patellar tendon and quadrupled hamstring grafts in anterior cruciate ligament reconstruction: a prospective randomized multicenter review of different fixation methods. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2010;18(8): 1085-1093.
Fergusson, C. M. Anterior cruciate ligament reconstruction: Middle-third bone/patella tendon/bone autograft. <i>Journal of the Royal College of Surgeons of Edinburgh</i> 2000;45(1): 33-43.
Gaweda, K., et al. Rehabilitation after one-stage anterior cruciate reconstruction and osteochondral grafting. <i>International Orthopaedics</i> 2006;30(3): 185-189.
Gladstone, J. N. and J. R. Andrews. Endoscopic anterior cruciate ligament reconstruction with patella tendon autograft. <i>Orthopedic Clinics of North America</i> 2002;33(4): 701-715, vii.
Graziano, J., et al. Return to Sport for Skeletally Immature Athletes After ACL Reconstruction: Preventing a Second Injury Using a Quality of Movement Assessment and Quantitative Measures to Address Modifiable Risk Factors. <i>Orthop J Sports Med</i> 2017;5(4): 2325967117700599.
Haillotte, G., et al. Early Strength Recovery After Anterior Cruciate Ligament Reconstruction Using The Fascia Lata. <i>Orthopaedics & Traumatology, Surgery & Research.</i> 2017;103(7);1021-1025.
Hartigan, E. H., et al. Preoperative predictors for noncopers to pass return to sports criteria after ACL reconstruction. <i>Journal of Applied Biomechanics</i> 2012;28(4): 366-373.
Hasebe, Y., et al. Anterior-cruciate-ligament reconstruction using doubled hamstring-tendon autograft. <i>J Sport Rehabil</i> 2005;14(4): 279-293.
Heijne, A. and S. Werner. Early versus late start of open kinetic chain quadriceps exercises after ACL reconstruction with patellar tendon or hamstring grafts: a prospective randomized outcome study. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2007;15(4): 402-414.
Hildebrandt, C., et al. Functional assessments for decision-making regarding return to sports following ACL reconstruction. Part I: development of a new test battery. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2015;23(5): 1273-1281.
Hill, G. N. and S. T. O'Leary. Anterior cruciate ligament reconstruction: the short-term recovery using the Knee Injury and Osteoarthritis Outcome Score (KOOS). <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2013;21(8): 1889-1894.
Hirschmann, M. T., et al. Surgical treatment of complex bicruciate knee ligament injuries in elite athletes: what long-term outcome can we expect? <i>American Journal of Sports Medicine</i> 2010;38(6): 1103-1109.

Hui, C., et al. Outcome of anatomic transphyseal anterior cruciate ligament reconstruction in Tanner stage 1 and 2 patients with open physes. <i>American Journal of Sports Medicine</i> 2012;40(5): 1093-1098.
Jagodzinski, M., et al. Biodegradable screw versus a press-fit bone plug fixation for hamstring anterior cruciate ligament reconstruction: a prospective randomized study. <i>American Journal of Sports Medicine</i> 2010;38(3): 501-508.
Jarocka, M. and A. Czaplicki. The influence of therapeutic training on changes in selected biomechanical variables after an anterior cruciate ligament reconstruction. <i>Polish Journal of Sport & Tourism</i> 2013;20(1): 13-18.
Jarvela, T., et al. Simple measurements in assessing muscle performance after an ACL reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2002;32(12): 638-638.
Jonsson, U. and L. Dahlstedt. Anterior cruciate ligament insufficiency treated by combined medial and lateral extra-articular reconstruction. <i>Archives of Orthopaedic and Trauma Surgery</i> 2002;104(2): 94-96.
Karasel, S., et al. Clinical and functional outcomes and proprioception after a modified accelerated rehabilitation program following anterior cruciate ligament reconstruction with patellar tendon autograft. <i>Acta Orthopaedica et Traumatologica Turcica</i> 2010;44(3): 220-228.
Keays, S. L., et al. Muscle strength and function before and after anterior cruciate ligament reconstruction using semitendinosus and gracilis. <i>Knee</i> 2001;8(3): 229-234.
Keays, S. L., et al. Strength and function before and after anterior cruciate ligament reconstruction. <i>Clinical Orthopaedics and Related Research</i> 2000;(373): 174-183.
Keays, S. L., et al. The relationship between knee strength and functional stability before and after anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic Research</i> 2003;21(2): 231-237.
Krych, A., et al. Adverse effect of femoral nerve blockade on quadriceps strength and function after ACL reconstruction. <i>Journal of Knee Surgery</i> 2015;28(1): 83-88.
Lambert, K. L. Vascularized patellar tendon graft with rigid internal fixation for anterior cruciate ligament insufficiency. <i>Clinical Orthopaedics and Related Research</i> 1983;172: 85-89.
Laorueangthana, A., et al. Clinical comparison between six-strand hamstring tendon and patellar tendon autograft in arthroscopic anterior cruciate ligament reconstruction: a prospective, randomized clinical trial. <i>Journal of the Medical Association of Thailand</i> 2009;92(4): 491-497.
Lee, S., et al. Anterior cruciate ligament reconstruction with use of autologous quadriceps tendon graft. <i>Journal of Bone and Joint Surgery (American Volume)</i> 2007;89 Suppl 3: 116-126.
Lentz, T. A., et al. Factors associated with function after anterior cruciate ligament reconstruction. <i>Sports Health</i> 2009;1(1): 47-53.
Lephart, S. M., et al. Quadriceps strength and functional capacity after anterior cruciate ligament reconstruction. Patellar tendon autograft versus allograft. <i>American Journal of Sports Medicine</i> 1993;21(5): 738-743.
Li, H., et al. MRI-based ACL graft maturity does not predict clinical and functional outcomes during the first year after ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2017;25(10): 3171-3178.
Logerstedt, D., et al. Self-reported knee function can identify athletes who fail return-to-activity criteria up to 1 year after anterior cruciate ligament reconstruction: a delaware-oslo ACL cohort study. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2014;44(12): 914-923.
Logerstedt, D., et al. Symmetry restoration and functional recovery before and after anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2013;21(4): 859-868.
Macleod, T. D., et al. Early regeneration determines long-term graft site morphology and function after reconstruction of the anterior cruciate ligament with semitendinosus-gracilis autograft: a case series. <i>International Journal of Sports Physical Therapy</i> 2013;8(3): 256-268.
Marcacci, M., et al. Arthroscopic intra- and extra-articular anterior cruciate ligament reconstruction with gracilis and semitendinosus tendons. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 1998;6(2): 68-75.
McCarroll, J. R., et al. Anterior cruciate ligament injuries in the young athlete with open physes. <i>American Journal of Sports Medicine</i> . 1988;16(1): 44-47.
Mueller, L. M., et al. Which outcome measures should be utilized to determine readiness to play after ACL reconstruction? <i>J Sport Rehabil</i> . 2014;23(2): 158-164.
Otsuka, H., et al. Comparison of three techniques of anterior cruciate ligament reconstruction with bone-patellar tendon-bone graft: Differences in anterior tibial translation and tunnel enlargement with each technique. <i>American Journal of Sports Medicine</i> . 2003;31(2): 282-288.

Paessler, H. H. and D. S. Mastrokalos. Anterior cruciate ligament reconstruction using semitendinosus and gracilis tendons, bone patellar tendon, or quadriceps tendon-graft with press-fit fixation without hardware. A new and innovative procedure. <i>The Orthopedic clinics of North America</i> . 2003;34(1): 49-64.
Panigrahi, R., et al. Outcome of Simultaneous Arthroscopic Anterior Cruciate Ligament and Posterior Cruciate Ligament Reconstruction With Hamstring Tendon Autograft: A Multicenter Prospective Study. <i>Asian Journal of Sports Medicine</i> . 2016;7(1)
Raviraj, A., et al. A comparison of early and delayed arthroscopically-assisted reconstruction of the anterior cruciate ligament using hamstring autograft. <i>Journal of Bone and Joint Surgery (British Volume)</i> . 2010;92(4): 521-526.
Ross, M. D., et al. The relationship between participation restrictions and selected clinical measures following anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> . 2002;10(1): 10-19.
Senanayake, S. M., et al. 3-D kinematics and neuromuscular signals' integration for post ACL reconstruction recovery assessment. Conference Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society. 2013: 7221-7225.
Sepulveda, F., et al. Anterior Cruciate Ligament Injury: Return to Play, Function and Long-Term Considerations. <i>Current Sports Medicine Reports</i> . 2017;16(3): 172-178.
Setuain, I., et al. Acceleration and Orientation Jumping Performance Differences Among Elite Professional Male Handball Players With or Without Previous ACL Reconstruction: An Inertial Sensor Unit-Based Study. <i>Physical Medicine and Rehabilitation</i> . 2015;7(12): 1243-1253.
Shino, K., et al. Anatomic ACL reconstruction: rectangular tunnel/bone-patellar tendon-bone or triple-bundle/semitendinosus tendon grafting. <i>Journal of Orthopaedic Science</i> 2015;20(3): 457-468.
Sigward, S. M., et al. Knee loading asymmetries during gait and running in early rehabilitation following anterior cruciate ligament reconstruction: A longitudinal study. <i>Clinical Biomechanics (Bristol, Avon)</i> . 2016;32: 249-254.
Stearns, K. M. and C. D. Pollard. Abnormal frontal plane knee mechanics during sidestep cutting in female soccer athletes after anterior cruciate ligament reconstruction and return to sport. <i>American Journal of Sports Medicine</i> . 2013;41(4): 918-923.
Tegner, Y., et al. A performance test to monitor rehabilitation and evaluate anterior cruciate ligament injuries. <i>American Journal of Sports Medicine</i> . 1986;14(2): 156-159.
Terry, G. C. and E. W. Janssen. ACL reconstruction with patellar tendon graft. <i>Sports Medicine and Arthroscopy Review</i> . 1997;5(2): 136-140.
Tirmik, U., et al. The results of reconstruction of the ACL using the cross-pin femoral system and four-strand hamstring tendon autografts. <i>Acta Orthopaedica et Traumatologica Turcica</i> . 2011;45(4): 233-239.
Webster, K. E. and J. A. Feller. Younger Patients and Men Achieve Higher Outcome Scores Than Older Patients and Women After Anterior Cruciate Ligament Reconstruction. <i>Clinical Orthopaedics and Related Research</i> . 2017.
Wells, L., et al. Adolescent anterior cruciate ligament reconstruction: a retrospective analysis of quadriceps strength recovery and return to full activity after surgery. <i>Journal of Pediatric Orthopedics</i> 2009;29(5): 486-489.
Wellsandt, E., et al. Limb Symmetry Indexes Can Overestimate Knee Function After Anterior Cruciate Ligament Injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2017;47(5): 334-338.
White, K., et al. Anterior cruciate ligament- specialized post-operative return-to-sports (ACL-SPORTS) training: a randomized control trial. <i>BMC Musculoskeletal Disorders</i> . 2013;14: 108.
White, K., et al. Gait Asymmetries Persist 1 Year After Anterior Cruciate Ligament Reconstruction. <i>Orthop J Sports Med</i> 2013;1(2).
Williams, D., et al. Anterior Cruciate Ligament Functional Sports Assessment. <i>Operative Techniques in Sports Medicine</i> 2016;24(1): 59-64.

Conference abstracts n= 74

Amarasooriya, M., et al. Return to play following anterior cruciate ligament reconstruction: Sri Lankan context. <i>Journal of Science and Medicine in Sport</i> 2015;19: e40.
Archer, K., et al. Impact of preoperative expectations and fear of movement on return to sport and KOOS scores at 6 months following ACL reconstruction. <i>Orthop J Sports Med</i> 2015;3(7).
Ardern, C., et al. Medium-term return to sport outcomes after ACL reconstruction surgery: Uptake,

maintenance or cessation of sport? <i>Journal of Science and Medicine in Sport</i> 2011;14: e97.
Ardern, C., et al. The psychological responses of athletes before, and early after recovery from ACL reconstruction predict returning to the pre-injury level sport. <i>Journal of Science and Medicine in Sport</i> 2013;16: e90.
Brophy, R. H., et al. Recovery of postural stability after anterior cruciate ligament reconstruction. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> 2013;29(10): e183.
Broccatelli, M., et al. Functional recovery after anterior cruciate ligament reconstruction in patients over 40. <i>Journal of Orthopaedics and Traumatology</i> 2011;12: S79.
Buda, R., et al. All-inside anatomical ACL reconstruction: Surgical technique and preliminary results. <i>Journal of Orthopaedics and Traumatology</i> 2012;13: S50.
Cordasco, F. A., et al. All-inside, all-epiphyseal ACL reconstruction in skeletally immature athletes: Incidence of second surgery and two-year clinical outcomes. <i>Orthop J Sports Med</i> 2015;3(7).
Cordasco, F., et al. All-inside, all-epiphyseal ACL reconstruction in skeletally immature athletes: Return to play, incidence of second surgery and two-year clinical outcomes. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> 2015;31(6): e1.
Corradini, C., et al. Kinesthetic recovery during the first 12 months after ACL reconstruction in athletes. <i>Journal of Orthopaedics and Traumatology</i> 2013;14(1): S71.
Culvenor, A., et al. Factors associated with anterior knee pain 12 months follow hamstring tendon autograft ACL reconstruction. <i>Journal of Science and Medicine in Sport</i> 2014;18: e44.
De Carlo, M., et al. ACL surgery and accelerated rehabilitation: Revisited. <i>J Sport Rehabil</i> 1997;6(2): 144-156.
DeMille, P., et al. Quality of movement for athletes six months post ACL reconstruction. <i>Orthop J Sports Med</i> 2016;4(7).
Edison, B. R., et al. Side-step cutting after anterior cruciate ligament reconstruction in pediatric athletes. <i>Clinical Journal of Sport Medicine</i> 2015;25(2): 205.
Eriksson, K. and B. Barenius. Acute vs delayed ACL reconstruction. Early differences and preliminary two year results: A randomized controlled trial. <i>Orthop J Sports Med</i> 2016;4(2).
Failla, M., et al. Outcomes 2 years after ACLR in athletes: Function, return to sport rates, and re-injury rates from the delaware-oslo ACL cohort study. <i>Orthop J Sports Med</i> 2015;3(7).
Gajjar, S. M., et al. The melbourne return to sports score (MRSS)-an assessment tool for return to sports following anterior cruciate ligament reconstruction. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> 2013;29(10): e87-e88.
Gleizes Cervera, S., et al. Double Bundle versus single bundle ACL reconstruction: Isokinetic and functional parameters comparison. <i>Annals of Physical and Rehabilitation Medicine</i> 2011;54: e83.
Goh, M. R., et al. Gait, kinesiophobia and functional scores post anterior cruciate ligament reconstruction. <i>Physiotherapy (United Kingdom)</i> 2015;101: eS461-eS462.
Harris, M. D., et al. Multi-joint compensations during landing and cutting at least one year after return to sport following ACL reconstruction. <i>Journal of Orthopaedic Research</i> 2017;35.
Hartigan, E. Knee function after ACL rupture and reconstruction effects of neuromuscular training. <i>University of Delaware</i> ; 2009.
Kaya, D., et al. Do the patients really need concentrated and/or accelerated rehabilitation program after ACL surgery? A new lightened rehabilitation program: A prospective - Double blind - Randomized-controlled study. <i>Orthop J Sports Med</i> 2014;2(11).
Kılınc, E., et al. Functional, clinical and biomechanical comparison of anterior cruciate ligament reconstruction with anatomical placement of transtibial tunnel placement. <i>Orthop J Sports Med</i> 2014;2(11).
Kim, J. G., et al. Vertical jump test as a functional test after anterior cruciate ligament reconstruction. <i>Orthop J Sports Med</i> 2016;4(7).
Kitaguchi, T., et al. Ipsilateral graft and contralateral ACL rupture following ACL reconstruction in highly active athletes. <i>Physiotherapy (United Kingdom)</i> 2015;101: eS762-eS763.
Kobayashi, T., et al. Relationship between myosin heavy chain expression and leg extension force following anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic Research</i> 2017;35.
Kuenze, C. M. Lower extremity function in active individuals following ACL reconstruction. <i>University of Virginia</i> : 2013: p.229
Lee, J. H., et al. Longitudinal changes of isokinetic strength and dynamic performances in patients with anterior cruciate ligament reconstruction. <i>PM and R</i> 2013;5(9): S212.

McCarroll, J. R., et al. Anterior cruciate ligament injuries in young athletes. Recommendations for treatment and rehabilitation. <i>Sports Medicine</i> . 1995;20(2): 117-127.
McClelland, J., et al. Altered trunk movements during landing in people with anterior cruciate ligament reconstruction. <i>Journal of Science and Medicine in Sport</i> . 2015;19: e60.
McIntosh, A. L., et al. Femoral Nerve Blockade is Associated with Persistent Strength Deficits at Six Months Post ACL Reconstruction in Pediatric and Adolescent Patients. <i>Orthop J Sports Med</i> . 2014;2.
Milewski, M. D., et al. Asymmetric Knee Kinematics And Kinetics After ACL Reconstruction In Adolescent Athletes. <i>Orthop J Sports Med</i> . 2014;2.
Milewski, M. D., et al. Recovery of psychological readiness may differ between genders following ACL reconstruction in adolescent athletes. <i>Orthop J Sports Med</i> . 2016;4(7).
Moon, K. T. Early vs. Delayed treatment of anterior cruciate ligament tears. <i>American Family Physician</i> . 2011;83(7): 842-844.
Murdick, S., et al. Adolescent athletes are unable to maintain preinjury activity levels 1 year after ACL reconstruction. <i>Clinical Journal of Sport Medicine</i> . 2012;22(2): 180.
Nawabi, D. H., et al. Return to Play and Clinical Outcomes after All-Inside, Anterior Cruciate Ligament Reconstruction in Skeletally Immature Athletes. <i>Orthop J Sports Med</i> . 2014;2.
Pace, J. L., et al. Effect of ACL graft type on side-step cutting in young athletes. <i>Orthop J Sports Med</i> . 2016;4(7).
Pace, J. L., et al. Comparison of three-dimensional motion during side-step cutting in pediatric athletes with recent ACL reconstruction and those with no ACL surgical history. <i>Orthop J Sports Med</i> . 2015;3(7).
Pairot De Fontenay, B., et al. Kinematic and kinetic deficits on the non-injured leg after anterior cruciate ligament reconstruction: Analysis of the single-leg squat jump. <i>Physiotherapy (United Kingdom)</i> . 2015;101
Paterno, M. V., et al. Effect of osteochondral and meniscal injury on patient reported outcomes at return to sport following anterior Cruciate ligament reconstruction in young athletes. <i>Orthop J Sports Med</i> . 2015;3(7).
Perraton, L., et al. Modifiable lower limb biomechanics associated with worse knee function after anterior cruciate ligament reconstruction. <i>Journal of Science and Medicine in Sport</i> . 2014;18: e91.
Perraton, L., et al. Quadriceps and hamstring strength, control and activation after anterior cruciate ligament reconstruction: Relationship with knee function. <i>Journal of Science and Medicine in Sport</i> . 2014;18: e91.
Perraton, L., et al. Quadriceps muscle force control is related to knee function 12 months after anterior cruciate ligament reconstruction. <i>Journal of Science and Medicine in Sport</i> . 2013;16: e91-e92.
Pinczewski, L., et al. A fifteen year prospective comparison of patellar and hamstring tendon grafts for ACL reconstruction. <i>Journal of Science and Medicine in Sport</i> . 2010;13: e3-e4.
Pomphrey, M. M., Jr. Aggressive rehabilitation following anterior cruciate reconstruction: an update and protocol. <i>Missouri Medicine</i> . 1992;89(6): 358-361.
Popieluch, M., et al. Comparison between strength of muscles rotating the knee in healthy individuals and patients one year after an ACL reconstruction. <i>Orthop J Sports Med</i> . 2014;2(11).
Pottkotter, K., et al. Improvements in thigh strength symmetry are related to changes in self-reported function after anterior cruciate ligament reconstruction. <i>Orthop J Sports Med</i> . 2015;3(7).
Prince, M., et al. Return to Sport: Does excellent 6-month strength and function following ACL reconstruction predict mid-term outcomes? <i>Arthroscopy: Journal of Arthroscopic and Related Surgery</i> . 2015;31(6)
Queen, R. M. and C. T. Franck. Side-to-side asymmetry during recovery following anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic Research</i> . 2017;35.
Rambaud, A., et al. Functional tests can they help in the decision to return to sports after anterior cruciate ligament? Example with Hop tests. <i>Annals of Physical and Rehabilitation Medicine</i> . 2016;59
Rambaud, A., et al. Interest of a biomechanical analysis of running on a treadmill coupled with functional testing and isokinetic evaluation during return to sport after anterior cruciate ligament reconstruction. <i>Annals of Physical and Rehabilitation Medicine</i> . 2016;59
Reid, D., et al. Adolescent anterior cruciate ligament injuries-a 10 year review. <i>Physiotherapy (United Kingdom)</i> . 2015;101: eS1269.
Reuben, S. S., et al. Effect of initiating a preventative multimodal analgesic regimen upon long-term patient outcomes after anterior cruciate ligament reconstruction for same-day surgery: A 1200-patient case series. <i>Acute Pain</i> . 2005;7(2): 65-73.

Ruffilli, A., et al. ACL reconstruction anatomic all-inside technique: Surgical technique and results. <i>Journal of Orthopaedics and Traumatology</i> . 2013;14(1): S71-S72.
Savalli, L., et al. Interest of the monopodal jump as an indirect means of assessing muscle recovery distance of an ACL reconstruction. <i>Annals of Physical and Rehabilitation Medicine</i> . 2012;55: e70+e76.
Satoh, M., et al. Determining predictors to safely initiate jogging after anterior cruciate ligament reconstruction. <i>Physiotherapy (United Kingdom)</i> . 2011;97: eS1101.
Schmitt, L., et al. ACL Reconstruction: Functional Performance 6 Months Following Return To Sport. <i>Journal of Orthopaedic & Sports Physical</i> . 2011;41(1): A41.
Schmitt, L. C., et al. Functional performance at the time of return to sport following ACL reconstruction: the impact of quadriceps strength asymmetry. <i>Journal of Orthopaedic and Sports Physical Therapy</i> . 2009;39(1): A103-A103.
Semay, B., et al. Evolution of the anteroposterior laxity by GnRB at 6, 9 and 12 months post-surgical anterior cruciate ligament reconstruction. <i>Annals of Physical and Rehabilitation Medicine</i> . 2013;59: e19.
Shousha, T., et al. Energy expenditure and fatigue induced proprioceptive defect following rehabilitation of the reconstructed anterior cruciate ligament. <i>Physiotherapy (United Kingdom)</i> . 2011;97: eS1139-eS1140.
Thomeè, P., et al. Return to pre-injury knee strenuous sports after an acl reconstruction, a report from a rehabilitation outcome registry on patient characteristics. <i>Physiotherapy (United Kingdom)</i> 2015;101: eS1515.
Tiefenböck, T. M., et al. Anatomic ACL all-inside repair for anterior cruciate ligament tear-A prospective study with a minimum follow-up of two years. <i>Sport-Orthopädie - Sport-Traumatologie</i> . 2015;31(2): 169-170.
Ucay, O., et al. Barriers to the return to sport after anterior cruciate ligament tear in operative vs. Conservative patients. <i>Annals of Physical and Rehabilitation Medicine</i> . 2015;58: e68.
Ucay, O., et al. How to evaluate precisely return to sport after anterior cruciate ligament tear with operative or conservative treatment on patients with moderate sport level? <i>Annals of Physical and Rehabilitation Medicine</i> 2016;59: e19.
Webster, K., et al. Effect of fatigue on landing biomechanics following anterior cruciate ligament reconstruction surgery. <i>Journal of Science and Medicine in Sport</i> 2010;13: e26.
Webster, K., et al. Double-bundle versus single-bundle anterior cruciate ligament reconstruction-Is the double bundle technique really better? <i>Journal of Science and Medicine in Sport</i> 2011;14: e107.
Wellsandt E. Lower hop scores related to gait asymmetries after ACL injury: Identifying associations related to the development of early onset knee OA. <i>Osteoarthritis and Cartilage</i> . 2015;23: A279.
White, K. The effects of neuromuscular training on the ability to return-to-activity for the ACL reconstructed athlete. <i>University of Delaware</i> . 2014;153 p-153 p.
White, K., et al. ACL-RSI and KOOS Measures Predict Normal Knee Function after ACL-SPORTS Training. <i>Orthop J Sports Med</i> 2014;2.
Willimon, S. C., et al. Micheli ACL reconstruction in prepubescent youths: A retrospective outcomes study. <i>Orthop J Sports Med</i> 2013;1(4).
Yabroudi, M. A., et al. Comparison of two methods to measure return to sports after anterior cruciate ligament (ACL) reconstruction. <i>Orthop J Sports Med</i> 2013;1(4).
Yilmaz, G. and G. Baltaci. Evaluation of knee strength, functional performance and sports activity level 18-24 months after anterior cruciate ligament reconstruction. <i>Physical Therapy in Sport</i> 2006;7(4): 178-179.
Zaffagnini, S., et al. Does Chronic MCL Laxity in the Setting of ACL Reconstruction Influence Clinical Results? A Prospective Evaluation from Surgery to Minimum 3 years Follow-Up. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> 2011;27(10): e156.
Zehir, S., et al. Evaluation of short term outcomes of transportal femoral cross pin fixation in anterior cruciate ligament reconstruction. <i>Orthop J Sports Med</i> 2014; 2(11).

ACL repair, augmentation, or allograft n= 17

Adachi, N., et al. Harvesting hamstring tendons for ACL reconstruction influences postoperative hamstring muscle performance. <i>Archives of Orthopaedic and Trauma Surgery</i> 2003;123(9): 460-465.
Almqvist, K. F., et al. A long-term study of anterior cruciate ligament allograft reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> 2009;17(7): 818-822.
Arundale, A. J., et al. Report of the Clinical and Functional Primary Outcomes in Men of the ACL-

SPORTS Trial: Similar Outcomes in Men Receiving Secondary Prevention With and Without Perturbation Training 1 and 2 Years After ACL Reconstruction. <i>Clinical Orthopaedics and Related Research</i> . 2017; 475(10):2523-2534.
Fujimoto, E., et al. An early return to vigorous activity may destabilize anterior cruciate ligaments reconstructed with hamstring grafts. <i>Archives of Physical Medicine and Rehabilitation</i> 2004;85(2): 298-302.
Gardinier, E. S., et al. Knee contact force asymmetries in patients who failed return-to-sport readiness criteria 6 months after anterior cruciate ligament reconstruction. <i>American Journal of Sports Medicine</i> 2014;42(12): 2917-2925.
Garrison, J. C., et al. The reliability of the vail sport test as a measure of physical performance following anterior cruciate ligament reconstruction. <i>International Journal of Sports Physical Therapy</i> 2012;7(1): 20-30.
Hall, M. P., et al. Neuromuscular Evaluation With Single-Leg Squat Test at 6 Months After Anterior Cruciate Ligament Reconstruction. <i>Orthop J Sports Med</i> 2015;3(3): 2325967115575900.
Hartigan, E. H., et al. Time line for noncopers to pass return-to-sports criteria after anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 2010;40(3): 141-154.
Lentz, T. A., et al. Comparison of physical impairment, functional, and psychosocial measures based on fear of reinjury/lack of confidence and return-to-sport status after ACL reconstruction. <i>American Journal of Sports Medicine</i> 2015;43(2): 345-353.
Majima, T., et al. Rehabilitation after hamstring anterior cruciate ligament reconstruction. <i>Clinical Orthopaedics and Related Research</i> 2002;397: 370-380.
McGrath, T. M., et al. An Ecological Study of Anterior Cruciate Ligament Reconstruction, Part 1: Clinical Tests Do Not Correlate With Return-to-Sport Outcomes. <i>Orthop J Sports Med</i> . 2016;4(11).
McGrath, T. M., et al. An Ecological Study of Anterior Cruciate Ligament Reconstruction, Part 2: Functional Performance Tests Correlate With Return-to-Sport Outcomes. <i>Orthop J Sports Med</i> . 2017;5(2).
Nawasreh, Z., et al. Do Patients Failing Return-to-Activity Criteria at 6 Months After Anterior Cruciate Ligament Reconstruction Continue Demonstrating Deficits at 2 Years? <i>American Journal of Sports Medicine</i> . 2017;45(5): 1037-1048.
Nyland, J., et al. Lower extremity neuromuscular compensations during instrumented single leg hop testing 2-10 years following ACL reconstruction. <i>Knee</i> . 2014;21(6): 1191-1197.
Risberg, M. A., et al. Assessment of functional tests after anterior cruciate ligament surgery. <i>Journal of Orthopaedic and Sports Physical Therapy</i> 1994;19(4): 212-217.
Sousa, P. L., et al. Return to sport: Does excellent 6-month strength and function following ACL reconstruction predict midterm outcomes? <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> . 2017;25(5): 1356-1363.
Yasuda, K., et al. Effects of initial graft tension on clinical outcome after anterior cruciate ligament reconstruction: autogenous doubled hamstring tendons connected in series with polyester tapes. <i>American Journal of Sports Medicine</i> 1997;25(1): 99-106.

Full text unavailable n= 11

Aglietti, P., et al. A comparison between patellar tendon and doubled semitendinosus/gracilis tendon for anterior cruciate ligament reconstruction. A minimum five-year follow-up. <i>Journal of Sports Traumatology and Related Research</i> 1997;19(2): 57-68.
Brief, L. P. Anterior cruciate ligament reconstruction without drill holes. <i>Arthroscopy</i> 1991;7(4): 350-357.
Comba, D., et al. Arthroscopic reconstruction of the ACL with double-looped semitendinosus and gracilis tendons and fixation with bioabsorbable interference screws. Medium-term results. <i>Journal of Sports Traumatology & Related Research</i> 1999;21(3): 145-154.
Czamara, A. Functional benchmarking of rehabilitation outcomes following anterior cruciate ligament reconstruction. <i>Ortop Traumatol Rehabil</i> 2010;12(6): 519-533.
Gillquist, J. and M. Odensten. Arthroscopic reconstruction of the anterior cruciate ligament. <i>Arthroscopy</i> 1988;4(1): 5-9.
Islam, M. Z., et al. Arthroscopic Anterior Cruciate Ligament Reconstruction with Triplet Autograft of Semitendinosus Tendon. <i>Mymensingh Medical Journal</i> 2017;26(3): 545-550.
Stanish, W. D., et al. Reconstruction of the anterior cruciate ligament with a quadriceps patellar tendon

graft. Preliminary results. <i>Canadian Journal of Applied Sport Sciences</i> . 1984;9(1): 21-24.
Tudisco, C., et al. Arthroscopic anterior cruciate ligament reconstruction with the patellar tendon for chronic laxity: Follow-up of 69 cases. <i>Journal of Sports Traumatology and Related Research</i> 1997;19(4): 198-209.
Wilk, K. E., et al. Anterior cruciate ligament reconstruction rehabilitation: a 12-week follow-up of isokinetic testing in recreational athletes. <i>Isokinetics and Exercise Science</i> 1992;2(2): 82-91.
Yasuda, K., et al. Can anterior cruciate ligament reconstruction surgery using autogenous quadriceps and patellar tendon substitute improve fundamental sports ability? <i>Sports Exercise and Injury</i> 1998;4(1): 22-27.
Zaccherotti, G., et al. Long-term isokinetic evaluation of quadriceps and hamstrings strength following ACL reconstruction. A case-control study. <i>Journal of Sports Traumatology and Related Research</i> 1997;19(3): 141-158.

Duplicate records n = 2

Ellera Gomes, J. L. and L. R. Marczyk Anterior cruciate ligament reconstruction with a loop or double thickness of semitendinosus tendon. <i>American Journal of Sports Medicine</i> 1984;12(3): 199-203.
Knezevic, O. M., et al. Alternating Consecutive Maximum Contraction as a Test of Muscle Function in Athletes Following ACL Reconstruction. <i>J Hum Kinet</i> 2012;35: 5-13.