

Subgroup	No. studies	Effect estimate (95% confidence interval)	I ² statistic
Pediatric			
ACL-R			-
>2 years FU	0		
2-5 years FU	1	0.05 (0.91-0.23)	
5-10 years FU	0		
<10 years FU	1	0.04 (0.01-0.18)	
Non-Op	0		-
>2 years FU			
2-5 years FU			
5-10 years FU			
<10 years FU			
Mixed			-
>2 years FU	0		
2-5 years FU	0		
5-10 years FU	1	0.34 (0.22)	
<10 years FU	0		
Adult:			
ACL-R			77%
>2 years FU	4	0.03 (0.0-0.12)	
2-5 years FU	6	0.04 (0.01-0.08)	
5-10 years FU	1	0.12 (0.08-0.18)	
<10 years FU	1	0.12 (0.06-0.23)	
Non-Op			85%
>2 years FU	2	0.18 (0.13-0.23)	
2-5 years FU	1	0.29 (0.16-0.47)	
5-10 years FU	2	0.20 (0.11-0.31)	
<10 years FU	0		
Mixed			
>2 years FU	0		
2-5 years FU	1	0.10 (0.05-0.19)	
5-10 years FU	0		
<10 years FU	1	0.25 (0.17-0.35)	
Mixed pop.:			
ACL-R			90%
>2 years FU	3	0.00 (0.0-0.03)	
2-5 years FU	9	0.04 (0.02-0.06)	
5-10 years FU	3	0.06 (0.04-0.08)	
<10 years FU	3	0.09 (0.06-0.13)	
Non-Op			83 %
>2 years FU	1	0.06 (0.03-0.12)	
2-5 years FU	3	0.08 (0.04-0.12)	
5-10 years FU	2	0.17 (0.09-0.26)	
<10 years FU	1	0.22 (0.17-0.28)	
Mixed			86%
>2 years FU	2	0.07 (0.04-0.13)	
2-5 years FU	0		
5-10 years FU	1	0.15	

<10 years FU	1	0.31	
--------------	---	------	--

Subgroup numbers of studies, point estimate, CI, I²