

# Effects of yoga on depressive symptoms in people with mental disorders: a systematic review and meta-analysis

Jacinta Brinsley <sup>1</sup>, Felipe Schuch,<sup>2</sup> Oscar Lederman,<sup>3</sup> Danielle Girard,<sup>1</sup> Matthew Smout,<sup>4</sup> Maarten A Immink,<sup>1</sup> Brendon Stubbs,<sup>5</sup> Joseph Firth,<sup>6</sup> Kade Davison,<sup>1</sup> Simon Rosenbaum <sup>7,8</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2019-101242>).

For numbered affiliations see end of article.

## Correspondence to

Jacinta Brinsley, School of Health Sciences, University of South Australia, Adelaide, SA 5001, Australia; [jacintabrinley.aep@gmail.com](mailto:jacintabrinley.aep@gmail.com)

Accepted 19 March 2020  
Published Online First  
18 May 2020

## ABSTRACT

**Objective** To assess whether physically active yoga is superior to waitlist control, treatment as usual and attention control in alleviating depressive symptoms in people with a diagnosed mental disorder recognised by the Diagnostic and Statistical Manual of Mental Disorders (DSM).

**Design** Systematic review and meta-analysis following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

**Data sources** Data were obtained from online databases (MEDLINE, EMBASE, PsychINFO, CENTRAL, EMCARE, PEDro). The search and collection of eligible studies was conducted up to 14 May 2019 (PROSPERO registration No CRD42018090441).

**Eligibility criteria for selecting studies** We included randomised controlled trials with a yoga intervention comprising  $\geq 50\%$  physical activity in adults with a recognised diagnosed mental disorder according to DSM-3, 4 or 5.

**Results** 19 studies were included in the review (1080 participants) and 13 studies were included in the meta-analysis (632 participants). Disorders of depression, post-traumatic stress, schizophrenia, anxiety, alcohol dependence and bipolar were included. Yoga showed greater reductions in depressive symptoms than waitlist, treatment as usual and attention control (standardised mean difference=0.41; 95% CI -0.65 to -0.17;  $p < 0.001$ ). Greater reductions in depressive symptoms were associated with higher frequency of yoga sessions per week ( $\beta = -0.44$ ,  $p < 0.01$ ).

modifiable lifestyle related risk factors, such as low levels of physical activity and high levels of sedentary behaviour.<sup>8</sup> Physical inactivity is associated with higher levels of depressive symptoms in the general population<sup>9</sup> and in those with major depressive disorder.<sup>10</sup> There is a clear need for interventions which are efficacious in improving both physical activity and depressive symptoms<sup>11 12</sup> and multicomponent lifestyle interventions incorporating a combination of physical activity, exercise and diet.<sup>13 14</sup> Treatment guidelines for mental disorders from leading international organisations now recommend the integration of physical activity based interventions as part of routine psychiatric care.<sup>11 15–17</sup> Despite these recommendations, translation of evidence into clinical practice and routine implementation of exercise interventions as part of standard care is limited.<sup>18</sup>

One form of exercise that is increasing in popularity in western culture, particularly among psychiatric patients,<sup>19</sup> is yoga. Yoga is a multicomponent mind–body practice comprising physical postures, movement, breathing control and techniques, relaxation, mindfulness and meditation.<sup>20</sup> Mindfulness, meditation and exercise are all effective in reducing depressive symptoms.<sup>21 22</sup> Yoga practice often includes a combination of each and therefore may have additional benefits beyond a single component (eg, mindfulness, meditation or exercise alone). Furthermore, yoga has demonstrated efficacy in improving exercise adherence and may provide an alternative engagement strategy for people who do not want to engage in conventional forms of exercise.<sup>23</sup>

People living with mental disorders commonly experience depressive symptoms—for example, 25% of people diagnosed with schizophrenia and 81% of people diagnosed with generalised anxiety disorder also have depression.<sup>24 25</sup> Several systematic reviews and meta-analyses have assessed a range of yoga interventions, using various styles of yoga, and have shown improvements in mental health across various mental disorders<sup>26 27</sup> (see [table 1](#)). However, no previous review has investigated the transdiagnostic benefits of yoga. Earlier reviews that focused on yoga for people with depression reported positive effects but all conclude that findings must be interpreted with caution due to the heterogeneity of the yoga interventions and poor methodological reporting.<sup>28–30</sup> This review differs from previous reviews as it includes a range of mental disorder

## INTRODUCTION

Globally, mental disorders are responsible for 32.4% of disability adjusted life years (a year of 'healthy' life lost), placing mental disorders at a distant first in the global burden of disease in terms of years lived with disability.<sup>1</sup> Depressive disorders, such as major depressive disorder, are the leading cause of disability worldwide, affecting more than 340 million people.<sup>2</sup> They are predicted to be the second largest contributor to the global burden of disease by the year 2020.<sup>3</sup> Depressive disorders are highly comorbid with other mental disorders, such as anxiety disorders<sup>4</sup> and psychotic disorders.<sup>5</sup> Physical comorbidities are also present in 65% of people with depressive disorders,<sup>6</sup> commonly including obesity, type 2 diabetes, metabolic syndrome and cardiovascular disease.<sup>7</sup> This is driven in part by



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Brinsley J, Schuch F, Lederman O, et al. *Br J Sports Med* 2021;**55**:992–1000.

**Table 1** Summary of previous literature reviews on yoga and mental disorders

Author	Year	Systematic review	Meta-analysis	Population	Assessment of risk of bias	DSM/ICD diagnostic criteria	Reference to PRISMA statement	General result
Balasubramaniam <i>et al</i> <sup>68</sup>	2013	Yes	No	Varied psychiatric disorders	Yes	No	Yes	High quality evidence for positive effect on depression and sleep
Broderick <i>et al</i> <sup>69</sup>	2015	No	No	Schizophrenia	Yes	Yes	Yes	Evidence too weak
Cramer <i>et al</i> <sup>66</sup>	2013	Yes	Yes	Schizophrenia	Yes	Yes	Yes	Moderate evidence for positive short term effect on quality of life
Cramer <i>et al</i> <sup>70</sup>	2017	Yes	No	Depression	Yes	Yes	Yes	Comparable effects of yoga compared with exercise and medication
Cramer <i>et al</i> <sup>71</sup>	2018	Yes	Yes	PTSD	Yes	No	Yes	Weak recommendation for yoga as an adjunctive treatment
Cramer <i>et al</i> <sup>72</sup>	2018	Yes	Yes	Anxiety	Yes	No	Yes	Moderate evidence for medium–large effect compared with no treatment and active comparator control
Cramer <i>et al</i> <sup>28</sup>	2013	Yes	Yes	Depression	Yes	Yes	Yes	Limited to moderate evidence for improvements in depression and anxiety
Da Silva <i>et al</i> <sup>73</sup>	2009	No	No	Varied psychiatric disorders	No	No	No	Moderate evidence for positive effect on mood disorders
Gallegos <i>et al</i> <sup>74</sup>	2017	No	Yes	PTSD	Yes	No	No	Small to moderate effect on PTSD symptoms
Gong <i>et al</i> <sup>60</sup>	2015	Yes	Yes	Prenatal depression	No	No	No	Low–moderate quality evidence for reducing depressive symptoms
Kirkwood <i>et al</i> <sup>27</sup>	2005	Yes	No	Anxiety	No	No	No	High quality evidence for positive effect on OCD
Pilkington <i>et al</i> <sup>29</sup>	2005	No	No	Depression	No	No	No	Evidence too weak
Uebelacker <i>et al</i> <sup>30</sup>	2010	No	No	Depression	No	No	No	Evidence too weak
Vancampfort <i>et al</i> <sup>75</sup>	2012	Yes	No	Schizophrenia	No	Yes	Yes	Moderate evidence for improved psychopathology, positive and negative symptoms, HRQL

DSM, Diagnostic and Statistical Manual for Mental Disorders; GAD, generalised anxiety disorder; HRQL, health related quality of life; ICD, International Classification of Diseases; OCD, obsessive compulsive disorder; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; PTSD, post-traumatic stress disorder.

diagnoses and considers the transdiagnostic potential for yoga across all mental disorder groups, thus allowing for a more comprehensive assessment of the potential benefits for depressive symptoms across the range of mental disorders.

Hence the primary aim of this review was to determine the effects of physically active yoga on depressive symptoms compared with control in people with any type of mental disorder. Additionally, the review examines the physical health outcomes of yoga interventions for people with mental disorders. We also sought to identify the key components of yoga based interventions that are associated with better outcomes.

## METHODS

This systematic review and meta-analysis was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines<sup>31</sup> and was prospectively registered with the PROSPERO database (ID: CRD42018090441).

### Data sources and searches

The following sources were searched from database inception to 14 May 2019: MEDLINE (Ovid), EMBASE (Ovid), PsychINFO (Ovid), the Cochrane Central Register of Controlled Trials (CENTRAL), EMCARE (Ovid) and the Physiotherapy Evidence Database (PEDro). MeSH headings used were ‘mental disorders’ or ‘mental disease’ AND ‘yoga’, with all branches entered as keywords (Supplementary Table 1). Manual searches were conducted using the reference lists from recovered articles and relevant systematic reviews and meta-analyses.

### Study selection/inclusion criteria

#### Participants

Studies of adults aged 18 years or older with a current Diagnostic and Statistical Manual of Mental Disorders (DSM)-3, 4 or 5<sup>32</sup> diagnosis of any mental disorder, including depressive disorders, anxiety disorders, affective disorders, trauma and stress related

disorders, psychotic disorders, panic disorders and alcohol and substance use disorders were included. Participants with other clinical health conditions in addition to a mental disorder were not excluded.

#### Interventions

Yoga interventions were defined as a structured, formal and premeditated form of physical activity involving the integration of specific body movements (asana) with breathing (pranayama) and/or mindfulness (including meditation),<sup>27 29</sup> where the movement component (physical activity) made up more than 50% of the total intervention.

#### Control conditions

Treatment as usual, waitlist or attention controls were included. Trials that compared one type of physical activity with another (eg, yoga vs aerobic exercise) without an adequate control group were excluded.<sup>33</sup>

#### Outcome measures

The primary outcome measure was change in depressive symptoms across all mental disorders (mean±SD), measured on validated scales, scored from baseline to post-intervention. Secondary outcomes of interest were exercise capacity (eg, measures of cardiorespiratory fitness and strength), biomarkers (eg, cortisol), quality of life, cognition, sleep, anthropometry (eg, body mass index, waist circumference), remission rates, attendance, dropout rates and adverse events.

#### Study design

Only randomised controlled trials (RCTs) were eligible.

#### Exclusion criteria

In the case of multiple publications from the same study, outcome measures were collated to create a ‘complete study’. Trials were

not excluded because of nationality or clinical setting (inpatient, outpatient, community setting or mixed). Studies not available in English were excluded.

### Data extraction

After removal of duplicates, two researchers (JB and OL) independently screened titles and abstracts using Covidence (Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia). Both authors applied eligibility criteria, and a final list of included articles was developed through consensus. A third reviewer (SR) was available for mediation throughout this process.

Data were extracted from the included articles using a pre-determined data extraction form by two authors (JB and DG) and cross-checked for accuracy by a third author (OL). The data extracted included participant and trial characteristics (diagnosis, type of yoga, control condition, programme length, session duration, frequency, intensity, supervision) and yoga effects on mental disorder severity, as per disorder specific scales. Information regarding the yoga interventions was extracted for meta-regression analyses. The type of yoga was determined by the authors' specifications in the protocol. A simple 'yes', 'no' or 'not specified' was used to determine whether the following aspects of yoga were reported in the intervention: specific yoga postures, consistent sequencing of postures, yogic philosophy, meditation, mindfulness ('attention to breath' met this criterion), breathing techniques ('pranayama' or other controlled breathing exercises met this criterion) and/or encouragement of home practice.

### Study quality

Study quality was reported using the PEDro checklist<sup>34</sup> and scores as reported on the PEDro database. Two authors cross-checked 35% of the included studies against PEDro scoring for quality assurance. A study was deemed to be of high quality with a score of 6–10, of fair quality with a score of 4–5 and of low quality with a score <4. The PEDro scores for articles included in this review are presented in table 2.

### Data analysis

Random effects meta-analysis was conducted with Comprehensive Meta-Analysis software (CMA, V.3) in the following steps. Firstly, changes in mental disorder symptoms were calculated using pre- and post-means (SD) of depressive symptom severity scores of the yoga and control groups, calculating the standardised mean difference (SMD) and 95% CI as the effect size. Secondly, subgroup analyses of yoga modality (eg, vinyasa, chair, kripalu), diagnoses, intervention supervision (yes vs home practice) and intensity, as specified by the authors (eg, gentle hatha) were conducted. In the final step, exploratory meta-regression analyses investigating the moderating role of trial duration, number of sessions per week, trial quality, session length and number of components of yoga used in the intervention on the effect size were conducted. In the absence of sufficient data to conduct meta-analyses, a narrative synthesis of results was conducted. Statistical heterogeneity was assessed using the  $I^2$  statistic. An  $I^2$  value >75% was considered to indicate considerable heterogeneity,  $I^2$  50–75% was considered to indicate substantial heterogeneity and  $I^2$  <40% was considered to indicate limited heterogeneity.<sup>35</sup> Publication bias was assessed with Egger<sup>36</sup> and Begg and Mazumdar,<sup>37</sup> and corrected using Duval and Tweedie trim and fill analysis.<sup>38</sup>

## RESULTS

### Search results and flow of trials through the review

A total of 3880 records (excluding duplicates) were identified, and the titles and abstracts were screened. Full text versions were retrieved for 80 articles, of which 19 were eligible for inclusion. Thirteen of those reported changes in depressive symptoms and were included in the meta-analysis. Six studies were excluded from the quantitative analyses as they did not report depression symptom scores. Figure 1 shows the flow of studies through the review and reasons for exclusion.

### Characteristics of included trials

Characteristics of participant sample, primary diagnosis, diagnostic criteria, yoga intervention and control group are shown in table 3. Analyses of physical health outcomes were not performed because of insufficient data.

Of the 19 included RCTs, 12 originated from the USA,<sup>19 39–49</sup> two from India,<sup>50 51</sup> two from Japan<sup>52 53</sup> and one each from China,<sup>54</sup> Germany<sup>55</sup> and Sweden.<sup>56</sup> Nine trials included patients with a depressive disorder (antenatal=5<sup>39–41 43 47</sup>; bipolar depression=1<sup>48</sup>; major depressive disorder=3<sup>44 46 55</sup>), five included a diagnoses of schizophrenia,<sup>19 50–52 57</sup> three with a post-traumatic stress disorder (PTSD) diagnosis,<sup>42 45 49</sup> one with diagnosed alcohol dependence<sup>56</sup> and one with a range of psychiatric disorders.<sup>53</sup> A total of 1080 participants were included across the 19 RCTs; sample sizes of individual studies ranged from 18 to 122 (median=57; mean=57), with 69% of the total sample being women.

Nine studies<sup>39–41 43 45–48 55</sup> reported using the Structured Clinical Interview for DSM for diagnostic purposes. One study used the Mini International Neuropsychiatric Interview,<sup>44</sup> one used the Clinician Administered PTSD Scale for DSM<sup>49</sup> and another used scores above 57 on the PTSD Checklist for DSM.<sup>42</sup> One study used the Structured Clinical Interview for the Positive and Negative Syndrome Scale (PANSS),<sup>19</sup> two studies used the International Classification of Diseases-10,<sup>52 53</sup> two studies used psychiatrist interviews to confirm DSM criteria<sup>50 51</sup> and two studies did not specify how the diagnosis was made.<sup>54 56</sup> Tools used to measure changes in depression symptoms were the Quick Inventory of Depressive Symptomatology (QIDS SR-16),<sup>46–48</sup> Depression Anxiety and Stress Scale (DASS-21),<sup>42</sup> Centre for Epidemiologic Studies Depression Scale (CES-D),<sup>40 41 58</sup> Hamilton Depression Rating Scale (HAM-D/HDRS),<sup>39 55</sup> Beck Depression Inventory (BDI),<sup>44 49</sup> Hospital Anxiety and Depression scale (HAD),<sup>56</sup> PANSS Depression subscale<sup>19</sup> and the Calgary Depression Scale (CDS).<sup>54</sup>

### Interventions


In total, n=578 participants were assigned to yoga and n=502 to control conditions. Seven RCTs utilised attention control groups, including health education,<sup>43 46 47 49</sup> yoga education,<sup>44</sup> social support<sup>40</sup> and bibliotherapy.<sup>48</sup> Seven RCTs utilised waitlist control<sup>19 39 42 45 50 51 59</sup> and five utilised treatment as usual.<sup>41 52 53 55 56</sup> Yoga interventions were on average 2.4 months long (range 1.5–2.5 months) with 1.6 sessions per week (range 1–3 sessions) of a duration of 60 min (range 20–90 min). Types of yoga were hatha,<sup>44 46 48 49 52 53 55</sup> vinyasa,<sup>39</sup> SVYASA (Swami Vivekananda Yoga Anusandhana Samsthana),<sup>50 51</sup> kundalini,<sup>42</sup> kripalu<sup>45</sup> or were unspecified.<sup>19 40 41 43 47 54 56</sup> Seventeen interventions were supervised by yoga practitioners<sup>19 39–49 52–56</sup> and two interventions were primarily unsupervised as they were home practices.<sup>50 51</sup>

**Table 2** Physiotherapy Evidence Database table for included reviews

Trial	Eligibility criteria specified	Random allocation	Concealed allocation to groups	Groups similar at baseline	Blinding of all subjects	Blinding of therapists who administered therapy	Blinding of assessors who measured at least one key outcome	Measures obtained for more than 85% of subjects/adequate follow-up	Intention to treat analysis	Between group statistical comparisons reported	Both point measures and measures of variability reported	PEDro score
Behere <i>et al</i> <sup>51</sup>	Yes	Yes	No	Yes	No	No	Yes	No	No	Yes	Yes	5
Buttner <i>et al</i> <sup>52</sup>	Yes	Yes	No	Yes	No	No	Yes	No	Yes	Yes	Yes	6
Field <i>et al</i> <sup>40</sup>	Yes	Yes	No	Yes	No	No	No	Yes	No	Yes	Yes	5
Field <i>et al</i> <sup>41</sup>	Yes	Yes	No	No	No	No	Yes	No	No	Yes	Yes	4
Hallgren <i>et al</i> <sup>56</sup>	Yes	Yes	No	Yes	No	No	No	No	No	Yes	Yes	4
Ikai <i>et al</i> <sup>57</sup>	Yes	Yes	Yes	Yes	No	No	Yes	No	No	Yes	Yes	6
Ikai <i>et al</i> <sup>53</sup>	Yes	Yes	No	Yes	No	No	Yes	Yes	Yes	Yes	Yes	7
Jindani <i>et al</i> <sup>42</sup>	Yes	Yes	No	Yes	No	No	No	No	No	Yes	Yes	4
Lin <i>et al</i> , 2015 <sup>54</sup>	Yes	Yes	Yes	Yes	No	No	No	No	Yes	Yes	Yes	6
Mitchell <i>et al</i> <sup>43</sup>	Yes	Yes	No	Yes	No	No	No	No	No	Yes	Yes	4
Prathikanti <i>et al</i> <sup>44</sup>	Yes	Yes	Yes	Yes	No	No	Yes	No	Yes	Yes	Yes	7
Reinhardt <i>et al</i> <sup>45</sup>	Yes	Yes	No	No	No	No	Yes	No	No	Yes	Yes	6
Sarubin <i>et al</i> <sup>45</sup>	No	Yes	No	No	No	No	Yes	No	No	Yes	Yes	6
Uebelacker <i>et al</i> <sup>46</sup>	Yes	Yes	No	Yes	No	No	Yes	Yes	Yes	Yes	Yes	7
Uebelacker <i>et al</i> <sup>47</sup>	Yes	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	5
Van der Kolk <i>et al</i> <sup>49</sup>	Yes	Yes	No	Yes	No	No	Yes	Yes	No	Yes	Yes	7
Varambally <i>et al</i> <sup>50</sup>	Yes	Yes	Yes	Yes	No	No	Yes	No	Yes	Yes	Yes	6
Visciglia <i>et al</i> <sup>19</sup>	Yes	Yes	No	Yes	No	No	Yes	No	No	Yes	Yes	5
Weinstock <i>et al</i> <sup>48</sup>	Yes	Yes	No	Yes	No	No	Yes	No	No	Yes	Yes	5

PEDro, Physiotherapy Evidence Database.





**Figure 1** Identification and selection of studies for the review. Adapted from Moher *et al.*<sup>31</sup> PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; RCT, randomised controlled trial.

### Study quality

PEDro scale total scores ranged from 4 to 7, indicating that the studies demonstrated fair to high quality (high=53%, fair=47%) (table 2). Only 21% of trials reported concealed allocation and 37% used an intention to treat analysis.

### Publication bias

The funnel plot for the analysis appeared to be roughly symmetrical. Publication bias was tested using Egger's regression method<sup>36</sup> and the Begg–Mazumdar test,<sup>37</sup> with a  $p$  value  $<0.05$  suggesting the presence of bias. The effect size was not modified by the Duval and Tweedie trim and fill test. Additionally, funnel plots in figure 2 display study specific effect estimates in relation to the SE, assessing the potential presence of publication bias. Heterogeneity was limited ( $I^2=33\%$ ,  $p=0.11$ ).

### Assessment of overall effect

**Primary outcomes: effect of yoga on depressive symptoms in people with a mental disorder**

A moderate effect of yoga on depressive symptoms compared with all control groups was found (SMD=−0.41; 95% CI −0.65 to −0.17,  $p<0.001$ ; heterogeneity:  $I^2=50\%$ ;  $Q=23.86$ ;  $p=0.02$ ; RCT=13; yoga  $n=333$ ; control  $n=299$ ) (figure 2).

### Subgroup analyses

#### Control group

Subgroup analyses based on control groups revealed evidence for a moderate effect of yoga on depressive symptoms compared with waitlist control ( $n=4$ , SMD=−0.58; 95% CI −1.03 to −0.12,  $p<0.05$ ), a small, but not statistically significant, effect compared with treatment as usual ( $n=2$ , SMD=−0.39; 95% CI −1.14 to 0.36,  $p=0.31$ ) and to attention controls ( $n=8$ , SMD=−0.21; 95% CI −0.54 to 0.12,  $p=0.22$ ).

### Diagnostic category

Subgroup analyses based on diagnostic groups revealed evidence for a moderate effect of yoga on depressive symptoms in depressive disorders ( $n=8$ , SMD=−0.40; 95% CI −0.67 to −0.13,  $p<0.01$ ), no effect in PTSD ( $n=2$ , SMD=−0.01; 95% CI, −0.51 to 0.48,  $p=0.95$ ), a small, but not statistically significant, effect in alcohol use disorders ( $n=1$ , SMD=−0.24; 95% CI −1.40 to 0.92,  $p=0.69$ ) and a large effect in schizophrenia ( $n=2$ , SMD=−0.90; 95% CI −1.44 to −0.35,  $p<0.01$ ).

### Secondary outcomes: effect of yoga on physical health

Insufficient data were available to pool the effects of yoga on physical health outcomes. Seven studies included measures of physical health. One study<sup>52</sup> examined cardiometabolic health (fasting blood glucose and total cholesterol) and found no significant differences between the yoga and control groups over the 8 week intervention. Another study by the same authors<sup>53</sup> examined physical fitness (strength) by measuring hand grip strength, lower limb muscle endurance and knee flexion strength to body weight ratio. The yoga group had significantly greater improvements across all three measures compared with the treatment as usual group. Lin *et al.*<sup>57</sup> assessed aerobic fitness with a  $VO_{2max}$  test and found no significant changes between groups. While not statistically significant, Field *et al.*<sup>40</sup> reported reduced cortisol levels within a single session (acute response) for both yoga and social support groups, and increased cortisol levels from baseline to last session (chronic response). Three studies measured physical health/functioning with self-report questionnaires. Uebelacker *et al.*<sup>46</sup> reported no significant improvements for Short Form Health Survey (SF-20) scores between groups. Buttner *et al.*<sup>39</sup> however used the SF-36 to measure health related quality of life and reported that the yoga group experienced a steeper linear increase in scores compared with the waitlist control. Visceglia *et al.*<sup>19</sup> reported that the yoga group had significant improvements in the physical health domain of the World Health Organization Quality of Life Instrument (WHOQOL-BREF) whereas no improvement was reported for the control group.

### Meta-regression

Meta-regression analyses using a mixed effects model revealed that number of sessions per week had a significant effect on depressive symptoms, with higher session frequency leading to a greater improvement in symptoms ( $\beta=-0.44$ ;  $p<0.001$ ; CI −0.66 to −0.21) (Supplementary Table 2). Other intervention variables, such as trial quality, session duration, intervention length, supervision and number of yoga components included (as reported) did not influence depressive symptoms.

### Remission rates

Three studies reported remission rates where symptom severity scores returned to below the diagnostic threshold. van der Kolk *et al.*<sup>49</sup> reported that 52% of participants no longer had clinically significant PTSD symptoms compared with 21% in the health education control group; Prathikanti, *et al.*<sup>44</sup> reported that 60% of participants reached remission in the yoga group compared with 10% in the control group; and Uebelacker *et al.*<sup>46</sup> reported no significant difference in remission rates between the groups at the end of the intervention but at the 3 month follow-up, 39% of participants in the yoga group were in remission compared with 24% in the control group.


### Adverse events, attendance and dropout rates

Seven studies reported on adverse events.<sup>19 44 46–48 53 54</sup> All reported that no adverse events occurred from the intervention. Four studies

**Table 3** Description of the 19 included trials

First author, year	Location, setting	Diagnosis; diagnostic tool	Age (years) (mean (SD))		% Women	Type	Duration and frequency	Home practice	Control	Outcome(s)/statistic
			Y	C						
Behere <i>et al.</i> , 2011 n=60 <sup>31</sup>	India, psychiatric outpatients	Schizophrenia; psychiatrist interview	31.3 (9.3)	33.6 (9.9)	67	SVYASA yoga*	60 min, 5x week for 12 weeks	Yes	WLC	PANSS; mean, SD
Buttner <i>et al.</i> , 2015 n=57 <sup>39</sup>	USA, community	Postpartum depression; SCID and HDRS (>12)	29.8 (5.2)	32.5 (4.8)	100	Vinyasa yoga†	60 min, 2x week for 8 weeks	Yes	WLC	HDRS; mean, SD
Field <i>et al.</i> , 2013 n=92 <sup>40</sup>	USA, community	Prenatal depression; SCID	24.4 (4.7)	24.5 (5.0)	100	Yoga	20 min, 1x week for 12 weeks	NS	Social support	CES-D, salivary cortisol; mean, SD
Field <i>et al.</i> , 2012 n=84 <sup>41</sup>	USA, community	Prenatal depression/dysthymia; SCID	–	–	100	Yoga	20 min, 1x week for 12 weeks	NS	TAU/ massage	CES-D; mean, SD
Hallgren <i>et al.</i> , 2014 n=18 <sup>45</sup>	Sweden, outpatient	Alcohol dependence; DSM criteria	–	–	–	Yoga	90 min, 1x week for 10 weeks	Yes	TAU	HAD-C; mean, SD
Ikai <i>et al.</i> , 2014 n=50 <sup>52</sup>	Japan, outpatients	Schizophrenia F20-F25; ICD-10	53.5 (9.9)	48.2 (12.3)	34	Hatha yoga‡	60 min, 1x week for 8 weeks	No	TAU +walking group	PANSS, FBGL, cholesterol; mean, SD
Ikai <i>et al.</i> , 2017 n=56 <sup>53</sup>	Japan, inpatients	Psychiatric disorders F0-F9; ICD-10	55.5 (11.4)	55 (15.8)	36	Chair hatha yoga	20 min, 2x week for 12 weeks	NS	TAU	PANSS, hand grip strength, lower limb endurance; mean, SD
Jindani <i>et al.</i> , 2015 n=80 <sup>44</sup>	USA, community	PTSD; PCL-17 (>57)	–	–	88	Kundalini yoga¶	90 min, 1x week for 8 weeks	Yes	WLC	DASS-21; mean, SD
Lin <i>et al.</i> , 2015 n=85 <sup>46</sup>	China, community	Psychosis; DSM-IV criteria	23.8 (6.8)	25.3 (8.1)	100	Integrated yoga therapy	60 min, 3x week for 12 weeks	NS	WLC	CDS total, PANSS, V <sub>02</sub> max; mean, SD
Mitchell <i>et al.</i> , 2012 n=24 <sup>43</sup>	USA, community	Prenatal depression; SCID	–	–	100	Yoga	20 min, 2x week for 12 weeks	NS	Education	CES-D total; mean, SD
Prathikanti <i>et al.</i> , 2017 n=38 <sup>45</sup>	USA, community	Major depressive disorder; MINI and BDI (14-28)	43.1 (15.2)	43.8 (14.7)	68	Hatha yoga	90 min, 2x week for 8 weeks	NS	Education	BDI; mean SD
Reinhardt <i>et al.</i> , 2018 n=51 <sup>45</sup>	USA, military	PTSD; SCID-CT	44.1 (14)	46.6 (12.7)	12	Kripalu yoga**	90 min, 2x week for 10 weeks	Yes	WLC	CAPS total; mean, SD
Sarubin <i>et al.</i> , 2014 n=60 <sup>55</sup>	Germany, inpatient	Major depressive disorder; SCID	37.3 (11.9)	42.4 (12.9)	28	Hatha yoga	60 min, 1x week for 5 weeks	–	TAU	HAM-D-21
Uebelacker <i>et al.</i> , 2017 n=122 <sup>46</sup>	USA, community	Major depressive disorder; DSM and SCID	46.8 (12.3)	46.2 (12.1)	84	Hatha yoga	80 min, 1-2x week for 10 weeks	Yes	Healthy living workshop	QIDS, SF-20; mean change?
Uebelacker <i>et al.</i> , 2016 n=20 <sup>47</sup>	USA, community	Antenatal depression; SCID	28.0 (5.9)	28.9 (6.0)	100	Yoga	75 min, 1x week for 9 weeks	Yes	Education	QIDS; mean, SD
Van der Kolk <i>et al.</i> , 2014 n=64 <sup>49</sup>	USA, community	PTSD; CAPS	41.5 (12.2)	44.3 (11.9)	100	Trauma informed yoga	60 min, 1x week for 10 weeks	NS	Education	BDI; mean, SD
Varembally <i>et al.</i> , 2012 n=83 <sup>50</sup>	India, outpatient	Schizophrenia; psychiatrist interview (DSM criteria)	32.8 (1.0)	33.6 (9.5)	13	Yogasana	45 min, 6x week for 12 weeks	Yes	WLC	PANSS; median, IQR
Visciglia <i>et al.</i> , 2011 n=18 <sup>9</sup>	USA, inpatient	Schizophrenia (axis I or II); SCI-PANSS	37.4 (13.7)	48.1 (11.2)	33	Yoga therapy	45 min, 2x week for 8 weeks	NS	WLC	PANSS Depression; mean, SD
Weinstock <i>et al.</i> , 2016 n=18 <sup>48</sup>	USA, community	Bipolar depression; SCID	41.3 (14.3)	33.6 (13.3)	56	Hatha yoga	80 min, 2x week for 10 weeks	Yes	Bibliotherapy	QIDS; mean, SD

N, sample size calculated as number of participants allocated to treatment intervention groups relevant to this study (eg, exercise group excluded from total).  
 \*SVYASA (Swami Vivekananda Yoga Anusandhana Samsthana) refers to the yoga practised from the University Research Institute influenced by Swami Vivekananda.  
 †Included in meta-analysis for depressive symptoms.  
 ‡Vinyasa translates to 'to place (the body) in a special way'; postures are typically flowing and synchronised with breath.  
 §Kundalini yoga focuses on awakening Kundalini energy through practices comprising gentle yoga postures, breathing techniques, meditation and chanting of mantras.  
 ¶Kripalu yoga is a gentle form of hatha yoga, with attention to modifying poses in a way that is best for your body.  
 \*\*Kripalu yoga is a gentle form of hatha yoga, with attention to modifying poses in a way that is best for your body.  
 ††Beck Depression Inventory; CAPS, Clinician Administered PTSD Scale for DSM-5; CDS, Carroll Depression Scale; CES-D, Centre for Epidemiological Studies Depression; DASS, Depression, Anxiety and Stress Scale; DSM, Diagnostic and Statistical Manual for Mental Disorders; FBGL, fasting blood glucose levels; HAD-C, Hospital Anxiety and Depression Scale; HAM-D/HDRS, Hamilton Depression Rating Scale; ICD-10, International Classification of Diseases 10th edition; MINI, Mini International Neuropsychiatric Interview; NS, not stated; PANSS, Positive and Negative Syndrome Scale; PCL-17, PTSD checklist-17; PTSD, post-traumatic stress disorder; QIDS, Quick Inventory of Depressive Symptomatology; SCID, Structured Clinical Interview for DSM-5; SCID-CT, Structured Clinical Interview for DSM-5-Clinical Trials Version; SCI-PANSS, Structured Clinical Interview-Positive and Negative Syndrome Scale; SF-20, Short Form Health Survey; TAU, treatment as usual; WLC, waitlist control.



**Figure 2** Effects of yoga versus treatment as usual, waitlist control or attention control interventions on depressive symptoms across a range of mental disorders, showing estimates of effect size with 95% CIs and relative weight (% weight) for each trial.

reported on some form of intervention fidelity (eg, participant fidelity such as attendance or adherence, or instructor fidelity, such as adherence to intervention protocol) (table 4).<sup>44–48</sup> Dropout rates are reported in table 4. Total dropout rate ranged from 9% to 48%. All but two studies reported dropout rates.<sup>19,43</sup>

## DISCUSSION

The purpose of this systematic review and meta-analysis was to examine the effects that a predominantly physical yoga practice has on depressive symptoms in people with a diagnosed mental disorder. The results demonstrated that yoga has a moderate effect on reducing depressive symptoms across a range of diagnosed mental disorders, thus suggesting yoga may be a viable transdiagnostic intervention for the management of depressive symptoms in people with mental disorders.

The magnitude of the effect of yoga on depressive symptoms compared with treatment as usual, waitlist and attention control combined (SMD = -0.41) was comparable with findings in previous meta-analyses of yoga for depressive symptoms in people with depressive disorders (Cramer *et al*,<sup>28</sup> SMD = -0.69, 95% CI = -0.99 to 0.39; Gong *et al*, SMD = -0.59, 95% CI = -0.94 to -0.25). Gong and colleagues<sup>60</sup> included six trials in their meta-analysis, of which four were also included in the current review. Cramer and colleagues included nine trials in their meta-analysis, of which four were also included in the current review. The variance in results is likely a reflection of the larger sample size with additional studies of high methodological quality, resulting in a more conservative effect. However, the current review has reported the most precise estimate so far, as indicated by narrower confidence intervals. Subgroup analyses

**Table 4** Dropout, attendance and adverse events of the included studies

Study	Dropout	Intervention fidelity	Adverse events
Behere <i>et al</i> <sup>51</sup>	Total 18% (20% yoga; 15% control)	–	–
Buttner <i>et al</i> <sup>39</sup>	Total 11% (14% yoga; 7% control)	–	–
Field <i>et al</i> <sup>40</sup>	Total 14%	–	–
Field <i>et al</i> <sup>41</sup>	Total 25%	–	–
Hallgren <i>et al</i> <sup>56</sup>	Total 22% (33% yoga; 11% control)	–	–
Ikai <i>et al</i> <sup>62</sup>	Total 28%	–	–
Ikai <i>et al</i> <sup>63</sup>	Total 9% (7% yoga; 5% control)	–	0
Jindani <i>et al</i> <sup>42</sup>	37.5% (all from yoga)	–	–
Lin <i>et al</i> , 2015 <sup>54</sup>	Total 23.40%	–	0
Mitchell <i>et al</i> <sup>43</sup>	–	–	–
Prathikanti <i>et al</i> <sup>44</sup>	Total 44%	Total 65% adherence rate (74% yoga; 51% control)	0
Reinhardt <i>et al</i> <sup>45</sup>	Total 48%	–	–
Sarubin <i>et al</i> <sup>55</sup>	Total 12%	–	–
Uebelacker <i>et al</i> <sup>46</sup>	Total 15% (8% yoga; 22% control)	Mean 8.9 (SD 5.1) classes attended/10 min (20 max)	0
Uebelacker <i>et al</i> <sup>47</sup>	Total 15%	Average 92% fidelity of yoga instructor to intervention protocol, 18 item checklist (yoga group only)	0
Van der Kolk <i>et al</i> <sup>49</sup>	1.6% yoga; 4.7% control	–	–
Varambally <i>et al</i> <sup>50</sup>	Total 20%	–	–
Visciglia <i>et al</i> <sup>19</sup>	–	–	0
Weinstock <i>et al</i> <sup>48</sup>	Total 22%	Average 4.8 (SD 5.12) classes attended/10 min (20 max)	0

–, not specified.

in the reviews of both Gong *et al* (2015) and Cramer *et al* (2013) indicated larger effects on depressive symptoms for meditation based yoga interventions compared with physically active yoga interventions. This may partially explain the smaller overall effect seen in the present study, as only physically active yoga interventions were included. While the present review yields a smaller result than physical activity reviews, a moderate effect size indicates that yoga is an efficacious modality of physical activity that should be considered when offering interventions to psychiatric populations, particularly considering the importance of preference and enjoyment for long term adherence and behaviour change.<sup>61</sup> Determining the benefits of mindful exercise, such as yoga, against conventional exercise should be explored further. Preliminary data suggest that combining mindfulness with conventional exercise is effective for reducing depressive symptoms.<sup>62</sup> There is strong evidence to support that both mindfulness<sup>21</sup> and aerobic and/or resistance exercise<sup>14 63</sup> independently reduce depressive symptoms. Exploring whether the effects are strengthened when these are combined, or whether movement allows for greater engagement in mindfulness for novices, is a novel and integral topic. Identifying the mechanisms by which yoga may reduce depressive symptoms is of clinical importance yet is beyond the scope of this review. Nonetheless, a recent comprehensive review<sup>64</sup> of the potential mechanisms by which physical activity may reduce depressive symptoms identified a range of biopsychosocial pathways which are likely to have broad relevance to understanding the mechanisms of yoga.

Extrapolating an effective, and even optimal 'dose' of yoga to facilitate clinically significant change would help inform clinical practice. Further exploration of yoga in comparison with conventional or purposefully non-mindful exercise may help determine effective components of yoga practice. In order to explore a dose–response relationship between yoga practice and the longevity of its benefits, future research should consider collecting follow-up data at varying time points post-intervention. Only six included articles reported follow-up data, with time points ranging from 12 weeks (2 weeks post intervention),<sup>46</sup> 16 weeks (8 weeks post intervention),<sup>51 52</sup> 18 weeks (6 weeks post intervention),<sup>53</sup> 6 months (14 weeks post intervention)<sup>46 56</sup> and 18 months (15 months post intervention).<sup>54</sup> Therefore, conclusions about the potential long term effects of yoga on depressive symptoms were unable to be made.

We found that the number of sessions per week moderated the effect of yoga on depressive symptoms. This is an important finding and should be considered in the design of future yoga interventions targeted at depressive symptoms in people with mental disorders. Interventions should aim to increase the frequency of their sessions per week, as opposed to the duration of each session or the overall duration of the intervention. Further investigation into the duration of effects, and whether particular types of yoga elicit better effects, would be a valuable research question to explore.

Structured exercise in depression and schizophrenia has been shown to have better outcomes when delivered by a qualified health professional.<sup>65 66</sup> However, the impact of qualification in yoga practice and outcomes is unclear. The qualification of yoga practitioners who delivered the interventions in the included RCTs was not reported. Currently, yoga teacher training and education courses are not regulated or standardised, nor accredited within any institutes of higher education.<sup>67</sup> This may have implications when dealing with clinical populations who have chronic and complex conditions that require in-depth

understanding of pathology and psychopathology as well as contraindications and safe prescription of exercise. Furthermore, additional training and upskilling in mental health first aid should be considered for any exercise professional or yoga practitioner delivering interventions for this population. Future research should report the qualification level and type, as well as the experience of the yoga practitioner delivering the intervention, and further analyses should explore the effect this has on intervention outcomes/effectiveness.

Along with efficacy, this review also demonstrated the feasibility of yoga for people with mental disorders. Attendance and dropout did not differ significantly from control groups, with some studies reporting better adherence to yoga than control condition.<sup>46 50 51</sup> A common barrier included 'difficulties making it to classes',<sup>48</sup> thus having yoga services available within inpatient and outpatient mental health facilitates may address this barrier and is likely to have positive effects on symptoms of depression, as well as increasing physical activity levels and promoting greater physical health.

### Limitations

The present review included a range of yoga interventions but due to the relatively small number of trials identified and the poorly defined interventions, it was not possible to conduct separate analyses based on the type and intensity of the interventions. Results from the control and diagnostic category subgroup analyses must also be interpreted with caution because of the small number of studies included. This review did not look at comparisons with true active bona fide interventions and only a small number of studies were available for each kind of control group, thus limiting the comparability of yoga against other interventions for depressive symptoms.

This review has limitations that relate to the primary trials included, such as non-standardised reporting of intervention protocols, lack of intervention description (eg, type of yoga and components of yoga prescribed), poor methodological quality (eg, inadequate blinding of assessors, inadequate follow-up, lack of intention to treat analysis), mix of interviewer rated and self-report outcome measures, and lack of reporting intervention adherence and adverse events. More high quality studies with sufficient sample sizes based on predetermined power analyses are required.

It is essential that future yoga research is reported in a standardised manner whereby the components of yoga can be clearly determined; mindfulness, meditation and physical activity. For further understanding of the mechanism by which yoga has an effect on mental and physical health, intervention variables such as type of yoga, intensity, environment, instructor qualification, specific postures, cueing, philosophical focuses, mindfulness techniques and breathing techniques should be adequately reported. Additionally, the question of longevity of the effects of yoga following intervention cessation requires further investigation; future studies should include long term follow-up. The effects of yoga may be moderated by participant specific factors, such as comorbidities and time since diagnosis, and therefore these factors warrant reporting and investigation in future studies.

Meaningful measures of physical health should be measured so that yoga intervention data can be compared with other modalities of physical activity and independently examined with regard to its physical health benefits, particularly among individuals with a mental disorder.



## CONCLUSION

This review found evidence of a positive effect of yoga beyond usual care for reducing depressive symptoms in people with a range of mental disorders. There was a dose–response relationship between the number of yoga sessions per week and improvements in depressive symptoms. Consideration of yoga as an evidence based exercise modality alongside conventional forms of exercise is warranted given the positive results of this review. Yoga may provide an additional or alternative strategy to engage people experiencing depression in meaningful physical activity.

### What is already known?

- ▶ There is an urgent need to improve physical and mental health for people with diagnosed mental disorders. This is a priority for clinicians, healthcare services, researchers and policy makers.
- ▶ There is evidence to support exercise and mindfulness as singular therapies for improving depressive symptoms.

### What are the new findings?

- ▶ Physically active yoga has a moderate positive effect on improving depressive symptoms in people with a range of mental disorders.
- ▶ Session frequency per week significantly influences the magnitude of depressive symptom reduction.

### Author affiliations

<sup>1</sup>Alliance for Research in Exercise, Nutrition and Activity, University of South Australia: Allied Health and Human Performance, Adelaide, South Australia, Australia

<sup>2</sup>Department of Sports Methods and Techniques, Federal University of Santa Maria, Santa Maria, Brazil

<sup>3</sup>Keeping the Body In Mind, South Eastern Sydney Local Health District, Sydney, New South Wales, Australia

<sup>4</sup>School of Psychology, Social Work and Social Policy, University of South Australia, Adelaide, South Australia, Australia

<sup>5</sup>Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

<sup>6</sup>Division of Psychology and Mental Health, University of Manchester, Manchester, UK

<sup>7</sup>School of Psychiatry, University of New South Wales, Sydney, New South Wales, Australia

<sup>8</sup>Black Dog Institute, Randwick, New South Wales, Australia

**Twitter** Jacinta Brinsley @theyogidoc\_ and Simon Rosenbaum @simon\_rosenbaum

**Contributors** JB and SR conceived the study. OL screened papers. FS, BS and JF assisted with data analysis. All authors contributed to data interpretation and drafting of the manuscript and approved the final version.

**Funding** Simon Rosenbaum is funded by an NHMRC Fellowship APP1123336. Brendon Stubbs is supported by a Clinical Lectureship (ICA-CL-2017-03-001) jointly funded by Health Education England (HEE) and the National Institute for Health Research (NIHR). Brendon Stubbs is part funded by the NIHR Biomedical Research Centre at South London and Maudsley NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the (partner organisation), the NHS, the NIHR or the Department of Health and Social Care. Joseph Firth is supported by a University of Manchester Presidential Fellowship.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

### ORCID iDs

Jacinta Brinsley <http://orcid.org/0000-0002-2588-9649>

Simon Rosenbaum <http://orcid.org/0000-0002-8984-4941>

## REFERENCES

- 1 Vigo D, Thornicroft G, Atun R. Estimating the true global burden of mental illness. *Lancet Psychiatry* 2016;3:171–8.

- 2 Greden JF. The burden of recurrent depression: causes, consequences, and future prospects. *J Clin Psychiatry* 2001;62(Suppl 22):5–9.
- 3 Murray CJ, Lopez AD, World Health Organization. *The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020: summary*, 1996.
- 4 Rebar AL, Stanton R, Rosenbaum S. Comorbidity of depression and anxiety in exercise research. *Lancet Psychiatry* 2017;4:519.
- 5 Buckley PF, Miller BJ, Lehrer DS, et al. Psychiatric comorbidities and schizophrenia. *Schizophr Bull* 2009;35:383–402.
- 6 Royal Australian New Zealand College of Psychiatrists. The economic cost of serious mental illness and comorbidities in Australia and New Zealand. Melbourne RANZCP; 2016.
- 7 Vancampfort D, Stubbs B, Mitchell AJ, et al. Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. *World Psychiatry* 2015;14:339–47.
- 8 Vancampfort D, Firth J, Schuch FB, et al. Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. *World Psychiatry* 2017;16:308–15.
- 9 Farmer ME, Locke BENZ, MOŚCICKI EVEK, et al. Physical activity and depressive symptoms: the NHANES I epidemiologic follow-up study. *Am J Epidemiol* 1988;128:1340–51.
- 10 Schuch F, Vancampfort D, Firth J, et al. Physical activity and sedentary behavior in people with major depressive disorder: a systematic review and meta-analysis. *J Affect Disord* 2017;210:139–50.
- 11 Stubbs B, Vancampfort D, Hallgren M, et al. EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and position statement from the European Psychiatric Association (EPA), supported by the International Organization of Physical Therapists in Mental Health (IOPTMH). *Eur Psychiatry* 2018;54:124–44.
- 12 Firth J, Siddiqi N, Koyanagi A, et al. The Lancet psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *Lancet Psychiatry* 2019;6:675–712.
- 13 Teasdale SB, Ward PB, Rosenbaum S, et al. Solving a weighty problem: systematic review and meta-analysis of nutrition interventions in severe mental illness. *Br J Psychiatry* 2017;210:110–8.
- 14 Schuch FB, Vancampfort D, Richards J, et al. Exercise as a treatment for depression: a meta-analysis adjusting for publication bias. *J Psychiatr Res* 2016;77:42–51.
- 15 National Mental Health Commission. *Equally well consensus statement: improving the physical health and wellbeing of people living with mental illness in Australia*. Sydney, Australia, 2016.
- 16 Malhi GS, Bassett D, Boyce P, et al. Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders. *Aust N Z J Psychiatry* 2015;49:1087–206.
- 17 Kennedy SH, Lam RW, McIntyre RS, et al. Canadian network for mood and anxiety treatments (CANMAT) 2016 clinical guidelines for the management of adults with major depressive disorder: section 3. pharmacological treatments. *Can J Psychiatry* 2016;61:540–60.
- 18 Schuch FB, Morres ID, Ekkekakis P, et al. Exercise works for depression: bridging the implementation gap and making exercise a core component of treatment. *Acta Neuropsychiatr* 2017;29:124–6.
- 19 Visceglia E, Lewis S. Yoga therapy as an adjunctive treatment for schizophrenia: a randomized, controlled pilot study. *J Altern Complement Med* 2011;17:601–7.
- 20 de Manincor M, Bensoussan A, Smith C, et al. Establishing key components of yoga interventions for reducing depression and anxiety, and improving well-being: a Delphi method study. *BMC Complement Altern Med* 2015;15:85.
- 21 Hofmann SG, Sawyer AT, Witt AA, et al. The effect of mindfulness-based therapy on anxiety and depression: a meta-analytic review. *J Consult Clin Psychol* 2010;78:169–83.
- 22 Gotink RA, Chu P, Busschbach JVV, et al. Standardised mindfulness-based interventions in healthcare: an overview of systematic reviews and meta-analyses of RCTs. *PLoS One* 2015;10:e0124344.
- 23 Ussher M, Stanbury L, Cheeseman V, et al. Physical activity preferences and perceived barriers to activity among persons with severe mental illness in the United Kingdom. *Psychiatr Serv* 2007;58:405–8.
- 24 Siris SG. Depression in schizophrenia: perspective in the era of "atypical" antipsychotic agents. *Am J Psychiatry* 2000;157:1379–89.
- 25 Brown TA, Campbell LA, Lehman CL, et al. Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. *J Abnorm Psychol* 2001;110:585–99.
- 26 Cramer H, Lauche R, Klose P, et al. Yoga for schizophrenia: a systematic review and meta-analysis. *BMC Psychiatry* 2013;13:32.
- 27 Kirkwood G, Ramps H, Tuffrey V, et al. Yoga for anxiety: a systematic review of the research evidence. *Br J Sports Med* 2005;39:884–91.
- 28 Cramer H, Lauche R, Langhorst J, et al. Yoga for depression: a systematic review and meta-analysis. *Depress Anxiety* 2013;30:1068–83.
- 29 Pilkington K, Kirkwood G, Ramps H, et al. Yoga for depression: the research evidence. *J Affect Disord* 2005;89:13–24.

- 30 Uebelacker LA, Epstein-Lubow G, Gaudiano BA, *et al.* Hatha yoga for depression: critical review of the evidence for efficacy, plausible mechanisms of action, and directions for future research. *J Psychiatr Pract* 2010;16:22–33.
- 31 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 32 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (DSM-5®)*. Washington, DC: American Psychiatric Pub, 2013.
- 33 Honey EP. I shrunk the pooled SMD! Guide to critical appraisal of systematic reviews and meta-analyses using the Cochrane review on exercise for depression as example. *Ment Health Phys Act* 2015;8:21–36.
- 34 Maher CG, Sherrington C, Herbert RD, *et al.* Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther* 2003;83:713–21.
- 35 The Cochrane Collaboration Higgins JPT, Green S, eds. *Cochrane Handbook for systematic reviews of interventions version 5.1.0*, 2011. www.handbook.cochrane.org
- 36 Egger M, Davey Smith G, Schneider M, *et al.* Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- 37 Begg CB, Mazumdar M. Operating characteristics of a RANK correlation test for publication bias. *Biometrics* 1994;50:1088–101.
- 38 Duval S, Tweedie R. Trim and fill: a simple Funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56:455–63.
- 39 Buttner MM, Brock RL, O'Hara MW, *et al.* Efficacy of yoga for depressed postpartum women: a randomized controlled trial. *Complement Ther Clin Pract* 2015;21:94–100.
- 40 Field T, Diego M, Delgado J, *et al.* Yoga and social support reduce prenatal depression, anxiety and cortisol. *J Bodyw Mov Ther* 2013;17:397–403.
- 41 Field T, Diego M, Hernandez-Reif M, *et al.* Yoga and massage therapy reduce prenatal depression and prematurity. *J Bodyw Mov Ther* 2012;16:204–9.
- 42 Jindani F, Turner N, Khalsa SBS. A yoga intervention for posttraumatic stress: a preliminary randomized control trial. *Evid Based Complement Alternat Med* 2015;2015:351746:1–8.
- 43 Mitchell J, Field T, Diego M, *et al.* Yoga reduces prenatal depression symptoms. *Psychology* 2012;3:782–6.
- 44 Prathikanti S, Rivera R, Cochran A, *et al.* Treating major depression with yoga: a prospective, randomized, controlled pilot trial. *PLoS One* 2017;12:e0173869.
- 45 Reinhardt KM, Noggle Taylor JJ, Johnston J, *et al.* Kripalu yoga for military veterans with PTSD: a randomized trial. *J Clin Psychol* 2018;74:93–108.
- 46 Uebelacker LA, Tremont G, Gillette LT, *et al.* Adjunctive yoga V. health education for persistent major depression: a randomized controlled trial. *Psychol Med* 2017;47:2130–42.
- 47 Uebelacker LA, Battle CL, Sutton KA, *et al.* A pilot randomized controlled trial comparing prenatal yoga to perinatal health education for antenatal depression. *Arch Womens Ment Health* 2016;19:543–7.
- 48 Weinstock LM, Broughton MK, Tezanos KM, *et al.* Adjunctive yoga versus bibliotherapy for bipolar depression: a pilot randomized controlled trial. *Ment Health Phys Act* 2016;11:67–73.
- 49 van der Kolk BA, Stone L, West J, *et al.* Yoga as an adjunctive treatment for posttraumatic stress disorder: a randomized controlled trial. *J Clin Psychiatry* 2014;75:e559–65.
- 50 Varambally S, Gangadhar BN, Thirithalli J, *et al.* Therapeutic efficacy of add-on yogasana intervention in stabilized outpatient schizophrenia: randomized controlled comparison with exercise and waitlist. *Indian J Psychiatry* 2012;54:227–32.
- 51 Behere RV, Arasappa R, Jagannathan A, *et al.* Effect of yoga therapy on facial emotion recognition deficits, symptoms and functioning in patients with schizophrenia. *Acta Psychiatr Scand* 2011;123:147–53.
- 52 Ikai S, Suzuki T, Uchida H, *et al.* Effects of weekly one-hour hatha yoga therapy on resilience and stress levels in patients with schizophrenia-spectrum disorders: an eight-week randomized controlled trial. *J Altern Complement Med* 2014;20:823–30.
- 53 Ikai S, Uchida H, Mizuno Y, *et al.* Effects of chair yoga therapy on physical fitness in patients with psychiatric disorders: a 12-week single-blind randomized controlled trial. *J Psychiatr Res* 2017;94:194–201.
- 54 Lin J, Chan SK, Lee EH, *et al.* Aerobic exercise and yoga improve neurocognitive function in women with early psychosis. *NPJ Schizophr* 2015;1:15047.
- 55 Sarubin N, Nothdurfter C, Schüle C, *et al.* The influence of hatha yoga as an add-on treatment in major depression on hypothalamic-pituitary-adrenal-axis activity: a randomized trial. *J Psychiatr Res* 2014;53:76–83.
- 56 Hallgren M, Romberg K, Bakshi A-S, *et al.* Yoga as an adjunct treatment for alcohol dependence: a pilot study. *Complement Ther Med* 2014;22:441–5.
- 57 JIX L, EHM L, Chang WC, *et al.* Aerobic exercise and yoga hold promises for improving neuro-cognition and symptom in early psychosis. *Schizophr Bull* 2015;1:S320.
- 58 Mitchell KS, Dick AM, DiMartino DM, *et al.* A pilot study of a randomized controlled trial of yoga as an intervention for PTSD symptoms in women. *J Trauma Stress* 2014;27:121–8.
- 59 Lin S-L, Huang C-Y, Shiu S-P, *et al.* Effects of yoga on stress, stress adaption, and heart rate variability among mental health professionals-a randomized controlled trial. *Worldviews Evid Based Nurs* 2015;12:236–45.
- 60 Gong H, Ni C, Shen X, *et al.* Yoga for prenatal depression: a systematic review and meta-analysis. *BMC Psychiatry* 2015;15:14.
- 61 Ekkekakis P. The dual-mode theory of affective responses to exercise in metatheoretical context: I. initial impetus, basic postulates, and philosophical framework. *Int Rev Sport Exerc Psychol* 2009;2:73–94.
- 62 Alderman BL, Olson RL, Brush CJ, *et al.* MAP training: combining meditation and aerobic exercise reduces depression and rumination while enhancing synchronized brain activity. *Transl Psychiatry* 2016;6:e726.
- 63 Hoffman BM, Babyak MA, Craighead WE, *et al.* Exercise and pharmacotherapy in patients with major depression: one-year follow-up of the SMILE study. *Psychosom Med* 2011;73:127–33.
- 64 Kandola A, Ashdown-Franks G, Hendrikse J, *et al.* Physical activity and depression: towards understanding the antidepressant mechanisms of physical activity. *Neurosci Biobehav Rev* 2019;107:525–39.
- 65 Vancampfort D, Stubbs B, Sienaert P, *et al.* What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. *Psychiatr Danub* 2015;27:210.
- 66 Stubbs B, Vancampfort D, Rosenbaum S, *et al.* Dropout from exercise randomized controlled trials among people with depression: a meta-analysis and meta regression. *J Affect Disord* 2016;190:457–66.
- 67 National Assessment and Accreditation Council. *The global yoga accreditation Summit*. New York, 2019.
- 68 Balasubramaniam M, Telles S, Doraiswamy PM. Yoga on our minds: a systematic review of yoga for neuropsychiatric disorders. *Front Psychiatry* 2012;3:117.
- 69 Broderick J, Knowles A, Chadwick J, *et al.* Yoga versus standard care for schizophrenia. *Cochrane Database Syst Rev* 2015:CD010554.
- 70 Cramer H, Anheyer D, Lauche R, *et al.* A systematic review of yoga for major depressive disorder. *J Affect Disord* 2017;213:70–7.
- 71 Cramer H, Anheyer D, Saha FJ, *et al.* Yoga for posttraumatic stress disorder - a systematic review and meta-analysis. *BMC Psychiatry* 2018;18:72.
- 72 Cramer H, Lauche R, Anheyer D, *et al.* Yoga for anxiety: a systematic review and meta-analysis of randomized controlled trials. *Depress Anxiety* 2018;35:830–43.
- 73 da Silva TL, Ravindran LN, Ravindran AV. Yoga in the treatment of mood and anxiety disorders: a review. *Asian J Psychiatr* 2009;2:6–16.
- 74 Gallegos AM, Crean HF, Pigeon WR, *et al.* Meditation and yoga for posttraumatic stress disorder: a meta-analytic review of randomized controlled trials. *Clin Psychol Rev* 2017;58:115–24.
- 75 Vancampfort D, Vansteelandt K, Scheewe T, *et al.* Yoga in schizophrenia: a systematic review of randomised controlled trials. *Acta Psychiatr Scand* 2012;126:12–20.