

267

A SYSTEMATIC REVIEW OF MUSCULOSKELETAL INJURIES IN PROFESSIONAL GOLFERS

¹Patrick Robinson, ¹Iain Murray, ¹Andrew Duckworth, ²Roger Hawkes, ²Danny Glover, ²Nigel Tilley, ²Rob Hillman, ^{1,3}Chris Oliver, ²Andrew Murray. ¹Department of Orthopaedics and Trauma, Edinburgh, UK; ²European Tour Performance Institute, Virginia Water, UK; ³Department of Sports and Exercise/Physical Activity for Health, Edinburgh, UK

10.1136/bjsports-2021-IOC.246

Background The distribution of professional golfing injuries is poorly understood.

Objective The aim of the study was to perform a systematic review to describe the epidemiology of musculoskeletal injuries in professional golfers.

Design A systematic review using Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guidelines. The databases used were Pubmed, SportDiscus and Embase. The inclusion criteria was published observational research articles relating to the incidence or prevalence of musculoskeletal injuries in professional golfers, which were written in the English language and not restricted by age or gender.

Setting Professional golf.

Main Outcome Measurements Data collected included age, sex, data collection methods, diagnosis, region of injury, side of injury, incidence/prevalence of injury, definition of injury, nature of injury, severity of injury, mechanism of injury, risk factors, length of golfing career, injury management and time to return to sport.

Results Of the 1863 studies identified on the initial search, five studies were found to satisfy the inclusion criteria for analysis. The mean age of the golfers in these studies was 34.8 (\pm 3.6) years. The gender of patients in included studies compromised 72% males and 28% females. Four studies reported that lumbar spine injuries were the most common (range 22–34%). Excluding injuries to the spine (lumbar, thoracic and cervical), the hand/wrist was the next most common region of injury (range 6–37%). The quality of the studies was relatively poor with no study satisfying >50% of the quality assessment tool questions and only one study giving a clear definition of how they defined injury.

Conclusions There is a paucity of well-designed epidemiological studies evaluating musculoskeletal injuries affecting professional golfers. Injuries to the spine are the most frequently affected region, followed by the hand/wrist. This study has identified targeted areas of future research that aims to improve the management of injuries among professional golfers.

268

A RANDOMISED CONTROLLED TRIAL INVESTIGATING THE CROSS-EDUCATION OF STRENGTH AND POWER FOLLOWING AT-HOME UNILATERAL CALF EXERCISES

^{1,2}Niall Simmons, ^{1,2}Sumona Mandal, ^{1,2}Liang Zhi Wong, ^{1,2}Andreas Mirallais, ¹Flaminia Ronca, ¹Bhavesh Kumar. ¹The Institute of Sport Exercise and Health, London, UK; ²University College London, London, UK

10.1136/bjsports-2021-IOC.247

Background Cross-education (CE) describes contralateral muscle function gain following unilateral limb training. CE has the potential to reduce strength and power losses following unilateral limb immobilisation. Despite the potential for

wider rehabilitative application, few CE studies have utilised home-based interventions.

Objective Can unilateral home-based strength training elicit bilateral strength and power increases in the calf?

Design A prospective, randomised controlled trial utilising single-blinding.

Setting The Institute of Sport Exercise & Health, London

Participants Thirty-five healthy participants (21.0 ± 1.5 years) with no resistance training 6 months prior to the study were recruited; thirty-four completed the study.

Intervention Participants were randomised to Intervention ($n=20$) or Control ($n=15$). Intervention completed 8 weeks of unilateral straight and bent-knee calf raises at-home, using resistance tubing. Load was self-progressed, sustaining a BORG CR10 score of seven.

Main Outcome Measurements Concentric and eccentric calf peak torque was measured using isokinetic dynamometry at 30°/s. Power output was measured during a single-leg jump using ChronoJump® (Chronojump-Boscosystem). Measurements were taken PRE, MID and POST-intervention.

Results The trained leg increased in strength PRE-POST for concentric ($25.8 \pm 29.3\%$, $p < 0.01$) and eccentric ($19.5 \pm 17.8\%$, $p < 0.001$) contractions. The untrained leg also increased in strength PRE-POST for concentric ($20.5 \pm 25.6\%$, $p < 0.05$) and eccentric ($14.8 \pm 19.2\%$, $p < 0.05$) contractions.

Power increased PRE-POST in the trained leg ($19.3 \pm 4.1\%$, $p < 0.01$). The untrained leg displayed PRE-POST power increases ($18.8 \pm 3.4\%$, $p < 0.01$). Controls displayed no significant strength or power changes.

In the untrained leg, strength increases were significant PRE-MID ($16.3 \pm 16.9\%$, $p < 0.01$) but not MID-POST ($1.3 \pm 13.7\%$, $p > 0.05$). The reverse was true for power with significant increases MID-POST ($10.3 \pm 12.6\%$, $p < 0.01$) but not PRE-MID ($8.3 \pm 13.3\%$, $p > 0.05$).

Conclusions This study demonstrates that unilateral, home-based strength training elicits significant bilateral calf strength and power increases. Additionally, the CE of strength seen prior to power indicates that outcomes secondary to the training type require greater training durations to reach significance. These findings may have rehabilitative potential, however further work is required in clinical populations.

269

COMMERCIALY-AVAILABLE INERTIAL MEASUREMENT UNIT UNDERESTIMATES NUMBER OF JUMPS FOR FEMALES MORE THAN MALES: IMPLICATIONS FOR LOAD MONITORING AND INJURY PREVENTION

¹Tyler J Tait, ¹Lauren C Benson, ¹Kimberley Befus, ¹John Choi, ¹Colin Hillson, ¹Carlyn Stilling, ¹Sagar Grewal, ²Kerry MacDonald, ^{1,3,4,5}Kati Pasanen, ^{1,4,5,6,7}Carolyn Emery. ¹Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Calgary, Canada; ²Volleyball Canada, Ottawa, Canada; ³Tampere Research Center of Sports Medicine, UKK Institute, Tampere, Finland; ⁴Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Canada; ⁵McCraig Bone and Joint Institute, Cumming School of Medicine, University of Calgary, Calgary, Canada; ⁶Department of Community Health Sciences, Cumming School of Medicine, University of Calgary, Calgary, Canada; ⁷Department of Pediatrics, Cumming School of Medicine, University of Calgary, Calgary, Canada

10.1136/bjsports-2021-IOC.248

Background A high incidence of overuse knee injuries among youth basketball players may be attributed to improperly-