031

INJURY SURVEILLANCE AT THE 17/U & 19/U AUSTRALIAN NATIONAL NETBALL CHAMPIONSHIPS AND THE EFFECT OF A FOOT BLISTER PREVENTION INTERVENTION

^{1,2}Erin Smyth, ³Laura Piromalli, ³Alanna Antcliff, ¹Phillip Newman, ^{1,2}Gordon Waddington, ⁴Juanita Weissensteiner, ^{2,5}Michael Drew. ¹University of Canberra, Canberra, Australia; ²Australian Institute of Sport, Canberra, Australia; ³Netball Australia, Canberra, Australia; ⁴New South Wales Office of Sport, Sydney, Australia; ⁵Australian Collaboration for Research into Injury in Sport and its Prevention (ACRISP), Perth, Australia

10.1136/bjsports-2021-IOC.29

Background Pre-elite athletes are at increased risk of injury/illness due to rapid increases in training load, insufficient rest and recovery, growth and a lack of support services compared to elite athletes.

Objective Surveillance of incidence, site, nature and mechanisms of injury/illness at the 2018 & 2019 17/U & 19/U Australian Netball National Championships (ANNC) and reduce injuries at the 2019 ANNCs.

Design Prospective observational cohort study.

Setting 2018 & 2019 17/U & 19/U ANNC's.

Patients (or Participants) One hundred and ninety-two pre-elite athletes were observed each year.

Interventions (or Assessment of Risk Factors) A foot blister prevention pack and advice regarding an injury prevention program were provided to each athlete 6 weeks prior to 2019 ANNC's.

Main Outcome Measurements Injuries were recorded prospectively by team physiotherapists using medical attention data collection methods at the 2018 & 2019 ANNCs.

Results The most frequently recorded medical attention injury diagnoses in 2018 were ankle sprain (n=14, 13.6%), foot blisters (n=11, 10.7%), and lumbar pain (n=10, 9.7%). In 2019 there were 16 (16.8%) ankle sprains, followed by 12 (12.6%) episodes of lumbar pain and 8 (8.4%) foot blisters. In 2018 there were 22 (21%) sports incapacity injuries with ankle sprain (n=4, 18.2%), anterior cruciate ligament (ACL) rupture (n=3, 13.6%) and concussion (n=3, 13.6%) recorded most frequently. In 2019, there were 12 (12.6%) sports incapacity injuries with concussion (n=5, 5.3%) recorded most frequently followed by ACL rupture (n=4, 4.2%). The blister prevention intervention reduced the number of foot blisters at the 2019 ANNCs (IRR 0.73 0.58–0.91, p=0.002).

Conclusions This is the first publication in recent times to articulate tournament injury rates for pre-elite netball athletes. Ankle sprains are the highest medical attention injury in pre-elite netball athletes. This has not changed over the past 30 years indicating current injury prevention interventions for ankle sprain are not effective. Blister prevention packs and advice reduced the number of foot blisters.

032

DOES MENTAL FATIGUE NEGATIVELY AFFECT FUNCTIONAL PERFORMANCE TESTS USED TO SCREEN FOR LOWER EXTREMITY INJURY RISK?

¹Jo Verschueren, ¹Bruno Tassinon, ¹Jeroen Van Cutsem, ¹Bart Roelands, ²Evert Verhagen, ^{1,3}Romain Meeusen. ¹Human Physiology and Sports Physiotherapy Research Group, Faculty of Physical Education and Physiotherapy, Vrije Universiteit Brussel, Brussels, Belgium; ²Amsterdam Collaboration on Health and Safety in Sports, Department of Public and Occupational Health, Amsterdam Movement Sciences, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, Netherlands; ³Strategic Research Program 'Exercise and the Brain in Health and Disease: the added value of Human-Centered Robotics', Vrije Universiteit Brussel, Brussels, Belgium

10.1136/bjsports-2021-IOC.30

Background A lot of controversy exists about how fatigue affects injury risk. Within the fatigue spectrum, mental fatigue impairs visuomotor skills, reaction time and decision-making. Recently, adaptability has been put forward as an important driver in understanding injury occurrence and injury prevention. It is hypothesised that mental fatigue decreases an athlete's adaptability by means of an increased visuomotor reaction time and diminished ability for decision-making and visuomotor skill execution.

Objective To assess the impact of mental fatigue on functional tests used in injury risk profiling and if mental fatigue decreases an athlete adaptability.

Design Randomized counterbalanced cross-over design.

Setting Primary prevention.

Participants Fourteen healthy participants (age = 22 ± 1 years; height = 176.9 ± 8.4 cm; weight = 69.7 ± 10.4 kg).

Interventions Mental fatigue was induced by a 90-minute 100% incongruent Stroop color word test, while the control task included watching a 90-minute documentary.

Main Outcome Measurements Y-balance test (YBT), Reactive balance test (RBT), Single leg hop test (SLH) and counter-movement jump (CMJ) performance was evaluated pre-post mental fatigue. Mental fatigue was evaluated using the M-VAS, Stroop and Eriksen-Flanker task performance outcomes.

Results Mental fatigue was successfully induced at the subjective level with a significant increase in M-VAS (p<0.001), with no decrease in performance on the Stroop and Eriksen-Flanker task. Mental fatigue did not affect YBT, SLH and the CMJ performance. However, RBT accuracy significantly decreased in mental fatigue condition (p=0.024), while the visuomotor reaction time of the RBT remained unaffected.

Conclusions This is the first experiment to illustrate that mental fatigue affects functional adaptability, as accuracy in response to environmental stimuli decreases. However, traditional functional test performance remains unaffected. Therefore, both neurocognitive functional tests and mental fatigue responses should be considered within injury risk profiling strategies. Prospective study designs within the injury prevention domain should include neurocognitive functional tests to investigate the possible relationship of these tests with sports injuries.

033

STRONG PREDICTIVE POWER OF MUSCLE FIBRE TYPOLOGY ON HAMSTRING STRAIN INJURY RISK IN PROFESSIONAL SOCCER

¹Eline Lievens, ¹Kim Van Vossel, ¹Frek Van de Castele, ²Evi Wezenbeek, ²Erik Witvrouw, ¹Wim Derave. ¹Department of Movement and Sports Sciences, Ghent University, Ghent, Belgium; ²Department of Rehabilitation Sciences and Physiotherapy, Ghent University, Ghent, Belgium

10.1136/bjsports-2021-IOC.31

Background Hamstring strain injuries (HSI) are the most prevalent injuries in team sports and occur more frequently in the fatigued phases of a match. In the search for interindividual factors determining increased muscle fatigue-and thus HSI risk-the muscle fibre type composition (fast-twitch vs slow-twitch muscle typology) is a likely candidate.

Objective The objective of this study was to identify whether the muscle fibre typology is a risk factor for HSI.

Design Prospective cohort study over 3 consecutive seasons

Setting First division Belgian (discovery cohort) and UK (replication cohort) soccer.

Patients (or Participants) Players had to be adult out-field players, who did not sustain a HSI in the 6 months before the scan. In total, 96 participants finished the study.

Interventions (or Assessment of Risk Factors) The muscle fibre typology was non-invasively estimated using proton magnetic resonance spectroscopy. During the 1–3 years follow-up period, all sustained injuries were provided by the medical team.

Main Outcome Measurements A cox regression model was used to identify if muscle fibre typology is a risk factor to sustain a HSI. Receiver operating characteristic (ROC) curves were analysed.

Results Belgian players with a fast typology display a 6.7-fold higher risk than slow typology players in experiencing a new HSI (14 HSI; $p=0.024$). This was independently confirmed in the UK cohort (5.0-fold risk; 13 HSI; $p=0.023$). ROC curve analysis revealed an AUC of 0.63 in the total cohort and 0.76 in the discovery cohort, indicating poor to fair sensitivity and specificity.

Conclusions The muscle fibre typology is discovered as a novel and important risk factor to sustain a HSI in professional soccer. This opens a new perspective on the mechanism of HSI, possibly related to more pronounced muscle fatigue in fast typology players. Future personalised prevention strategies could take these interindividual muscle composition differences into account.

034 HIGHER NECK STRENGTH MAY LOWER HEAD ACCELERATION DURING PURPOSEFUL HEADING IN FOOTBALL: A SYSTEMATIC REVIEW

Kerry Peek, James M Elliott, Rhonda Orr. *Faculty of Health Sciences, The University of Sydney, Sydney, Australia*

10.1136/bjsports-2021-IOC.32

Background The long-term effect of repeated head impacts during purposeful heading in football is largely unknown. However, evidence suggests that head acceleration during purposeful heading may be reduced by players having strong neck musculature.

Objective To investigate the relationship between neck strength and head acceleration during purposeful heading in football.

Design This systematic review included a comprehensive search of five electronic databases: EMBASE, MEDLINE, CINAHL, SportsDiscus and Web of Science.

Setting All competitive levels of football regardless of setting (high school, football club or academy) were included.

Patients (or Participants) All football players regardless of age, sex, health status or playing level were included.

Interventions (or Assessment of Risk Factors) Only original research reporting on the relationship between neck strength and head acceleration during purposeful heading in football were included. For experimental research designs, studies were included if neck strengthening was the intervention of interest.

Main Outcome Measurements Neck strength measurement using hand-held dynamometry and head acceleration captured during purposeful heading using accelerometry or video analysis.

Results From an initial search of 1174 potentially eligible papers, five cross-sectional studies met the eligibility criteria for inclusion in this review. A total of 190 participants (67% Female) with a mean age of 19 years (range 15–24 years) were recruited across the five studies. Playing level ranged from high school football to division I and II collegiate level. Evidence from four level IV studies supported a statistically significant relationship between higher neck strength and lower head acceleration during purposeful heading in football ($p<0.05$; $r<-0.5$). The fifth study showed a weak association ($r=-0.10$).

Conclusions Data available from cross-sectional research supports that higher neck strength may lower head acceleration during purposeful heading in football. Further evidence is required to determine the most effective method to strengthen the neck musculature in football players and the impact this may have on the long-term health of players.

035 CAN WE PREDICT RECOVERY AND RE-INJURY FOLLOWING CALF MUSCLE STRAIN INJURY?

¹Brady Green, ²Monica Lin, ¹Anthony Schache, ¹Jodie McClelland, ¹Adam Semciw, ²Andrew Rotstein, ¹Jill Cook, ¹Tania Pizzari. *¹La Trobe Sport and Exercise Medicine Research Centre, Melbourne, Australia; ²Victoria House Imaging, Melbourne, Australia*

10.1136/bjsports-2021-IOC.33

Background Calf muscle strain injuries (CMSI) are prevalent in sport, but investigations into the variable recovery periods and re-injury are lacking.

Objective To determine whether clinical and MRI findings are predictive of recovery and re-injury following CMSI.

Design Retrospective audit of injury data (2014–2017).

Setting Data from the Soft Tissue injury Registry of the Australian Football League (STRAFL).

Patients (or Participants) 149 Australian Football League players.

Interventions (or Assessment of Risk Factors) Blinded radiologists assessed MRI characteristics including: the anatomical location of injury, waviness, and severity of aponeurotic disruption (AD): none (0%), mild (<50%), severe (50%–99%), complete (100%). Clinical factors included: age, mechanism, and injury history.

Main Outcome Measurements The recovery period consisted of four milestones: pain free walking, running at >90% of maximum speed, return to full training, return to play (RTP). Risk factors for re-injury were evaluated for early re-injury (<2 months), and overall (<2 seasons). Multi-variable regression was used to evaluate clinical and MRI predictors of recovery and re-injury.

Results 149 CMSI (114 index, 35 re-injuries) were included: 126 soleus, 17 gastrocnemius, and 6 other. The anatomical location of injury did not impact recovery (all, $p>0.05$). CMSI with severe AD took on average 11.9 days longer to RTP than injuries with no AD ($p=0.003$), and the presence of AD ($p=0.03$) and a running-related mechanism of injury ($p=0.01$) were the strongest predictors of a longer RTP for soleus injuries. Older age ($p=0.001$) and a history of a previous ankle injury ($p=0.03$) were risk factors for early re-injury, while older age ($p=0.01$) and a history of previous CMSI ($p=0.002$) were risk factors for re-injury overall.