Patients (or Participants) 192 elite female netball players from all 16 contesting teams at the NWC (average squad size 12 players).

Interventions (or Assessment of Risk Factors) Medical staff of each team recorded all illness (1440 player days) and training/match days lost during the 10-day tournament.

Main Outcome Measurements Incidence rates (IR) (per 1000 player-days; 95% CI) and severity (% time-loss ≥ 1 day) of all illness.

Results In total 11 illnesses were recorded. The overall crude IR of illness was 7.6 (3.1–12.2). Organ systems most affected was the respiratory- (27%), followed by gastrointestinal- (18%) and urogenital tract (18%). More common illness, by specific diagnosis, was acute upper respiratory infection (18%), non-infective gastroenteritis (18%) and dysmenorrhea (18%). Cause of illness was similar for non-infective (5.6; 1.7–9.4) and infective (2.1; 0.3–4.4) illness ($p=0.227$). Time-loss of ≥ 1 day was reported for 9% of all illness. Illness by player position was common in goal keepers (27%), goal defenders (27%), followed by goal shooters (18%). Illness was reported on a match day in 81% cases (match day with a training session=45%; match day without training=36%).

Conclusions There is a moderate incidence of illness during the 2019 NWC, but we note that >80% of players only report illness on match days. We suggest that players be educated to report illness early (at onset of first symptoms), so that treatment can commence before match day. We recommend implementing illness prevention programs prior to and at future international netball tournaments.

388

THE RELIABILITY AND ASSOCIATION OF THE REPEATED COPENHAGEN ADDUCTION EXERCISE TO GELIC FOOTBALL PLAYERS WITH A HISTORY OF GROIN INJURY: A PILOT STUDY

Rachel Tierney, Ahmad Salma, Ulrik McCarthy Persson. *University College Dublin, Dublin, Ireland*

10.1136/bjsports-2021-IOC.355

Background The Copenhagen adduction exercise (CAE) can significantly reduce the risk of developing Groin injuries (GIs) and may be of clinical utility in being used as a test to assess hip-adductor muscle function.

Objective To examine the relationship between hip-adductor muscle function using a modified, repeated version of the CAE with the adductor squeeze test and the function, sport and recreation subscale of the Copenhagen hip and groin outcome score (HAGOS) in male Gaelic football players (GFP) with a history of GIs. A secondary aim was to assess the inter-rater reliability of the CAE.

Design Two stage pilot study. Stage one assessed the inter-rater reliability of the CAE in active male university students. Stage 2 examined the relationship between hip-adductor muscle function using the CAE with the adductor squeeze test and level of function in male GFP with and without a history of GIs.

Setting University clinical skills laboratory and Gaelic football clubs.

Participants Recruitment on a volunteer basis. Thirteen physically active males over 18 with no history of GIs participated in stage one. Sixty-two males (previous GIs $n=30$, no previous GIs $n=32$) over 18 years, currently playing Gaelic

participated in stage two. To be included into the 'injured' group, participants needed to have a history of GIs diagnosed by a physiotherapist which resulted in complete absence from training for at least two weeks. Participants were excluded if they had an acute injury (<2 weeks) or were unable to participate in training sessions.

Main Outcome Measurements Maximum number of repetitions achieved during the CAE, adductor squeeze test and the HAGOS function, sport and recreation subscale.

Results The CAE test demonstrated good inter-rater reliability (ICC: 0.86, 95% CI: 0.61–0.96, $p<0.001$). Injured and uninjured Gaelic footballers did not differ in CAE or adductor squeeze scores ($p>0.05$). Injured players did not differ in CAE scores between their injured and uninjured leg ($p=0.127$). There was a significant correlation between the CAE score on the uninjured leg and the HAGOS ($\rho=0.405$; $p=0.026$) and adductor squeeze ($\rho=0.458$; $p=0.011$).

Conclusions The CAE has good inter-rater reliability. The repeated CAE test was unable to differentiate between injured and uninjured Gaelic football players or between injured and uninjured legs.

389

ABSTRACT WITHDRAWN

390

CAN A HEAT-AND-MOISTURE EXCHANGER ATTENUATE INFLAMMATORY RESPONSES TO EXERCISE IN SUB-ZERO CONDITIONS?

¹Helen G Hanstock, ¹Alasdair S Tutt, ²Hampus Persson, ¹Erik P Andersson, ³Mats Ainegren, ²Nikolai Stenfors. ¹Swedish Winter Sports Research Centre, Department of Health Sciences, Mid Sweden University, Östersund, Sweden; ²Unit of Medicine, Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden; ³Sports Tech Research Centre, Department of Quality Management and Mechanical Engineering, Mid Sweden University, Östersund, Sweden

10.1136/bjsports-2021-IOC.356

Background Heavy endurance training in sub-zero environments increases risk of exercise-induced asthma. Heat-and-moisture exchangers (HME) can prevent exercise-induced bronchoconstriction but it is not known whether they protect against inflammatory responses to exercise in healthy individuals.

Objective To investigate whether use of an HME during exercise in a sub-zero environment affects post-exercise inflammatory responses.

Design Investigator-blind randomised crossover trial.

Setting Environmental chamber at -15°C.

Participants 23 healthy, trained participants aged 18–53 (15 male, 8 female, VO_{2peak} 57 ± 6 and 50 ± 4 mL/kg/min; mean $\pm$ SD).

Interventions Two experimental trials with and without HME, consisting of 30-min moderate-intensity running followed by a 4-min maximal running time-trial. Plasma samples were obtained pre- and 1h-post-exercise and analysed for a panel of 10 cytokines using a multiplex immunoassay.

Main Outcome Measurements Plasma cytokine concentrations (GM-CSF, IL-1 β , IL-4, IL-5, IL-6, IL-8, IL-10, IL-13, IL-17E/25, TNF- α). Data were log-transformed then analysed using two-way repeated-measures ANOVA; one participant was an extreme outlier and excluded.