Objective To estimate the incidence and proportion of non-contact to total ACL injuries by sex, age, sport, participation level, and exposure type.

Design Systematic review, meta-analysis and meta-regression

Setting Team ball-sports (soccer, rugby, basketball, volleyball, American and Australian football, floorball, lacrosse, hockey, baseball, softball)

Participants Athletes participating in team ball-sports regardless of sex, age, and participation level

Interventions Six electronic databases were searched from inception to July 2020. Cohort studies of team ball-sports reporting number of knee injuries as a function of exposure and injury mechanism were included.

Main Outcome Measurements Proportion of non-contact to total ACL injuries and non-contact ACL injury incidence by unit of exposure, sex, age group, sport, participation level, and exposure type.

Results Forty-three studies covering 12 sports were included. The overall proportion of non-contact to total ACL injuries was 54% (95% CI 47–60%, $I^2=90\%$; females: 62%, 95% CI 54–70%, $I^2=83\%$; males: 47%, 95% CI 39–54%, $I^2=86\%$). Injury incidence was higher in females (0.20 per 1000 player-hours, 95% CI 0.12–0.33, $I^2=92\%$; 0.07 per 1000 player-exposures, 95% CI 0.04–0.14, $I^2=98\%$) than males (0.08 per 1000 player-hours, 95% CI 0.04–0.14, $I^2=94\%$; 0.05 per 1000 player-exposures, 95% CI 0.03–0.08, $I^2=97\%$). Injury incidence during competition was much higher (0.53 per 1000 player-hours, 95% CI 0.35–0.81, $I^2=80\%$; 0.35 per 1000 player-exposures, 95% CI 0.17–0.71, $I^2=96\%$) than training (0.03 per 1000 player-hours (95% CI 0.02–0.05, $I^2=60\%$; 0.02 per 1000 player-exposures, 95% CI 0.01–0.05, $I^2=86\%$). Heterogeneity between studies was generally high.

Conclusions Non-contact ACL injuries represented over half of all ACL injuries sustained. The proportion of non-contact to total ACL injuries and injury incidence were higher in females than males. Injuries mostly occurred in competition settings. More research is required to fully understand the influence of sport, age group, and participation level on injury proportion and incidence. difficult.

400

LONGITUDINAL CHANGES IN CUTTING KINEMATICS IN YOUNG FEMALE ATHLETES ACROSS PUBERTY

¹Lionel Chia, ²Timothy Hewett, ³Kevin Ford, ^{4,5}Gregory Myer, ¹Marnee McKay, ¹Justin Sullivan, ^{1,6}Pappas Evangelos. ¹Sydney School of Health Sciences, The University of Sydney, NSW, Australia; ²Hewett Global Consulting, MN, USA; ³Department of Physical Therapy, High Point University, NC, USA; ⁴Emory Sport Performance and Research Center, GA, USA; ⁵Emory University School of Medicine, GA, USA; ⁶University of Wollongong, NSW, Australia

10.1136/bjsports-2021-IOC.365

Background Young female athletes involved in high-speed cutting and change-of-direction sports are particularly susceptible to anterior cruciate ligament (ACL) injuries. Little is known if maturational changes in cutting technique contribute to the increased injury risk.

Objective To examine longitudinal changes in cutting kinematics in female athletes as they matured through puberty.

Design Longitudinal cohort study

Setting Biomechanics laboratory

Patients (or Participants) One-hundred and seventy-two healthy high-school athletes were tested while in at least two different pubertal stages (pre-, mid-, post-pubertal).

Interventions (or Assessment of Risk Factors) Trunk, hip, and knee sagittal, frontal and transverse plane range-of-motion, peak angles, and angles at initial-contact during an unanticipated 45° cut measured with a laboratory-based optoelectronic motion capture system. Pubertal stages were defined by Tanner Stages.

Main Outcome Measurements Changes in trunk, hip, and knee kinematics as girls mature across puberty (pre-to-midpubertal, pre-to-postpubertal, mid-to-postpubertal).

Results As girls matured through puberty, they displayed a decrease in sagittal-plane hip (1.6–4.4°, $p\leq 0.01$) and knee range-of-motion (ROM) (2.7–2.9°, $p\leq 0.01$), and decreased peak hip (2.9–3.2°, $p\leq 0.02$) and knee flexion angles (2.5–2.9°, $p\leq 0.01$), which is indicative of greater quadriceps dominance. Peak knee abduction angles also increased as girls progressed through puberty (0.9–1.4°, $p\leq 0.006$), suggesting greater ligament dominance. In terms of trunk dominance, there were mixed findings with a decrease in trunk frontal- (2.5–5.7°, $p< 0.01$) and sagittal-plane ROM (1.4–2.0°, $p\leq 0.05$), but an increase in trunk transverse-plane ROM (2.8–3.6°, $p\leq 0.01$) observed as girls mature. Other significant changes in cutting technique were decreased peak trunk flexion (3.8–7.8°, $p\leq 0.01$), and decreased hip flexion (2.9–3.3°, $p\leq 0.01$) and knee flexion angles (2.0–3.0°, $p\leq 0.02$) at initial contact, suggesting a more upright and stiffer cutting posture.

Conclusions As girls mature through puberty, they display a change in cutting strategy characterized by greater quadriceps and ligament dominance. Ameliorating these kinematic changes throughout maturation may further reduce ACL injuries in this high-risk group.

401

BONE BRUISE PATTERN AND MECHANISM OF ANTERIOR CRUCIATE LIGAMENT INJURY IN PROFESSIONAL FOOTBALL PLAYERS: CORRELATION BETWEEN MRI AND VIDEO ANALYSIS

¹Pieter D'Hooghe, ²Alberto Grassi, ³Francesco Della Villa, ¹Khalid Alkhelaifi, ¹Emmanouil Papakostas, ²Filippo Tosarelli, ²Stefano Zaffagnini. ¹Aspetar Hospital, Doha, Qatar; ²IRCCS Istituto Ortopedico Rizzoli, Bologna, Italy; ³Isokinetic Group, Bologna, Italy

10.1136/bjsports-2021-IOC.366

Background The presence of bone bruises (BB), especially in the lateral compartment of the knee, is a common finding in the context of anterior cruciate ligament (ACL) injury. However, different patterns have been described and a precise correlation with mechanism of ACL injury has never been determined.

Objective To correlate the MRI features of ACL injuries with the exact injury mechanism obtained from video-analysis. Such investigation, which has never been performed up to now, provides relevant insights in the understanding ACL injury mechanism and in the genesis of intra-articular lesions such as bone bruises.

Design Fifteen professional football players sustaining an ACL injury (while playing during an official match of First League Championship) were included in the study. The video of injury was obtained from the Television broadcast. Knee Magnetic Resonance Imaging (MRI) was obtained within 7 days