Interventions (or Assessment of Risk Factors) Risk of reinjury and osteoarthritis following knee surgery in professional folk dancers.

Results The dancers suffered 14 knee injuries requiring arthroscopic surgery (3 meniscus tears, 4 anterior cruciate ligament tears one of which is with posterolateral corner tear, 1 posterior cruciate ligament tear, 1 patellar dislocation, 1 infrapatellar bursitis, 2 Hoffa's fat pad syndromes, 2 symptomatic medial plicae) during a ten-year period. Following surgeries, the dancers could restart to perform live on the stage in 19.5 ± 12 (range: 5 to 52) weeks on average. Injuries and postoperative times to return to dance was 56.7 ± 23 (26–108) months to follow-up with the same clinic and same surgeon for the patients. One dancer had reoperation due to meniscus re-tear after 4 years. The rate of reinjury is 7.14% after knee surgery. All of the dancers who underwent arthroscopic knee surgery were evaluated for osteoarthritis according to the Kellgren Lawrence classification. The osteoarthritis were classified as G:0 in 7 patients, G:1 in 3 patients, and G:2 in 4 patients on final knee radiographs. All of the operated patients continued with their careers in dancing.

Conclusions Knee surgeries for the cases mentioned above do not necessarily put an end to their dancing career. This may be attributed to the dancers' balance and coordination skill focus compared to endurance focus during training.

124

BONE MINERAL DENSITY AND ASSOCIATED FACTORS: DO YOUNG FEMALE DANCERS AND OTHER RECREATIONAL SPORT ATHLETES DIFFER?

¹Meghan Critchley, ²Clodagh Toomey, ³Stacey M Lobos, ^{1,2,4}Luz Palacios-Derflingher, ^{1,2,5}Sarah J Kenny, ^{1,2,3,4,5}Carolyn Emery. ¹Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Calgary, Alberta, Canada; ²Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta, Canada; ³Human Performance Laboratory, Faculty of Kinesiology, Calgary, Alberta, Canada; ⁴Department of Community Health Sciences, Cumming School of Medicine, Calgary, Alberta, Canada; ⁵O'Brien Institute for Public Health, University of Calgary, Calgary, Alberta, Canada

10.1136/bjsports-2021-IOC.116

Background As an aesthetic activity, dancers are susceptible to Relative Energy Deficiency in Sport (RED-S), specifically low bone mineral density (BMD). Compared to sedentary groups, dancers have significantly higher BMD, but little is known about how BMD in dancers relates to other athletic populations. This knowledge may better inform comparative values and risk of bone injury.

Objective To evaluate associated factors of BMD among female dancers and sporting athletes.

Design Cross-sectional study.

Setting Human Performance Lab, University of Calgary.

Patients (or Participants) 275 females [138 pre-professional dancers (18.4 years IQR: 15.3, 19.9); 137 recreational sport athletes (23.2 years IQR: 21.9, 24.7)] participated.

Interventions (or Assessment of Risk Factors) Factors evaluated: Age (years), BMI (kg/m^2), supplement intake (Ca+, vitD; yes/no), stress fracture history (yes/no), irregular menses (yes/no), and MRI/bone scan (yes/no), one-year injury history (yes/no), and activity (dance/sport). All two-way interactions were evaluated.

Main Outcome Measurements Whole-body BMD (g/cm^2) was estimated from dual-energy x-ray absorptiometry. Lower extremity (LE) segmentation followed published protocols.

Results The mean whole-body BMD was 1.03 g/cm^2 (SD:0.10) in dancers and 1.14 g/cm^2 (SD:0.08) in athletes. LE BMD was 1.10 g/cm^2 (SD:0.11) in dancers and 1.19 g/cm^2 (SD:0.08) in athletes. Age was positively associated with whole-body BMD ($b=0.008, 97.5\% \text{ CI}; 0.004, 0.012$) and LE BMD ($b=0.007, 97.5\% \text{ CI}; 0.002, 0.010$). Activity modified the relationship between BMI and BMD ($p<0.001$), for example, for a BMI of 20.1 kg/m^2 (mean of participants) the difference between dancers and athletes is -0.389 g/cm^2 [97.5% CI: $-0.566, -0.195$] and -0.381 g/cm^2 [97.5% CI: $-0.552, -0.197$] for whole body and LE BMD respectively. No other variables were associated with BMD.

Conclusions Older age is associated with higher BMD in female pre-professional dancers and sport athletes. The relationship between BMI and BMD depends on activity type for both whole-body and lower extremity sites. Efforts to prevent bone injury should focus on younger dancers and athletes and consider the associations of changing body composition over a training season with low BMD.

125

INJURY ASSOCIATED WITH DANCE EDUCATION: A SYSTEMATIC REVIEW

¹Meghan Critchley, ^{1,2,3}Sarah Kenny, ⁴Ashleigh Ritchie, ^{4,5}Katy Chambers, ⁶Carly D McKay. ¹Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Calgary, Alberta, Canada; ²Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta, Canada; ³O'Brien Institute for Public Health, University of Calgary, Calgary, Alberta, Canada; ⁴Royal Academy of Dance, London, UK; ⁵Trinity Laban Conservatoire of Music and Dance, London, UK; ⁶Department for Health, University of Bath, Claverton Down, Bath, UK

10.1136/bjsports-2021-IOC.117

Background Several studies have investigated injury in specific genres (i.e., ballet, contemporary) and/or levels (i.e., pre-professional, professional) of dance. Less is known about injuries sustained during the teaching of dance, which accounts for the highest proportion of the global dance community.

Objective To determine the risk of injury associated with participation in organized dance education.

Design Systematic review.

Setting Formal dance education (e.g., community classes, dance schools/studios, university programs).

Patients (or Participants) Dance students and teachers.

Main Outcome Measurements Six electronic databases were searched to August 2020 (Medline, EMBASE, SportDiscus, CINAHL, SCOPUS, Cochrane). Inclusion criteria were: original data from dance teacher/student samples, injury and exposure related to formal dance classes, any dance genre. Studies were excluded if no estimate of exposure was reported, if injuries occurred during rehearsal/performance, or if dance was used as a therapeutic intervention or for exercise. Two reviewers independently assessed each paper for inclusion at abstract and full text screening stages. Study quality was assessed using the Joanna Briggs Institute Level of Evidence tool.

Results Twenty-five papers were included.

Most studies ($n=22$) focused on dance students only, two included only dance teachers, and one study included both. The quality of studies ranged from poor to moderate. For students and teachers, most injuries were overuse/chronic and involved the lower limb. Student injury rates were estimated at 0.8–9.3 injuries/1000 h. In dance teachers, a single study described 732 pain reports/1000 h and injury incidence