Main Outcome Measurements Factors being neglected in sports research will be discussed in this study, for example, safety justice is relating to players' risk-taking behaviours during match or training; whether opponents are 'co-workers' and players' safety attitudes towards co-workers can influence players' aggressiveness which relates to injuries and accidents.

Results The framework identified for evaluating OSH awareness will be presented from two dimensions including five themes: rugby management commitment (management safety priority, management safety empowerment and management safety justice) and rugby player involvement (players' safety priority and players' trust in co-workers' safety competence).

Conclusions The findings have theoretical implications for rugby organisations to design a survey to facilitate the development of appropriate behaviour interventions. Furthermore, the framework could be potentially applied in wider sports settings.

247

ABSTRACT WITHDRAWN

248

U.S. RUGBY-7S PLAYERS INJURY INCIDENCE, SEVERITY AND BURDEN EFFECTS BY POSITIONS AND LEVELS OF PLAY

^{1,2,3,4}Victor Lopez, ^{1,2,5}Shen-Ying Richard Ma, ^{1,6}Meryle G Weinstein, ²Patria Hume, ^{7,8,9,10,11}Robert C Cantu, ^{1,12}Christian Victoria, ¹Sophie C Queler, ^{1,13}Khalil JA Webb, ^{1,2,14,15,16}Answorth A Allen. ¹Rugby Research and Injury Prevention Group, affiliate of Hospital for Special Surgery, New York, USA; ²Sports Performance Research Institute New Zealand, Auckland University of Technology, Auckland, New Zealand; ³USA Rugby Empire and New England Geographic Union Rugby Football Unions, New York/Boston, USA; ⁴Northeast Rugby Academy, USA Rugby Development Program and USOC-Community Olympic Development Program, New York/Boston, USA; ⁵University of Missouri, Missouri Orthopaedic Institute and Thompson Laboratory for Regenerative Orthopaedics, Columbia, USA; ⁶New York University, Steinhardt School of Culture, Education and Human Development, New York, USA; ⁷Center for the Study of Traumatic Encephalopathy, Boston University School of Medicine, Boston, USA; ⁸Department of Neurosurgery and Sports Medicine, Emerson Hospital, Concord, USA; ⁹Neurologic Sports Injury Center, Brigham and Women's Hospital, Boston, USA; ¹⁰Concussion Legacy Foundation, Waltham, USA; ¹¹World Rugby, Independent Concussion Group, Dublin, Ireland; ¹²New York University, College of Global Public Health, Urban Epidemiology Lab, New York, USA; ¹³University of Arizona, Tucson, USA; ¹⁴Sports Medicine and Shoulder Service, Hospital for Special Surgery, New York, USA; ¹⁵National Basketball Association, New York Knickerbockers, New York, USA; ¹⁶USA Basketball, Colorado Springs, USA

10.1136/bjsports-2021-IOC.227

Background There are limited injury data for Rugby-7s, and even less data analysed by participation level or days return-to-sport after injury.

Objective To describe injury incidence, severity, and injury burden for three levels of Rugby-7s competition.

Design Prospective descriptive epidemiology study.

Setting U.S. Rugby-7s tournaments/series and championships (n=57; 2010–2014) over 72 tournament days; L1 elite, L2 sub elite, and L3 under-19/college/senior games (exposure=14,591 player-hours).

Participants 24,538 U.S. players (men=17,770; women=6,768; age 13–54 years).

Assessment of Risk Factors Intrinsic and extrinsic risk in match injuries.

Main measurement outcome Incidence (per/1000 player-hour (ph)) and mechanism of injury were captured using Rugby

Injury Survey & Evaluation (RISE) report methodology. Time-loss injuries, injury severity (days=d) from training/competition (including post tournament) were documented, and injury burden were calculated.

Results Injury incidence (n=491) was not significantly different between levels (L3:30.74/1000ph, CI:27.26–34.54; L2:36.24/1000ph, CI:30.84–42.31; L1:41.78/1000ph, CI:30.8–55.39). Mean injury burden was significantly lower for L3 than L1 and L2 (L3-vs-L2, $P<0.001$; L3 vs L1, $P<0.001$). Greater risk of lower limb injuries was noted in L3-vs-L1 (RR:0.59, CI:0.38–0.95, $P=0.024$). The cohort sustained high head/neck injury rates (22.6%; 13.3/1000ph). Backs had more injuries among levels than forwards (L1 backs 51.8/1000ph, forwards 26.4/1000ph, $P=0.034$; L2 backs 37.7/1000ph, forwards 29.6/1000ph, $P=0.152$; L3, backs 32.76/1000ph, forwards 24.8/1000ph, $P=0.029$; total cohort backs 35.74/1000ph, forwards 26.39/1000ph, RR:1.35; CI:1.12–1.65, $P=0.002$). Average days absent post injury=44.0d (37.8–50.1d) in 68.4% with follow-up data. A significant difference ($P=0.018$) in mean severity days absent from sport was between the L3 (57.1d) and L2 (27.9d) forwards.

Conclusions Competition level and playing position had significant effects on injury burden and nature of injury. The L1 and L2 had higher injury burden than the lower L3. The L1-vs-L3 cohort had a high proportion of head/neck injury risk compared with other injury locations. Backs sustained greater injury incidence rates among all three levels as compared to forwards.

249

THE EPIDEMIOLOGY OF HEAD, NECK AND FACE INJURIES OF ADULT MEN'S AND WOMEN'S U.S. RUGBY-7S PLAYERS

^{1,2,3}Shen-Ying Richard Ma, ^{2,4}Christian Victoria, ⁴Danielle C Ompad, ^{2,5}Meryle G Weinstein, ^{6,7,8,9,10}Robert C Cantu, ^{11,2}Jordan Genece, ^{2,3,12,13,14}Answorth A Allen, ^{2,3,15,16}Victor Lopez. ¹University of Missouri, Missouri Orthopaedic Institute and Thompson Laboratory for Regenerative Orthopaedics, Columbia, USA; ²Rugby Research and Injury Prevention Group, Hospital for Special Surgery, New York, USA; ³Sports Performance Research Institute New Zealand, Auckland University of Technology, Auckland, New Zealand; ⁴New York University, College of Global Public Health, Urban Epidemiology Lab, New York, USA; ⁵New York University, Steinhardt School of Culture, Education and Human Development, New York, USA; ⁶Center for the Study of Traumatic Encephalopathy, Boston University School of Medicine, Boston, USA; ⁷Department of Neurosurgery and Sports Medicine, Emerson Hospital, Concord, USA; ⁸Sports Neurology and Concussion Program, Brigham and Women's Hospital, Boston, USA; ⁹Concussion Legacy Foundation, Waltham, USA; ¹⁰World Rugby, Independent Concussion Group, Dublin, Ireland; ¹¹Mt. Sinai Hospital, Department of Dermatology, New York, USA; ¹²Sports Medicine and Shoulder Service, Hospital for Special Surgery, New York, USA; ¹³National Basketball Association, New York Knickerbockers, New York, USA; ¹⁴USA Basketball, Colorado Springs, USA; ¹⁵USA Rugby Empire and New England Geographic Union RFUs, New York/Boston, USA; ¹⁶Northeast Rugby Academy, USA Rugby Development Program and USOC-Community Olympic Development Program, New York/Boston, USA

10.1136/bjsports-2021-IOC.228

Background Head and neck injuries are a collision sport concern, however there is a lack of Rugby-7s injury data, particularly in emerging Rugby nations.

Objective To determine the head, neck, and face (HNF) match injury rates and characteristics in US Rugby-7s.

Design Prospective descriptive epidemiology study.

Setting USA Rugby-sanctioned tournaments/series and championships (2010–2016).