Background Previous research has shown that muscle imbalances may be an important risk factor for lower limb injuries in runners. No studies have evaluated hip musculature imbalance in sprinters. Identifying pre-season strength and imbalances of the hip gluteal musculature can have an important impact on injury prevention within this population.

Objective To describe the muscle performance and imbalances between limbs related to the hip extensor, external rotators and abductors muscles in sprinters.

Design Athletes were assessed during the pre-season period. Muscle function performance of the hip extensors, external rotators and abductors were tested.

Participants All participants (n=69) were federated at the Spanish athletics federation and involved in high-level track and field training sessions (minimum 5 sessions per week) for at least the last 3 years. All of them had undertaken similar training demands in the past.

Interventions We measured the maximum number of repetitions without compensatory movements of each leg during the straight knee bridge exercise (test 1), hip external rotation in prone (test 2) and hip abduction in side-line (test 3). Data analysis and comparison in between limbs were calculated.

Results No significant differences in between limbs were observed in all three tests ($p < 0.005$). The assessment of the hip external rotators (test 2), have shown the worse performance among this population with 30,8% of the athletes on the dominant limb and 35,3% on non-dominant limb not being able to perform even 1 rep without compensation.

Conclusions We did not observe significant imbalances in between limbs. The hip external rotators muscles demonstrated the worst performance with a high number of athletes not being able to perform the test. The weakness of the external rotators and compensations associated with it can have several different impacts on the lower limb function during running, therefore possible importance for injury prevention strategies.

255

ISOKINETIC PROFILE AND CONTRALATERAL DEFICIT OF THE LOWER LIMBS OF ARTISTIC GYMNASTICS ATHLETES

^{1,2}Stefânia Gonçalves Dias, ²Natália Batista Albuquerque Goulart Lemos, ³Morgana Lunardi, ²Fernando de Aguiar Lemos, ⁴Caroline Pieta Dias, ⁴João Carlos Oliva, ⁴Marco Aurélio Vaz. ¹University of Pernambuco, Petrolina, Brazil; ²Federal University of Vale do São Francisco, Petrolina, Brazil; ³Federal University of Santa Catarina, Florianópolis, Brazil; ⁴Federal University of Rio Grande do Sul, Porto Alegre, Brazil

10.1136/bjsports-2021-IOC.234

Background The hamstring/quadriceps (H/Q) functional ratio detects muscle imbalance and monitors knee joint stability, making it a predictor of injury, especially ACL. The injury of this ligament in gymnasts is one of the most incident and of greatest negative impact on recovery and return to sports.

Objective To evaluate functional H/Q at different angular velocities and contralateral deficit of the lower limbs of female artistic gymnastics athletes.

Design Descriptive study.

Setting Federated athletes, participants of competitions at national and international level, in which two were part of the Brazilian Selection of Artistic Gymnastics.

Patients (or Participants) Selected through a club in Porto Alegre, Brazil. They had an average of eight years of experience in the modality, minimum frequency of training of six times a

week, six hours daily. Participants with any lesion that interfered or made it impossible to perform the evaluations were excluded.

Interventions (or Assessment of Risk Factors) Functional H/Q was calculated by dividing the peak of the hamstring eccentric torque by the peak of the quadriceps concentric torque, analyzed using the isokinetic dynamometer (Biodex Medical System).

Main Outcome Measurements Isokinetic protocol at two speeds: 1) 60°/s and 2) 120°/s; and Waterloo questionnaire to determine dominant member (DM).

Results 10 elite gymnasts were evaluated, (mean±SD age 14.9 ±3.7 yrs; body mass 45.6±7.0 kg and height 1.51±0.04 m). The DM, as well as the non-dominant (NDM) at the moment 60°/s (0.71 ± 0.09 ; 0.77 ± 0.12), had values below those established in the literature (functional H/Q ≤ 0.90), at the moment 120°/s were closer to normal (0.85 ± 0.12 ; 0.88 ± 0.13). 5 of the 10 athletes obtained values within the normal range in the MND at 120°/s.

Conclusions All athletes presented significant thigh muscle imbalance, with functional H/Q values lower than those proposed in the literature, with proximity between the limbs. Strength training program of hamstrings should be considered in order to prevent injuries caused by muscle imbalance.

256

THE DIFFERENCES OF STATIC LOWER EXTREMITY ALIGNMENT BETWEEN FEMALE BALLET STUDENTS AND FEMALE NON-BALLET STUDENTS

^{1,2,3}Min Jin Kim, ^{1,2,3}Sae Yong Lee. ¹Department of Physical Education, Yonsei University, Seoul, South Korea (Republic of); ²YISSEM (Yonsei Institute of Sports Science and Exercise Medicine), Seoul, South Korea (Republic of); ³Institute of Convergence Science, Seoul, South Korea (Republic of)

10.1136/bjsports-2021-IOC.235

Background Ballet dancers experience changes in body alignment in repeated practice to achieve the ideal turn-out position, and they have different alignments from that of the non-ballet dancers. However, no quantitative data have been reported to support their differences in lower extremity alignment (LEA).

Objective To compare differences in clinical measures of static LEA between female ballet students and female non-ballet students.

Design Descriptive and cohort study.

Setting Controlled laboratory setting.

Patients (or Participants) Eighty-four female ballet students and fifty-one female non-ballet students were recruited.

Interventions (or Assessment of Risk Factors)

Every participant read and signed the consent form that approved by the Institutional Reviews Board (IRB). The static LEAs were measured using four instruments, Height gage, 6' and 12' Goniometers, PALM inclinometer, and Bubble inclinometer.

Main Outcome Measurements Twelve static LEAs including leg-length, quadriceps angle, tibiofemoral angle, prone rearfoot angle, forefoot angle, tibial torsion, tibial varum, hip anteversion, pelvic tilt, standing quadriceps angle, standing rearfoot angle and navicular drop were measured.

Results Ballet students showed greater mean in tibial torsion ($M_{diff}=4.55$, $SE=.83$, $t_{(266)}=5.51$, $p=.000$, 95% CI: 2.92 – 6.18), anterior-posterior pelvic tilt ($M_{diff}=1.93$, $SE=.46$, $t_{(228)}=4.10$, $p=.000$, 95% CI: 1.00 – 2.86), and standing rearfoot

angle ($M_{diff}=4.81$, $SE=.51$, $t_{(268)}=9.42$, $p=.000$, 95% CI: $3.81 - 5.82$) than non-ballet students. In contrast, ballet students had lesser mean than non-ballet students in tibiofemoral angle tibiofemoral angle ($M_{diff}=-1.05$, $SE=.31$, $t_{(268)}=-3.39$, $p=.001$, 95% CI: $-1.64 - -0.44$), prone rearfoot angle ($M_{diff}=-8.65$, $SE=.56$, $t_{(254.04)}=-16.16$, $p=.000$, 95% CI: $-9.71 - -7.60$), tibial varum ($M_{diff}=-2.52$, $SE=.23$, $t_{(159.90)}=-10.98$, $p=.000$, 95% CI: $-2.96 - -2.06$), hip anteversion ($M_{diff}=-11.47$, $SE=.72$, $t_{(156.15)}=-15.87$, $p=.000$, 95% CI: $-12.90 - -10.04$), and navicular drop ($M_{diff}=-4.45$, $SE=.42$, $t_{(182.22)}=-10.44$, $p=.000$, 95% CI: $-5.29 - -3.61$).

Conclusions Significant results from the alignment indicate that ballet movement and turn-out position may suggest changes in LEA, therefore, clinicians should consider these aspects while preventing and treating dancer injuries.

257 INFLUENCE OF LOWER QUARTER Y-BALANCE TEST™ SCREENING PROTOCOL ON DYNAMIC BALANCE OUTCOMES

¹Scott Lawrance, ¹Megan Jacobs, ¹Emily Boss, ¹Jennifer Popp, ¹Larry Leverenz, ²Michael Weller. ¹Purdue University, West Lafayette, IN, USA; ²Cedarville University, Cedarville, OH, USA

10.1136/bjsports-2021-IOC.236

Background The Lower Quarter Y-Balance Test™ (LQYBT) is commonly used to determine risk of injury or readiness for return to sport. However, clinicians conduct the assessment with differing test protocols potentially altering test outcomes.

Objective To determine if differences in reach distances, composite score, and limb symmetry exist between LQYBT testing protocols.

Design Prospective cohort study.

Setting Controlled laboratory research.

Participants 48 non-injured participants: 32 females, 16 males (21.4 ± 0.3 years, 170.6 ± 9.2 cm, 72.9 ± 14.2 kg).

Interventions Participants completed four testing protocols including barefoot with hands fixed at their hips, barefoot with hands free to move, shod with hands fixed at hips, and shod with hands free to move.

Main Outcome Measures Maximum reach distance was recorded for each limb in the anterior, posteromedial, and posterolateral directions. Limb length composite scores and limb symmetry index (LSI) were calculated for each LQYBT assessment.

Results No statistically significant differences were observed in reach distances, composite scores, or LSI between shod and barefoot protocols. Significant differences were observed in reach distances ($p=0.00$) and in limb length composite scores (right limb $p=0.00$, left limb $p=0.00$) between protocols comparing hands fixed at hips and hand free to move, although no differences were observed in LSI between these conditions when participants were shod ($p=0.27$) or barefoot ($p=0.49$).

Conclusions No differences were observed when participants wore athletic shoes or were barefoot during assessments. Reach distance and limb composite score differences were present when participants were allowed to move their arms and counterbalance their movement during the LQYBT, although no differences in LSI was observed. Results suggest if LSI is used to make clinical decisions, any LQYBT testing protocol can be used. However, testing protocols could influence clinical decisions if reach distances or composite scores are used to make patient care decisions.

258 SPORT PRE-PARTICIPATION HEALTH EVALUATION IN ELITE ATHLETES FROM A MULTISPORT CLUB: PROPOSAL FOR A PERSONALIZED PROTOCOL

¹Ramon Pi, ^{1,2}María Sanz de la Garza, ³Gonzalo Grazioli, ^{1,2}Gil Rodas, ¹Manel García, ^{1,2}Marta Sitges, ^{1,4}Francheck Drobnic. ¹FC Barcelona Medical Department, Barcelona, Spain; ²Hospital Clínic, Barcelona, Spain; ³Sports Medicine Unit, Consell Català de l'Esport, Barcelona, Spain; ⁴Olympic Training Centre, Sant Cugat del Valles, Barcelona, Spain

10.1136/bjsports-2021-IOC.237

Background In 2009 the IOC published a Consensus Statement highlighting the value of periodic health evaluation in elite athletes.

Objective The aim of this study was to evaluate the prevalence of pathological findings, and the preventive value of the method used.

Design Retrospective epidemiological study.

Setting and Participants A total of 2574 elite and professional athletes from Football Club Barcelona (FCB). FCB is a sports club with 13 different sports, 5 professional and 8 amateur.

Interventions (or Assessment of Risk Factors) Between 2008 and 2018, a total of 2574 athletes, with range from 12 to 35 years old, were evaluated through a personalized sport pre-participation health evaluation protocol (SPPHE) at the FCB Medical Department.

Main Outcome Measurements The protocol used was the 'Guía de la Revisión Médica del Fútbol Club Barcelona'. The SPPHE examination consisted of basic medical information, anthropometric data, physical examination, spirometry, basal 12-lead electrocardiography, submaximal cardiovascular exercise testing, and cardiac echocardiography.

Results In 2574 SPPHE, we recorded 750 medical findings (29.1%); including 495 (19.2%) pathological findings, 255 (9.9%) minor abnormal findings, and 958 (37.2%) previous sports injuries. Specific treatment was necessary in 6 cardiovascular diseases. We found 3 endocrinological diseases (type 1e Diabetes Mellitus). Periodic follow-up was specified in all cases, although none prevented the athlete from competing.

Conclusions The SPPHE was effective in identifying a wide range of pathologies (66.3%) in elite athletes from this multisport club. This allow us to provide fast treatment and implement ad hoc strategic preventive programs. We found no alteration that affected the athlete's development or hindered them in reaching the highest level of competition.

259 SHOULD THE SIDE-HOP TEST BE REDUCED FROM 40 CM TO 30 CM WHEN USED IN DEVELOPMENTAL ATHLETES?

^{1,2}Isabel Schneider, ³Nicolai Kammann, ^{1,3}Oliver Haertel. ¹German University of Health and Sport, Berlin, Germany; ²Physio-Motion, Grobenzell, Germany; ³Leichtathletik Foerderzentrum, Munich, Germany

10.1136/bjsports-2021-IOC.238

Background The side-hop test by Gustavsson et al. (2006) helps with return to play decision making after lower extremity injury.

Objective Determine whether a reduced width of 30cm (compared with 40cm) may be more suitable for use in developmental athletes.

Design Cross-sectional study.

Setting School gymnasium.