

Moderate-to-vigorous intensity physical activity from young adulthood to middle age and metabolic disease: a 30-year population-based cohort study

Jason M Nagata ¹, Eric Vittinghoff,² Kelley Pettee Gabriel,³ Andrea K Garber,¹ Andrew E Moran,⁴ Jamal S Rana,^{5,6} Jared P Reis,⁷ Stephen Sidney,⁶ Kirsten Bibbins-Domingo²

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2021-104231>).

¹Division of Adolescent and Young Adult Medicine, Department of Pediatrics, University of California San Francisco, San Francisco, California, USA

²Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco, California, USA

³Department of Epidemiology, The University of Alabama at Birmingham, Birmingham, Alabama, USA

⁴Division of General Medicine, Columbia University, New York, New York, USA

⁵Division of Cardiology, Kaiser Permanente Northern California, Oakland, California, USA

⁶Division of Research, Kaiser Permanente Northern California, Oakland, California, USA

⁷Division of Cardiovascular Sciences, National Heart Lung and Blood Institute, Bethesda, Maryland, USA

Correspondence to

Dr Jason M Nagata, Department of Pediatrics, University of California San Francisco, San Francisco, CA 94143, USA; jason.nagata@ucsf.edu

Accepted 28 August 2021
Published Online First
14 September 2021



© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Nagata JM, Vittinghoff E, Pettee Gabriel K, et al. *Br J Sports Med* 2022;**56**:847–853.

ABSTRACT

Objectives To determine the association between moderate-to-vigorous intensity physical activity (MVPA) trajectories (course over age and time) through the adult life course and onset of metabolic disease (diabetes and dyslipidaemia).

Methods We analysed prospective community-based cohort data of 5115 participants in the Coronary Artery Risk Development in Young Adults study, who were black and white men and women aged 18–30 years at baseline (1985–1986) at four urban sites, collected through 30 years of follow-up. Individualised MVPA trajectories were developed for each participant using linear mixed models.

Results Lower estimated MVPA score at age 18 was associated with a 12% (95% CI 6% to 18%) higher odds of incident diabetes, a 4% (95% CI 1% to 7%) higher odds of incident low high-density lipoprotein (HDL) and a 6% (95% CI 2% to 11%) higher odds of incident high triglycerides. Each additional annual 1-unit reduction in the MVPA score was associated with a 6% (95% CI 4% to 9%) higher annual odds of diabetes incidence and a 4% (95% CI 2% to 6%) higher annual odds of high triglyceride incidence. Analysing various MVPA trajectory groups, participants who were in the most active group at age 18 (over 300 min/week), but with sharp declines in midlife, had higher odds of high low-density lipoprotein and low HDL incidence, compared with those in the most active group at age 18 with subsequent gains.

Conclusion Given recent trends in declining MVPA across the life course and associated metabolic disease risk, young adulthood is an important time period for interventions to increase and begin the maintenance of MVPA.

INTRODUCTION

Metabolic disease, including type 2 diabetes and dyslipidaemia (low high-density lipoprotein cholesterol (HDL-C), high low-density lipoprotein cholesterol (LDL-C) and high triglycerides (TG)) are established risk factors for cardiovascular disease (CVD),^{1–3} the leading cause of mortality in the USA.⁴ Despite our understanding of the general benefits of moderate-to-vigorous intensity physical activity (MVPA) on preventing type 2 diabetes⁵ and dyslipidaemia,^{6–8} there is a paucity of longitudinal data regarding the specific trajectories (the course over age and time) of MVPA during young

adulthood and the association of MVPA trajectory groups with adult-onset CVD. In particular, there is a need to understand how MVPA in young adulthood affects the incidence of metabolic disease.

Young adulthood may set the baseline for later life physical activity trajectories and, thus, be an important time window for intervention.^{9–10} The 2018 US Department of Health and Human Services (HHS) Physical Activity Guidelines recommend a minimum of 150 min of moderate-intensity physical activity per week for adults 18–65 years.^{11–12} The Physical Activity Guidelines Scientific Report noted that young adults have unique growth and developmental needs similar to adolescents, who are recommended to have 60 min/day (420 min/week) of MVPA.¹¹ However, there was insufficient literature on physical activity and health outcomes in young adulthood to confirm current guidance or to support a change to the current approach.¹¹ Thus, the optimal dose and trajectory patterns of MVPA, particularly in young adulthood, to prevent metabolic disease remains unknown.^{13–14}

The objective of this study was to determine the independent associations between the young adult level of MVPA and subsequent changes in MVPA through the transition to midlife and incidence of metabolic disease (diabetes, high LDL-C, low HDL-C, high TG, dyslipidaemia). Second, we examined if specific MVPA trajectory patterns were associated with metabolic disease onset.

METHODS

Study population

The Coronary Artery Risk Development in Young Adults (CARDIA) study is a prospective cohort study that recruited black and white young adults at baseline from 1985 to 1986. Participants (N=5115) were recruited from four urban locations (Birmingham, AL; Chicago, IL; Minneapolis, MN and Oakland, CA) and have been followed up for more than 30 years (years 2, 5, 7, 10, 15, 20, 25, 30 with 90%, 86%, 81%, 77%, 74%, 72%, 72% and 71% retention, respectively). The cohort of participants was designed to be diverse in approximately equal parts by sex, race (black and white), age (18–24 years and 25–30 years at baseline) and educational level (high school or less or higher than high school) within each centre. After conducting the baseline examination, one participant requested to be excluded from all further

analyses. Further details about the study design have been previously published.¹⁵

Measures

Physical activity

Self-reported MVPA was assessed by the interviewer-administered CARDIA Physical Activity History Questionnaire at each of the nine examinations.^{16 17} Participants were asked about the frequency of participation in 13 different activity categories (8 of vigorous and 5 of moderate intensity) within the leisure time and occupational physical activity domains over the prior 12 months. Each activity's intensity was expressed as metabolic equivalents of task (METs), where one MET is defined as the energy used at rest (approximately an oxygen consumption of 3.5 mL/1 kg of body weight/min). Vigorous intensity activities (≥ 6 METs) included running, racquet sports, bicycling faster than 10 miles/hour, swimming, vigorous exercise classes, sports (eg, basketball, football), heavy lifting, carrying or digging on the job and home activities, such as snow shovelling and lifting heavy objects. Moderate intensity activities (3–5 METs) included non-strenuous sports (eg, softball), walking, bowling/golf, home maintenance (eg, gardening, raking) and callisthenics.¹⁸ Each activity was assigned a frequency based on whether it was performed for ≥ 1 hour or during any 1 month in the past year, the number of months it was performed at that level and the number of months it was performed on a frequent basis. Intensity scores (3–8 METs) and duration thresholds (2–5 hours/week) were assigned to each activity; activities above these levels of participation were considered frequent.¹⁷ An MVPA score was computed by multiplying the frequency (number of months) of participation by the intensity (METs) of the activity with a weighting factor for the months of more frequent participation.¹⁹ The MVPA score was the sum of all activities expressed in exercise units (EUs). For reference, an MVPA score of 300 EU estimates the HHS recommendations of approximately 150 min of moderate-intensity activity per week.^{12 20} Given recent evidence that occupational physical activity does not improve health and may be detrimental, constituting an occupational paradox,²¹ we excluded the occupational physical activity question (lifting, carrying or digging on the job) from the MVPA score. Convergent validity of the CARDIA Physical Activity History Questionnaire has been established using report-based measures, including physical activity diaries and detailed quantitative recall questionnaires^{16 19 22} and accelerometers.^{22–24} It has also been indirectly validated by showing expected relations with physical fitness and measures of body fat,^{18 19 25} and has demonstrated adequate test–retest reliability.¹⁹

Diabetes

Blood was drawn and processed at the central laboratory according to standard procedures at each of the nine CARDIA examinations. Glucose was assayed using the hexokinase method. Diabetes was defined as a fasting glucose ≥ 126 mg/dL or on diabetic medications but not pregnant for examinations before May 2011.^{15 17} Diabetes was defined as fasting glucose ≥ 126 mg/dL, 2-hour glucose tolerance test ≥ 200 mg/dL, haemoglobin A1c $\geq 6.5\%$, or being on diabetic medications but not pregnant for examinations after May 2011.

Cholesterol

Fasting lipid measures were measured at each of the nine examinations. Total cholesterol was measured enzymatically and defined as high if levels were ≥ 240 mg/dL.¹⁵ TG were measured

enzymatically and defined as high if levels were ≥ 200 mg/dL.¹⁵ HDL-C was determined after precipitation with dextran sulfate-magnesium chloride and defined as low if levels were < 35 mg/dL for males or < 45 mg/dL for females.²⁶ LDL-C was calculated using the Friedewald equation and defined as high if levels were ≥ 160 mg/dL.²⁷ Dyslipidaemia was defined as TG ≥ 150 mg/dL or HDL < 35 mg/dL for males, or TG ≥ 150 mg/dL or HDL < 45 mg/dL for females.

Covariates

Age (years), race (black or white), sex (male or female), smoking status (never, former or current smoker), alcohol use (mL of alcohol consumed/day), educational attainment (the highest grade of school completed), family history of diabetes or CVD (yes or no), medical history and medications were reported through a questionnaire. The use of diabetes or dyslipidaemia medications was assessed by self-report at each examination. Body mass index (BMI) was calculated based on measured height and weight at each examination.

Statistical analysis

Summarising physical activity

MVPA trajectories were modelled among all CARDIA participants. We developed a linear mixed model (LMM) for repeated measures of MVPA in order to generate succinct summaries of exercise patterns over time. The MVPA slopes use all observations of the MVPA scores prior to metabolic disease onset in order to use as much of the data for each participant as possible and to stabilise the best linear unbiased predictions. The LMM included fixed effects for a four-level categorisation of sex and race, with age as continuous, and their interactions, as well as random effects for participant and age, with unstructured covariance. Our inclusion of these covariates, along with observed outcomes, makes the LMM assumption of missingness at random more plausible, though this assumption is not ultimately verifiable. From the fixed and random effects estimates provided by this model, we calculated the expected MVPA level at age 18 and annual change for each participant. For ease of interpretation, we changed the sign of both summaries, so as to capture the associations of lower level and faster decline in MVPA with increased metabolic risk.

Modelling the association of lower MVPA with incident metabolic disease

Unadjusted cumulative incidence of metabolic diseases (diabetes, high LDL-C, low HDL-C, high TG, or dyslipidaemia) by sex and race/ethnicity were estimated using Kaplan-Meier methods. The data for each participant were then expanded to include a record for each age between study entry and either metabolic disease onset, which was assumed to occur at the first visit at which it was detected, or at censoring by the end of the study of loss to follow-up. Pooled logistic models were used to estimate the independent associations of the expected MVPA at age 18 and subsequent annual change with onset of metabolic disease, adjusting for potential confounders, including sex, race, family history of diabetes or CVD, years of education, smoking status, alcohol use and BMI (smoking status, alcohol use and BMI were time varying, with the last observation carried forward), which have been adjusted for in prior analyses of physical activity and CVD risk (directed acyclic graphs are shown in online supplemental appendix 1).^{17 28} We scaled the estimated MVPA score at age 18 from high to low per 100 EUs, corresponding to 0.45 SDs, for ease of interpretation. We kept annual reduction in total

Table 1 Baseline demographic and health characteristics of participants in the Coronary Artery Risk Development in Young Adults study

N	Total	White women	Black women	White men	Black men	P value
	5114	1307	1480	1170	1157	
Baseline demographic characteristics	Median (IQR)/n (%)	Median (IQR)/n (%)	Median (IQR)/n (%)	Median (IQR)/n (%)	Median (IQR)/n (%)	
Age (years)	25.0 (22.0–28.0)	26.0 (23.0–28.0)	24.0 (21.0–28.0)	26.0 (23.0–28.0)	24.0 (21.0–28.0)	<0.001
Highest grade of school completed	13.0 (12.0–16.0)	15.0 (12.0–16.0)	13.0 (12.0–14.0)	15.0 (12.0–16.0)	12.0 (12.0–14.0)	<0.001
Family history of diabetes	800 (15.6%)	166 (12.7%)	319 (21.6%)	131 (11.2%)	184 (15.9%)	<0.001
Family history of cardiovascular disease	1022 (20.0%)	250 (19.1%)	310 (20.9%)	227 (19.4%)	235 (20.3%)	0.62
Body mass index (BMI)	23.4 (21.2–26.4)	22.0 (20.3–24.6)	24.2 (21.2–28.9)	23.7 (21.9–26.0)	23.7 (21.7–26.4)	<0.001
<18.5 kg/m ²	237 (4.6%)	91 (7.0%)	95 (6.4%)	20 (1.7%)	31 (2.7%)	
18.5–<25 kg/m ²	3091 (60.6%)	921 (70.7%)	728 (49.4%)	741 (63.5%)	701 (60.8%)	
25–30 kg/m ²	1170 (23.0%)	195 (15.0%)	337 (22.8%)	334 (28.6%)	304 (26.4%)	
>30 kg/m ²	599 (11.8%)	95 (7.3%)	315 (21.4%)	72 (6.2%)	117 (10.1%)	
Smoking status						<0.001
Never	2856 (56.2%)	685 (52.7%)	885 (60.1%)	670 (57.8%)	616 (53.8%)	
Former	676 (13.3%)	261 (20.1%)	127 (8.6%)	182 (15.7%)	106 (9.3%)	
Current	1546 (30.4%)	355 (27.3%)	461 (31.3%)	307 (26.5%)	423 (36.9%)	
Alcohol (mL of alcohol consumed per day)	5.4 (0.9–15.5)	4.8 (0.9–12.1)	1.8 (0.0–6.9)	11.1 (3.7–23.2)	10.2 (2.0–25.2)	<0.001
MVPA score at enrolment (EU)	312.0 (168.0–528.0)	306.0 (183.0–502.0)	207.0 (90.0–348.0)	396.0 (228.0–612.0)	408.0 (235.0–656.0)	<0.001
Estimated MVPA score at age 18 (EU)	313.4 (208.1–477.5)	297.7 (204.5–431.6)	209.9 (138.6–303.9)	395.8 (271.6–545.4)	445.2 (305.8–611.9)	<0.001
Estimated MVPA score at age 18 (EU), mean (SD)	361.8 (209.5)	330.6 (168.9)	240.8 (144.6)	428.8 (205.9)	484.2 (229.8)	<0.001
Annual reduction in MVPA score (EU)	2.3 (0.3–4.8)	1.2 (–0.7–3.1)	1.6 (0.3–3.3)	2.2 (0.1–4.5)	5.5 (3.4–8.0)	<0.001
Annual reduction in MVPA score (EU), mean (SD)	2.6 (4.0)	1.2 (3.3)	1.8 (3.2)	2.3 (4.0)	5.8 (4.1)	<0.001
Diabetes	32 (0.6%)	7 (0.5%)	14 (1.0%)	6 (0.5%)	5 (0.4%)	0.29
High LDL cholesterol	1096 (21.7%)	260 (20.0%)	298 (20.4%)	298 (25.9%)	240 (21.0%)	0.001
LDL cholesterol	106.0 (87.0–127.0)	102.0 (85.0–123.0)	107.0 (88.0–129.0)	109.0 (89.0–129.0)	106.0 (85.0–127.0)	<0.001
Low HDL cholesterol	1853 (36.6%)	539 (41.5%)	672 (46.1%)	403 (34.7%)	239 (20.9%)	<0.001
HDL cholesterol	52.0 (44.0–61.0)	55.0 (47.0–64.0)	54.0 (46.0–64.0)	45.0 (40.0–53.0)	52.0 (44.0–61.0)	<0.001
High triglycerides	925 (18.3%)	203 (15.6%)	130 (8.9%)	378 (32.6%)	214 (18.7%)	<0.001
Triglycerides	62.0 (45.0–84.0)	62.0 (45.0–81.5)	56.0 (42.0–75.0)	72.0 (52.0–103.0)	60.0 (45.0–82.0)	<0.001
Dyslipidaemia*	2633 (52.0%)	690 (53.2%)	775 (53.1%)	689 (59.3%)	479 (41.8%)	<0.001

A total MVPA score of 300 EU approximates the Health and Human Services recommendations of 150 min of moderate-intensity activity per week.

*Dyslipidaemia is defined as triglycerides \geq 150 mg/dL or HDL <35 mg/dL for males or HDL <45 mg/dL for females.

EU, exercise units; HDL, high-density lipoprotein; LDL, low-density lipoprotein; MVPA, moderate-to-vigorous intensity physical activity.

MVPA per 1 EU, corresponding to 0.23 SDs. We tested if BMI category or sex and race modified the effect of MVPA (level and change) on incident metabolic disease. Pooled logistic models estimated the associations of meeting various MVPA thresholds at age 18 (<150, 150–300, 300–600, >600 EU) in combination with annual change categories (gain, loss of <2.5 EU/year, of loss >2.5 EU/year) and onset of metabolic disease, adjusting for confounders (sample sizes shown in online supplemental appendix A). We used Stata V.16.0 (StataCorp) for all analyses.

RESULTS

Table 1 shows the baseline demographic and health characteristics of 5114 participants included in the sample. The sample was 51.6% black and 45.5% male. Demographic and health characteristics of the sample at baseline are shown in table 1. Average MVPA declines from young adulthood in all race and sex groups, particularly in black men (figure 1). Incidence of diabetes and cholesterol outcomes by race and sex are presented in online supplemental figures A–E.

Table 2 shows pooled logistic regression model estimates for the associations of the two MVPA summaries (estimated MVPA

level at age 18 and subsequent declines in MVPA) with metabolic disease onset. Model 2.1 adjusted for age only, whereas model 2.2 adjusted for age, race, sex, education, family history, smoking status, alcohol and BMI. In the fully adjusted model (model 2.2), lower estimated MVPA score (per 100 units) at age 18 was associated with a 12% (95% CI 6% to 18%) higher odds of incident diabetes, a 4% (95% CI 1% to 7%) higher odds of incident low HDL, a 6% (95% CI 2% to 11%) higher odds of incident high TG and a 3% (95% CI 0% to 6%) higher odds of incident dyslipidaemia. Each additional annual 1-unit reduction in the MVPA score was associated with a 6% (95% CI 4% to 9%) higher annual odds of diabetes incidence, a 4% (95% CI 2% to 6%) higher annual odds of high triglyceride incidence and a 2% (95% CI 0% to 3%) higher annual odds of dyslipidaemia incidence. Pooled logistic regression model estimates stratified by BMI category (online supplemental appendix B) and race and sex (online supplemental appendix C) are shown in online supplemental appendix 1.

Associations between various MVPA thresholds at age 18 combined with categories of subsequent annual change in MVPA and onset of metabolic disease are shown in table 3. In fully




Figure 1 Average moderate-to-vigorous intensity physical activity (MVPA) trajectories, by race and sex. A total MVPA score of 300 exercise units (EU) approximates the Health and Human Services recommendations of 150 min of moderate-intensity activity.

adjusted models treating MVPA as additive (model 3.1), reductions in MVPA in midlife (compared with gains in MVPA) were associated with onset of all metabolic disease outcomes for any given MVPA threshold level at age 18. In fully adjusted models allowing MVPA categories to interact (model 3.2), certain MVPA combinations are notable. For instance, among participants with over twice the minimum recommended MVPA level (>600 EU) at age 18, those with steep subsequent losses (>2.5 EU/year) had 4.74 higher odds (95% CI 1.55 to 14.50) of low HDL incidence and 3.22 higher odds (95% CI 1.23 to 8.47) of high LDL than those with gains in MVPA. In addition, participants with 300–600 EU at age 18 and subsequent gains in MVPA did not

have higher odds of most metabolic disease onset compared with participants with >600 EU at age 18 and subsequent gains in MVPA.

DISCUSSION

In this prospective cohort study with 30 years of follow-up, we found that a high level of MVPA in young adulthood is a critical starting point for maintaining lifetime metabolic health. Young adult MVPA is associated with lower incidence of diabetes, low HDL-C and high TG, independent of MVPA levels across later adulthood. Maintaining high levels of MVPA in the adult life course is also important; for any given young adult MVPA set point, decline in MVPA through the adult life course is also associated with incident diabetes and high TG.

These findings add to prior literature on physical activity and metabolic disease^{5–8} by leveraging a large longitudinal cohort with 30 years of follow-up data to develop MVPA trajectories throughout the life course. Using these individualised trajectories, we find independent associations with the young adult level of MVPA and subsequent declines in the later adult level of MVPA with diabetes and high TG. It is notable that both estimated MVPA level and slope were independently associated with diabetes and triglyceride onset, even after adjusting for a number of potential confounders.

Our findings indicate that young adult MVPA levels provide protection from subsequent metabolic disease, independent of MVPA levels up through midlife. Thus, young adulthood is an important period for intervention to ensure adequate MVPA levels. Furthermore, our findings indicate that protective levels of activity are higher than the currently recommended minimum. This is an important finding since the HHS Physical Activity Guidelines Scientific Committee Report noted there was insufficient literature on physical activity to inform guidelines in young adults.¹¹ Physical activity typically declines in the transition from

Table 2 Associations between moderate-to-vigorous intensity physical activity (MVPA) trajectories and onset of metabolic disease in the Coronary Artery Risk Development in Young Adults study

	Model 2.1 (adjusted for age)*			Model 2.2 (fully adjusted)†		
	OR	95% CI	P value	OR	95% CI	P value
Diabetes						
Lower MVPA score (per 100 EUs) at age 18	1.26	1.19 to 1.33	<0.001	1.12	1.06 to 1.18	<0.001
Annual reduction in MVPA score (per 1 EU)	1.14	1.12 to 1.17	<0.001	1.06	1.04 to 1.09	<0.001
High LDL cholesterol						
Lower MVPA score (per 100 EUs) at age 18	0.99	0.96 to 1.02	0.65	0.99	0.95 to 1.02	0.49
Annual reduction in MVPA score (per 1 EU)	1.01	1.00 to 1.02	0.09	1.00	0.99 to 1.02	0.58
Low HDL cholesterol						
Lower MVPA score (per 100 EUs) at age 18	1.16	1.13 to 1.19	<0.001	1.04	1.01 to 1.07	0.019
Annual reduction in MVPA score (per 1 EU)	1.00	0.99 to 1.02	0.43	1.01	1.00 to 1.03	0.09
High triglycerides						
Lower MVPA score (per 100 EUs) at age 18	1.02	0.98 to 1.05	0.40	1.06	1.02 to 1.11	0.008
Annual reduction in MVPA score (per 1 EU)	1.05	1.03 to 1.06	<0.001	1.04	1.02 to 1.06	<0.001
Dyslipidaemia‡						
Lower MVPA score (per 100 EUs) at age 18	1.08	1.06 to 1.11	<0.001	1.03	1.00 to 1.06	0.034
Annual reduction in MVPA score (per 1 EU)	1.02	1.01 to 1.03	<0.001	1.02	1.00 to 1.03	0.026

We scaled estimated MVPA score at age 18 from high to low per 100 EUs, corresponding to 0.45 SDs, for ease of interpretation. We kept annual reduction in MVPA per 1 EU, corresponding to 0.23 SDs.

*Model 2.1 includes: estimated MVPA level at age 18, additional annual reduction in MVPA and age. Separate models are presented for each outcome (diabetes, high LDL, low HDL, high triglycerides, dyslipidaemia).

†Model 2.2 includes: estimated MVPA level at age 18, additional annual reduction in MVPA, age, race, sex, education, family history of diabetes or cardiovascular disease, smoking status, alcohol and body mass index. Separate models are presented for each outcome (diabetes, high LDL, low HDL, high triglycerides, dyslipidaemia).

‡Dyslipidaemia is defined as triglycerides ≥ 150 mg/dL or HDL <35 mg/dL for males or HDL <45 mg/dL for females.

EU, exercise unit; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

Table 3 Associations between moderate-to-vigorous intensity physical activity (MVPA) thresholds at age 18, expected annual reduction of MVPA categories, and onset of metabolic disease in the Coronary Artery Risk Development in Young Adults study

	Diabetes			High LDL			Low HDL			High TG			Dyslipidaemia		
	OR	95% CI	P value	OR	95% CI	P value	OR	95% CI	P value	OR	95% CI	P value	OR	95% CI	P value
Model 3.1 treating MVPA as additive, fully adjusted*															
Expected MVPA at age 18 ^b															
>600 EU	Reference			Reference			Reference			Reference			Reference		
300–600 EU	1.12	0.86 to 1.48	0.40	1.11	0.90 to 1.36	0.32	1.35	1.10 to 1.65	0.003	0.97	0.79 to 1.21	0.81	1.23	1.06 to 1.32	0.005
150–300 EU	1.24	0.93 to 1.65	0.15	0.95	0.76 to 1.19	0.64	1.11	0.90 to 1.38	0.33	1.01	0.80 to 1.28	0.93	1.06	0.90 to 1.24	0.49
0–150 EU	1.50	1.03 to 2.17	0.034	0.78	0.57 to 1.05	0.09	0.80	0.61 to 1.06	0.12	1.02	0.72 to 1.44	0.90	0.83	0.65 to 1.05	0.12
Expected annual reduction in MVPA score															
Gain	Reference			Reference			Reference			Reference			Reference		
Reduction 0–2.5 EU/year	1.80	1.38 to 2.35	<0.001	3.49	2.80 to 4.35	<0.001	4.27	3.54 to 5.15	<0.001	1.98	1.58 to 2.47	<0.001	3.30	2.83 to 3.85	<0.001
Reduction >2.5 EU/year	1.74	1.35 to 2.25	<0.001	2.11	1.70 to 2.62	<0.001	1.97	1.63 to 2.38	<0.001	1.50	1.20 to 1.87	<0.001	1.90	1.63 to 2.22	<0.001
Model 3.2 allowing MVPA categories to interact, fully adjusted*															
>600, gain	Reference			Reference			Reference			Reference			Reference		
>600, loss <2.5 EU/year	1.14	0.18 to 7.08	0.89	1.59	0.49 to 5.19	0.44	1.85	0.47 to 7.34	0.38	3.03	0.85 to 10.78	0.09	2.31	0.77 to 6.94	0.13
>600, loss >2.5 EU/year	2.12	0.52 to 8.71	0.30	3.22	1.23 to 8.47	0.018	4.74	1.55 to 14.50	0.006	3.14	0.98 to 10.04	0.054	4.95	1.86 to 13.18	0.001
300–600, gain	0.81	0.18 to 3.56	0.78	0.78	0.28 to 2.20	0.64	0.61	0.18 to 2.08	0.43	1.17	0.35 to 3.92	0.80	1.18	0.42 to 3.26	0.76
300–600, loss <2.5 EU/year	1.83	0.44 to 7.63	0.41	5.11	1.95 to 13.34	<0.001	10.84	5.57 to 32.87	<0.001	3.72	1.16 to 11.93	0.027	8.69	3.27 to 23.12	<0.001
300–600, loss >2.5 EU/year	2.53	0.63 to 10.26	0.19	3.38	1.30 to 8.80	0.012	6.01	1.99 to 18.19	0.001	3.16	0.99 to 10.04	0.051	6.16	2.33 to 16.34	<0.001
150–300, gain	1.48	0.35 to 6.17	0.59	1.46	0.54 to 3.91	0.45	2.69	0.87 to 8.28	0.09	2.26	0.70 to 7.32	0.18	2.81	1.04 to 7.57	0.041
150–300, loss <2.5 EU/year	3.11	0.76 to 12.68	0.11	5.04	1.93 to 13.15	<0.001	10.66	5.52 to 32.25	<0.001	4.48	1.40 to 14.35	0.011	9.6	3.62 to 25.50	<0.001
150–300, loss >2.5 EU/year	2.16	0.53 to 8.84	0.28	1.95	0.74 to 5.17	0.18	2.35	0.76 to 7.23	0.14	2.34	0.72 to 7.57	0.16	2.71	1.01 to 7.29	0.047
<150, gain	2.40	0.56 to 10.25	0.24	2.02	0.73 to 5.62	0.18	3.88	1.23 to 12.21	0.02	3.48	1.03 to 11.74	0.044	4.22	1.55 to 11.53	0.005
<150, loss <2.5 EU/year	3.03	0.74 to 12.49	0.12	3.25	1.22 to 8.67	0.018	5.76	1.87 to 17.71	0.002	3.34	1.01 to 11.04	0.048	5.23	1.94 to 14.14	0.001
<150, loss >2.5 EU/year	1.35	0.22 to 8.22	0.75	0.43	0.05 to 3.76	0.45	1.15	0.1 to 7.20	0.88	1.21	0.19 to 7.71	0.84	1.44	0.29 to 7.05	0.66
A total MVPA score of 300 exercise units (EU) approximates the Health and Human Services recommendations of 150 min of moderate-intensity activity per week.															
* Covariates: age, race, sex, education, family history of cardiovascular disease, smoking status, alcohol and body mass index.															
HDL, high-density lipoprotein; LDL, low-density lipoprotein; TG, triglycerides.															

adolescence to young adulthood due to educational, economic and social transitions.^{9 10 29} For instance, young adults may have fewer opportunities for team or organised sports when they transition to the workforce or college, compared with adolescents, who have physical activity requirements and more opportunities for organised team sports in school.³⁰ The transition to parenthood may also displace leisure time for physical activity.³¹

The MVPA trajectory analysis also identified notable MVPA patterns by race and sex through the life course. For instance, black women have the lowest MVPA levels through the adult life course. Although black men start with high average levels of MVPA in young adulthood, their levels persistently decline throughout the adult life course, similar to findings from the National Health and Nutrition Examination Surveys (NHANES).³² Physical activity interventions and messaging may particularly focus on black women through adulthood and preventing declines in black men. We also found that black adults have higher diabetes incidence throughout the life course compared with white adults, similar to findings in NHANES.^{4 33}

Clinical and public health implications

We find that reductions in MVPA during midlife are associated with an increased incidence of metabolic disease across several outcomes. Public health campaigns and clinicians should focus messaging on maintaining adequate levels of MVPA and preventing declines throughout the adult life course. It is also noteworthy that young adults who were in the most active group at age 18 (over 300 min/week), but had sharp declines in MVPA in midlife, had higher odds of high LDL and low HDL incidence, compared with those in the most active group at age 18 and subsequent gains. Young adults may be more willing to take on immediate risk (eg, not being physically active) if they perceive the outcome is too far into the future, a concept referred to as temporal discounting.³⁴ Young adults may respond to messaging that promotes optimising health beyond just the need to avoid risk.³⁵

Limitations and strengths

Limitations and strengths of this study should be noted. While we adjusted for several potential confounders including age, race, sex, education, family history, smoking status, alcohol and BMI (and behavioural and BMI covariates were time-varying), there is the possibility of unmeasured confounders, such as neighbourhood or genetic factors.³⁶ In addition, mixed models can produce biased effect estimates in the presence of time-varying confounding affected by prior exposure, which would require causal methods to adjust for this potential bias, such as inverse probability weighting.³⁷ There may be attenuation bias due to measurement error in MVPA and residual confounding due to measurement error in some confounders such as smoking and alcohol. There was a possibility of selection bias due to censoring, including losses-to-follow-up and competing risks. MVPA was measured simultaneously with other proposed confounders in one visit, which may lead to over-adjustment bias (adjustment for mediators rather than confounders). MVPA was based on self-report and may be subject to information and prevarication bias, and it did not collect information regarding activity intensity. Nonetheless, the same questionnaire was used across the 30-year follow-up period, which is a distinct strength. Because we scaled the estimated MVPA score at age 18 per 100 EUs for ease of interpretation, the ORs for the two MVPA estimates (level at age 18 and annual reduction) are not standardised and should be interpreted in different scales. The sampling design of CARDIA was not representative of all races or ethnicities in the USA, which may limit generalisability; however, the study specifically focused

What are the findings?

- ⇒ In this prospective longitudinal study with regular 30-year follow-up, we found that low young adult moderate-to-vigorous intensity physical activity (MVPA) was associated with higher odds of diabetes, high triglycerides and low HDL incidence.
- ⇒ Reductions in MVPA during midlife are associated with an increased incidence of metabolic disease.

How might it impact on clinical practice in the future?

- ⇒ Given recent trends in declining MVPA across the life course and associated metabolic disease risk, young adulthood and midlife are important time periods for interventions to increase and maintain MVPA.

on participants identifying as black or white race.¹⁵ Given the larger proportion of the sample with metabolic disease by age 60, the number of eligible participants in the analysis drops with age.

CONCLUSION

In conclusion, MVPA level in young adulthood and declines in later adulthood are each significantly and independently associated with later life metabolic disease onset. Public health and clinical programmes should emphasise, prioritise and develop interventions to promote MVPA in young adulthood, the time when individuals establish an MVPA set point. Future research could examine the mechanisms and mediators by which MVPA may be related to metabolic disease, such as insulin sensitivity. Regardless of young adult MVPA level, interventions to sustain or increase MVPA across adulthood remain another priority for lifetime metabolic health.

Twitter Jason M Nagata @jasonmnagata

Acknowledgements The authors thank Samuel E Benabou for editorial assistance. An abstract corresponding to this manuscript was presented at the Society for Adolescent Health and Medicine Annual Meeting on 10 March 2021.

Contributors JMN conducted the literature search, interpreted findings, and wrote, revised, and edited the manuscript. EV conceptualised the study, analysed data and edited the manuscript. KPG, SS, AKG, KB-D conceptualised the study, interpreted the data and provided critical revisions on the manuscript. AEM, JSR and JPR provided critical revisions on the manuscript. All authors approved the final draft.

Funding JMN is supported by the National Heart, Lung, and Blood Institute (K08HL159350) and the American Heart Association (CDA34760281). KB-D is supported by the National Institutes of Health (K24DK103992). The funders had no role in the study design. The Coronary Artery Risk Development in Young Adults Study (CARDIA) is conducted and supported by the National Heart, Lung, and Blood Institute in collaboration with the University of Alabama at Birmingham (HHSN2682018000051 and HHSN2682018000071), Northwestern University (HHSN2682018000031), the University of Minnesota (HHSN2682018000061) and the Kaiser Foundation Research Institute (HHSN2682018000041). This manuscript has been reviewed by CARDIA for scientific content.

Disclaimer The views expressed in this manuscript are those of the authors and do not necessarily represent the views of the National Heart, Lung, and Blood Institute; the National Institutes of Health or the US Department of Health and Human Services.

Competing interests The authors have no conflicts of interest to report.

Patient and public involvement statement Patients or the public were not involved in the design, conduct, reporting or dissemination plans of our research.

Patient consent for publication Not required.

Ethics approval The institutional review boards at each study site approved all study procedures. All participants provided written informed consent. The University of California, San Francisco institutional review board deemed this secondary data analysis project exempt (IRB 18-26523).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Data are available upon request and data use agreement with the CARDIA Study (<https://www.cardia.dopm.uab.edu/>).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iD

Jason M Nagata <http://orcid.org/0000-0002-6541-0604>

REFERENCES

- Henning RJ. Type-2 diabetes mellitus and cardiovascular disease. *Future Cardiol* 2018;14:491–509.
- Rader DJ, Hovingh GK. HDL and cardiovascular disease. *The Lancet* 2014;384:618–25.
- Nordesgaard BG, Varbo A. Triglycerides and cardiovascular disease. *The Lancet* 2014;384:626–35.
- Virani SS, Alonso A, Benjamin EJ, et al. Heart disease and stroke statistics—2020 update: a report from the American heart association. *Circulation* 2020;141:e139–596.
- Carnethon MR, Sternfeld B, Schreiner PJ, et al. Association of 20-year changes in cardiorespiratory fitness with incident type 2 diabetes: the coronary artery risk development in young adults (cardia) fitness study. *Diabetes Care* 2009;32:1284–8.
- Sarzynski MA, Schuna JM, Carnethon MR, et al. Association of fitness with incident dyslipidemias over 25 years in the coronary artery risk development in young adults study. *Am J Prev Med* 2015;49:745–52.
- Carnethon MR et al. Cardiorespiratory fitness in young adulthood and the development of cardiovascular disease risk factors. *JAMA* 2003;290:3092–100.
- Whitaker KM, Pettee Gabriel K, Buman MP, et al. Associations of accelerometer-measured sedentary time and physical activity with prospectively assessed cardiometabolic risk factors: the cardia study. *J Am Heart Assoc* 2019;8.
- Kwon S, Janz KF, Letuchy EM, et al. Developmental trajectories of physical activity, sports, and television viewing during childhood to young adulthood. *JAMA Pediatr* 2015;169:666–72.
- Li K, Haynie D, Lipsky L, et al. Changes in moderate-to-vigorous physical activity among older adolescents. *Pediatrics* 2016;138:e20161372.
- Physical Activity Guidelines Advisory Committee. 2018 physical activity guidelines Advisory Committee scientific report 2018;2018:Washington DC.
- U.S. Department of Health and Human Services. *Physical activity guidelines for Americans*. 2 edn, 2018.
- Eijssvogels TMH, Thompson PD. Exercise is medicine: at any dose? *JAMA* 2015;314:1915–6.
- Eijssvogels TM, Thompson PD. Are there clinical cardiac complications from too much exercise? *Curr Sports Med Rep* 2017;16:9–11.
- Friedman GD, Cutter GR, Donahue RP, et al. Cardia: study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol* 1988;41:1105–16.
- Pereira MA, FitzerGerald SJ, Gregg EW, et al. A collection of physical activity questionnaires for health-related research. *Med Sci Sports Exerc* 1997;29:1.
- Laddu DR, Rana JS, Murillo R, et al. 25-Year physical activity trajectories and development of subclinical coronary artery disease as measured by coronary artery calcium: the coronary artery risk development in young adults (cardia) study. *Mayo Clinic Proceedings* 2017;92:1660–70.
- Sidney S, Jacobs DR, Haskell WL, et al. Comparison of two methods of assessing physical activity in the coronary artery risk development in young adults (cardia) study. *Am J Epidemiol* 1991;133:1231–45.
- Jacobs DR, Hahn LP, Haskell WL, et al. Validity and reliability of short physical activity history: cardia and the minnesota heart health program. *J Cardiopulm Rehabil* 1989;9:448–59.
- Gabriel KP, Sidney S, Jacobs DR, et al. Convergent validity of a brief self-reported physical activity questionnaire. *Med Sci Sports Exerc* 2014;46:1570–7.
- Shephard RJ. Is there a 'recent occupational paradox' where highly active physically active workers die early? or are there failures in some study methods? *Br J Sports Med* 2019;53:1557–9.
- Jacobs DR, Ainsworth BE, Hartman TJ, et al. A simultaneous evaluation of 10 commonly used physical activity questionnaires. *Med Sci Sports Exerc* 1993;25:81–91.
- Petee Gabriel K, Sidney S, Jacobs DR, et al. Ten-year changes in accelerometer-based physical activity and sedentary time during midlife: cardia study. *Am J Epidemiol* 2018;187:2145–50.
- Sternfeld B, Gabriel KP, Jiang S-F, et al. Risk estimates for diabetes and hypertension with different physical activity methods. *Med Sci Sports Exerc* 2019;51:2498–505.
- Whitaker KM, Pereira MA, Jacobs DR, et al. Sedentary behavior, physical activity, and abdominal adipose tissue deposition. *Med Sci Sports Exerc* 2017;49:450–8.
- Warnick GR. High-Density lipoproteins: the neglected stepchildren whose importance as a risk factor continues to be defined. *Clin Chem* 2008;54:923–4.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499–502.
- Nagata JM, Vittinghoff E, Pettee Gabriel K, et al. Physical activity and hypertension from young adulthood to middle age. *Am J Prev Med* 2021;60:757–65.
- Schmitz KH, Jacobs DR, Leon AS, et al. Physical activity and body weight: associations over ten years in the cardia study. coronary artery risk development in young adults. *Int J Obes Relat Metab Disord* 2000;24:1475–87.
- Kwan MY, Cairney J, Faulkner GE, et al. Physical activity and other health-risk behaviors during the transition into early adulthood. *Am J Prev Med* 2012;42:14–20.
- Bellows-Riecken KH, Rhodes RE. A birth of inactivity? a review of physical activity and parenthood. *Prev Med* 2008;46:99–110.
- Armstrong S, Wong CA, Perrin E, et al. Association of physical activity with income, race/ethnicity, and sex among adolescents and young adults in the United States: findings from the National health and nutrition examination survey, 2007–2016. *JAMA Pediatr* 2018;172:732–80.
- Kwan MY, Thorpe RJ, McGinty EE, et al. Disparities in diabetes: the nexus of race, poverty, and place. *Am J Public Health* 2014;104:2147–55.
- de Water E, Cillessen AHN, Scheres A. Distinct age-related differences in temporal discounting and risk taking in adolescents and young adults. *Child Dev* 2014;103:n/a–97.
- Gooding HC, Milliren C, Shay CM, et al. Achieving cardiovascular health in young adulthood—which adolescent factors matter? *J Adolesc Health* 2016;58:119–21.
- Fleischer NL, Diez Roux AV, Roux AVD. Using directed acyclic graphs to guide analyses of neighbourhood health effects: an introduction. *J Epidemiol Community Health* 2008;62:842–6.
- Mansournia MA, Etmian M, Danaei G, et al. Handling time varying confounding in observational research. *BMJ* 2017;359:j4587.