

Supplementary material 6 - Summary of exercise training interventions and comparison group characteristics

Exercise training intensity was high (n = 3),¹⁻³ moderate (n = 5),⁴⁻⁸ and mixed intensity (n = 9).^{2,9-15} Thirteen studies did not report physical training intensity or reported a target intensity without an objective measure.¹⁶⁻²⁸ One study with two intervention groups was classified with different exercise training intensities (high and mixed intensity).² The frequency of exercise training ranged from two to seven sessions per week, with an average of three times a week. Each session lasted from 15 to 90 minutes, with an average of 43 minutes. The intervention length ranged from nine to 65 weeks, with an average of 18 weeks. The exercise training volume (min/week) ranged from 45 to 270 minutes, with an average of 129 minutes/week.

Exercise training interventions were compared with participants receiving usual care (n = 8);^{1,3-5,11,25,27} cognitive training (n = 3);^{3,17,21} active training (n = 3);^{10,12,14} other activities, such as card-playing, reading, craftwork, social activities, etc. (n = 11);^{2,9,16-18,20-24,28} no treatment (n = 4);^{7,13,19,26} or wait-list (n = 1).⁶ One study offered active training (i.e., light physical activity), reading, playing games, listening to music, and conversations.⁸ Three studies had two exercise training intervention groups^{2,6,9} and four studies had two comparison groups.^{3,17,21,24} We considered each of these studies with more than one intervention or comparison group twice (e.g., Bossers et al. (2015) AT vs. social program and Bossers et al. (2015) MT vs. social program).

References

1. de Oliveira Silva F, Ferreira JV, Plácido J, et al. Three months of multimodal training contributes to mobility and executive function in elderly individuals with mild cognitive impairment, but not in those with Alzheimer's disease: A randomized controlled trial.

- Maturitas*. 2019;126(February):28-33. doi:10.1016/j.maturitas.2019.04.217
2. Enette L, Vogel T, Merle S, et al. Effect of 9 weeks continuous vs. interval aerobic training on plasma BDNF levels, aerobic fitness, cognitive capacity and quality of life among seniors with mild to moderate Alzheimer's disease: A randomized controlled trial. *Eur Rev Aging Phys Act*. 2020;17(1):1-16. doi:10.1186/s11556-019-0234-1
 3. Fonte C, Smania N, Pedrinolla A, et al. Comparison between physical and cognitive treatment in patients with MCI and Alzheimer's disease. *Aging (Albany NY)*. 2019;11(10):3138-3155. doi:10.18632/aging.101970
 4. Hsu CL, Best JR, Davis JC, et al. Aerobic exercise promotes executive functions and impacts functional neural activity among older adults with vascular cognitive impairment. *Br J Sports Med*. 2018;52(3):184-191. doi:10.1136/bjsports-2016-096846
 5. Arcoverde C, Deslandes A, Moraes H, et al. Treinamento na esteira como um tratamento adicional para a doença de Alzheimer: Estudo piloto controlado randomizado. *Arq Neuropsiquiatr*. 2014;72(3):190-196. doi:10.1590/0004-282X20130231
 6. Ho RTH, Fong TCT, Chan WC, et al. Psychophysiological effects of dance movement therapy and physical exercise on older adults with mild dementia: A randomized controlled trial. *Journals Gerontol - Ser B Psychol Sci Soc Sci*. 2020;75(3):560-570. doi:10.1093/geronb/gby145
 7. Kwak YS, Um SY, Son TG, Kim DJ. Effect of regular exercise on senile dementia patients. *Int J Sports Med*. 2008;29(6):471-474. doi:10.1055/s-2007-964853
 8. Telenius EW, Engedal K, Bergland A. Effect of a high-intensity exercise program on physical function and mental health in nursing home residents with dementia: An assessor blinded randomized controlled trial. *PLoS One*. 2015;10(5):1-18. doi:10.1371/journal.pone.0126102
 9. Bossers WJR, Van Der Woude LH V, Boersma F, Hortobágyi T, Scherder EJA, Van Heuvelen MJG. A 9-Week Aerobic and Strength Training Program Improves Cognitive and Motor Function in Patients with Dementia: A Randomized, Controlled Trial. *Am J Geriatr Psychiatry*. 2015;23(11):1106-1116. doi:10.1016/j.jagp.2014.12.191
 10. Karssemeijer EGA, Aaronson JA, Bossers WJR, Donders R, Rikkert MGMO, Kessels RPC. The quest for synergy between physical exercise and cognitive stimulation via exergaming in people with dementia: A randomized controlled trial. *Alzheimers Res Ther*. 2019;11(1):1-13. doi:10.1186/s13195-018-0454-z
 11. Liu-Ambrose T, Best JR, Davis JC, et al. Aerobic exercise and vascular cognitive impairment: a randomized controlled trial. *Neurology*. 2016;87(20):2082-2090. doi:10.1212/WNL.0000000000003332
 12. Sanders LMJ, Hortobágyi T, Karssemeijer EGA, Van Der Zee EA, Scherder EJA, Van Heuvelen MJG. Effects of low- and high-intensity physical exercise on physical and cognitive function in older persons with dementia: A randomized controlled trial. *Alzheimer's Res Ther*. 2020;12(1):1-15. doi:10.1186/s13195-020-00597-3
 13. Kemoun G, Thibaud M, Roumagne N, et al. Effects of a physical training programme on

- cognitive function and walking efficiency in elderly persons with dementia. *Dement Geriatr Cogn Disord*. 2010;29(2):109-114. doi:10.1159/000272435
14. Yu F, Salisbury D, Mathiason MA. Interindividual differences in the responses to aerobic exercise in Alzheimer's disease: Findings from the FIT-AD trial. *J Sport Heal Sci*. 2020;00:1-8. doi:10.1016/j.jshs.2020.05.007
 15. Yu F, Vock DM, Zhang L, et al. Cognitive effects of aerobic exercise in Alzheimer's disease: A pilot randomized controlled trial. *J Alzheimer's Dis*. 2021;80(1):233-244. doi:10.3233/JAD-201100
 16. Cancela JM, Ayán C, Varela S, Seijo M. Effects of a long-term aerobic exercise intervention on institutionalized patients with dementia. *J Sci Med Sport*. 2016;19(4):293-298. doi:10.1016/j.jsams.2015.05.007
 17. Cheng ST, Chow PK, Song YQ, Yu ECS, Lam JHM. Can leisure activities slow dementia progression in nursing home residents? A cluster-randomized controlled trial. *Int Psychogeriatrics*. 2014;26(4):637-643. doi:10.1017/S1041610213002524
 18. Venturelli M, Scarsini R, Schena F. Six-month walking program changes cognitive and ADL performance in patients with Alzheimer. *Am J Alzheimers Dis Other Demen*. 2011;26(5):381-388. doi:10.1177/1533317511418956
 19. Stevens J, Killeen M. A randomised controlled trial testing the impact of exercise on cognitive symptoms and disability of residents with dementia. *Contemporary Nurse*. 2006;21(1):32-39. doi:10.5172/conu.2006.21.1.32
 20. Steinberg M, Sheppard Leoutsakos JM, Podewills LJ, Lyketsos CG. Evaluation of a home-based exercise program in the treatment of Alzheimer's disease: The Maximizing Independence in Dementia (MIND) study. *Int J Geriatr Psychiatry*. 2009;24(7):680-685. doi:10.1002/gps.2175
 21. Cheng ST, Chow PK, Song YQ, et al. Mental and physical activities delay cognitive decline in older persons with dementia. *Am J Geriatr Psychiatry*. 2014;22(1):63-74. doi:10.1016/j.jagp.2013.01.060
 22. de Souto Barreto P, Cesari M, Denormandie P, Armaingaud D, Vellas B, Rolland Y. Exercise or social intervention for nursing home residents with dementia: A pilot randomized, controlled trial. *J Am Geriatr Soc*. 2017;65(9):E123-E129. doi:10.1111/jgs.14947
 23. Harris JB, Johnson CS. The impact of physical versus social activity on the physical and cognitive functioning of seniors with dementia. *Act Adapt Aging*. 2017;41(2):161-174. doi:10.1080/01924788.2017.1306383
 24. Henskens M, Nauta IM, Van Eekeren MCA, Scherder EJA. Effects of physical activity in nursing home residents with dementia: A randomized controlled trial. *Dement Geriatr Cogn Disord*. 2018;46(1-2):60-80. doi:10.1159/000491818
 25. Huang N, Li W, Rong X, et al. Effects of a modified Tai Chi program on older people with mild dementia: A randomized controlled trial. *J Alzheimer's Dis*. 2019;72(3):947-956. doi:10.3233/JAD-190487

26. Miu DKY, Szeto SL, Mak YF. A randomised controlled trial on the effect of exercise on physical, cognitive and affective function in dementia subjects. *Asian J Gerontol Geriatr.* 2008;3(1):8-16. file:///C:/Users/guimo/Box Sync/Lab Drive/Post-doc/Teresa Liu-Ambrose/SystReview_Exercise&Dementia/EligibleStudies/Individual/20.Miu_et_al(2008)_EffectOfExerciseOnPhysical,Cognitive&AffectiveFunctionInDementiaSubjects.pdf
27. Pitkälä KH, Pöysti MM, Laakkonen ML, et al. Effects of the Finnish Alzheimer Disease Exercise Trial (FINALEX): A randomized controlled trial. *JAMA Intern Med.* 2013;173(10):894-901. doi:10.1001/jamainternmed.2013.359
28. Toots A, Littbrand H, Boström G, et al. Effects of exercise on cognitive function in older people with dementia: A randomized controlled trial. *J Alzheimer's Dis.* 2017;60(1):323-332. doi:10.3233/JAD-170014