





# Unilateral tests of lower-limb function as prognostic indicators of future knee-related outcomes following anterior cruciate ligament injury: a systematic review and meta-analysis of 13 150 adolescents and adults

Thomas J West <sup>1,2</sup>, Andrea M Bruder <sup>1,2</sup>, Kay M Crossley <sup>1,2</sup>, Adam G Culvenor <sup>1,2</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2022-105736>).

<sup>1</sup>La Trobe Sport and Exercise Medicine Research Centre, School of Allied Health, Human Services and Sport, La Trobe University, Bundoora, Victoria, Australia

<sup>2</sup>Australian IOC Research Centre, La Trobe University, Bundoora, Victoria, Australia

## Correspondence to

Dr Adam G Culvenor, La Trobe University, Melbourne, VIC 3086, Australia; [a.culvenor@latrobe.edu.au](mailto:a.culvenor@latrobe.edu.au)

Accepted 13 December 2022  
Published Online First  
20 January 2023

## ABSTRACT

**Objective** To investigate the prognostic capacity of individual hop tests, hop test batteries and other unilateral functional performance tests following anterior cruciate ligament (ACL) injury.

**Design** Systematic review with meta-analysis.

**Data sources** Six databases searched up to June 2021.

**Eligibility criteria** Studies reporting associations between unilateral lower-limb function (eg, hop tests) following ACL injury and future ( $\geq 3$  months) knee-related outcomes.

**Results** Of 42 included studies (13 150 participants), all assessed the single-forward hop test and 32 assessed a repeated-forward hop test (crossover hop, triple hop, 6m-timed hop), mostly within a year after ACL injury/reconstruction. Results of meta-analyses indicated that higher single-forward and repeated-forward hop limb symmetry were associated with higher odds of return-to-sport 1–3 years post-ACL reconstruction (OR 2.15; 95% CI 1.30 to 3.54; OR 2.11; 95% CI 1.23 to 3.60, respectively). Higher single-forward and repeated-forward hop limb symmetry was associated with better self-reported symptoms and function 1–37 years after ACL injury (OR 2.51; 95% CI 1.62 to 3.88; OR 4.28; 95% CI 1.65 to 11.08, respectively). Higher limb symmetry on a repeated-forward hop does not appear to be associated with higher odds of successful rehabilitation without ACL reconstruction (OR 1.51; 95% CI 0.94 to 2.44). Achieving  $\geq 90\%$  limb symmetry on the single-forward hop was associated with reduced odds of knee osteoarthritis 5–37 years after ACL injury (OR 0.46; 95% CI 0.23 to 0.94).

**Conclusion** Very low certainty evidence suggests single-forward and repeated-forward hop tests are prognostic indicators for important knee-related outcomes in individuals after ACL injury and may help stratify individuals at risk of poor outcomes to target rehabilitation interventions.

**PROSPERO registration number** CRD42018092197.

## INTRODUCTION

Anterior cruciate ligament (ACL) rupture is a devastating diagnosis for athletes given the lengthy rehabilitation, absence from sports participation and high risk of early-onset osteoarthritis (OA).<sup>1</sup> The primary target of rehabilitation, irrespective of operative or non-operative management, is addressing lower-limb muscle strength, neuromuscular control

## WHAT IS ALREADY KNOWN?

- ⇒ Unilateral measures of lower-limb function are recommended for return to sport clearance after anterior cruciate ligament (ACL) injury.
- ⇒ Hop test results are important indicators of functional recovery after ACL injury.

## WHAT ARE THE NEW FINDINGS?

- ⇒ Very low certainty evidence indicates that a higher Limb Symmetry Index for hop tests (irrespective of specific hop test assessed) is a prognostic indicator for returning to competitive sport but may not infer a reduction of future injury risk.
- ⇒ Very low certainty evidence indicates that higher Limb Symmetry Index on single-forward, repeated-forward hop tests and the one-leg rise test are prognostic for better self-reported symptoms and function 1–37 years after ACL injury.
- ⇒ Very low certainty evidence indicates that a battery of hop tests is associated with reduced odds of knee reinjury though sensitivity analysis revealed conflicting findings with different test batteries.
- ⇒ Very low certainty evidence indicates that achieving  $\geq 90\%$  Limb Symmetry Index on a single-forward hop test between 6 months and 4 years after ACL injury were associated with reduced odds of future knee osteoarthritis.

and functional deficits that can otherwise persist for many years.<sup>2</sup> Functional performance testing is recommended during ACL injury rehabilitation to assess lower-limb function and guide return to sport.<sup>3 4</sup> Functional performance measures also provide valuable feedback to enhance patient motivation and exercise adherence,<sup>5 6</sup> integral to achieving desired outcomes following ACL injury and reconstruction.<sup>7</sup>

Hop tests are common performance measures used during ACL injury and reconstruction rehabilitation<sup>8</sup> and given that they require few resources and are quick to administer, are widely used clinically. Many hop tests have been described (eg, hop for distance (single-forward hop), triple hop, crossover hop), with all having excellent inter-rater reliability.<sup>9–11</sup> Hop tests underpin return to sport



© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** West TJ, Bruder AM, Crossley KM, et al. *Br J Sports Med* 2023;**57**:855–863.

test batteries and meeting acceptable standards (eg,  $\geq 90\%$  Limb Symmetry Index (LSI)) is a goal of rehabilitation.<sup>3</sup>

Some findings suggest that lower-limb functional test scores assessed during and following ACL rehabilitation are associated with future knee-related outcomes.<sup>12–14</sup> For example, poor hop test performance is associated with worse future quality of life and fewer one-leg rises is related to development of radiographic OA.<sup>15–16</sup> However, other studies found no association between functional performance and future outcomes.<sup>17–19</sup> Systematic review evidence is also conflicting. Passing a return to sport test battery (including hop tests) is shown to lower future reinjury risk in some reviews<sup>20–21</sup> but not others,<sup>22</sup> though it is hard to compare findings due to variability in the components of such batteries as well as their criteria for ‘passing’.<sup>23</sup> The prognostic capacity of tests of lower-limb function on future knee-related outcomes has not been synthesised systematically. Such evidence synthesis and critical appraisal may inform clinical management following ACL injury and identify those at risk for worse outcomes to implement preventive strategies. This systematic review aimed to investigate the prognostic capacity of individual hop tests, hop test batteries and other unilateral functional performance tests on knee-related outcomes following ACL injury.

## METHODS

### Search strategy and selection criteria

This systematic review is reported using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement<sup>24</sup> and registered prospectively on PROSPERO (CRD42018092197). We performed an electronic search of six databases (MEDLINE, EMBASE, CINAHL, Scopus, Web of Science, SPORTDiscus) without language restriction from inception to June 2021. The searches combined MeSH terms and keywords related to ACL and hop tests (online supplemental appendix 1). Studies reporting the association (or lack thereof) between unilateral lower-limb functional performance following ACL injury and future outcomes were eligible for inclusion. Specifically, included studies had to: (1) report on individuals after ACL injury treated operatively or non-operatively (no limits to age or type of surgery); (2) include a unilateral functional test (quantitative measure); (3) able to be assessed in a clinical setting (ie, no laboratory or video-recording equipment needed) and (4) report the outcome  $\geq 3$  months following functional testing. We also included studies that assessed a hop test battery (ie, multiple hops combined into one overarching assessment). As it is common for hop test batteries to incorporate a measure of muscle strength,<sup>25–26</sup> we also included studies that evaluated a hop test battery with muscle strength assessment as part of that battery. Cohort studies (prospective and retrospective), case-control studies and randomised controlled trials (RCTs) were eligible for inclusion. Any knee-related outcomes were considered (eg, patient-reported, return to sport, knee reinjury, OA).

Two reviewers (TJW and AGC) independently assessed all titles and abstracts for eligibility and screened reference lists of all relevant articles identified. When eligibility could not be confirmed from title and abstract, full texts were reviewed. Disagreements between reviewers were discussed until consensus, as described previously.<sup>27</sup> During screening of titles, abstracts and full texts, articles published in languages other than English were excluded.

### Data extraction

Two reviewers (TJW and AGC) independently extracted data using a predefined excel spreadsheet. Data extracted included

participant characteristics (eg, age, sex, body mass index), functional test(s) assessed, outcome measures and results of functional tests and knee-related outcome. We contacted study authors for additional information as required.

### Risk of bias and level of evidence

Two reviewers (TJW and AMB) independently assessed risk of bias using a modified version of the Newcastle-Ottawa scale.<sup>28</sup> As we expected most eligible studies to be observational, we a priori developed a modified Newcastle-Ottawa scale for this review, consisting of ten items across four constructs (online supplemental appendix 2). The four constructs were: (1) participant selection; (2) definition of exposure; (3) comparability (exposed and nonexposed groups and (4) outcome assessment. Consistent with Cochrane Handbook for Systematic Reviews recommendations,<sup>29</sup> each item was ranked as high or low risk of bias. Grade of Recommendations Assessments, Development and Evaluation (GRADE) assessed overall certainty of evidence.<sup>30</sup> As part of the GRADE tool, where meta-analysis of any outcome included  $\geq 10$  studies, publication bias secondary to small study effects was assessed using funnel plots and the Eggers test.

### Data synthesis and analysis

Unilateral functional test data was calculated as an LSI defined as ACL injured limb performance  $\div$  contralateral limb performance  $\times 100$ . We analysed LSI primarily as a continuous variable (mean and SD) but also explored the influence of dichotomised LSI results (ie,  $\geq 90\%$  limb symmetry) as this is a common threshold to define success.<sup>25</sup> Knee-related outcome data were dichotomised to define an unsuccessful vs successful outcome using established criteria (table 1). These criteria were those most commonly used in the included studies (online supplemental appendix 3). Where results were not reported using these criteria, authors were contacted for data set out using these thresholds to facilitate meta-analysis.

Based on findings in a recent systematic review,<sup>31</sup> we expected the single-forward hop to be most commonly assessed with sufficient data available for a meta-analysis. Given the similarity of repeated-forward hops (ie, triple hop, crossover hop and/or 6m-timed hop) we grouped these hops into one ‘repeated-forward hop’ category. Crossover hop results were preferred as it comprises forward and lateral movements and is comparable to others for reliability.<sup>9</sup> If data from a crossover hop was not available, we used (in hierarchical order): (1) triple hop, (2) 6m-timed hop. To assess if one repeated-forward hop test was more strongly associated with outcomes than another, we performed sensitivity analysis using data from each individual hop test. Other functional tests (eg, side hop, vertical hop, one-leg rise, step-down test) were analysed separately as they measure different constructs of lower-limb function. We also assessed the relationship between a test battery (ie, including hop tests, and in some studies measures of strength or other biomechanical measures of function) and future knee-related outcomes for studies that combined several functional tests into an overall battery. We planned an a priori secondary analysis to separately assess subgroups (eg, male vs female, non-operative vs operative). Meta-regression was also used to assess the effect of follow-up time and proportion of females on the relationship between single-forward and repeated-forward hop tests and each outcome.

When two or more studies reported the results from the same hop test category (ie, single-forward hop or repeated-forward hop) and association to the same knee-related outcome,

**Table 1** Definitions of dichotomised knee-related outcome

Knee-related outcome	Definition of unsuccessful outcome
Return to sport	<ul style="list-style-type: none"> <li>▶ Not returning to previous level of sport or</li> <li>▶ Not maintaining level of sport (after returning to sport)</li> </ul>
Self-reported symptoms and function	<ul style="list-style-type: none"> <li>▶ The Knee injury and Osteoarthritis Outcome Score 'Englund symptomatic knee criteria' (Englund <i>et al</i>)<sup>32</sup>: QOL subscale <math>\leq 87.5</math> and at least two other subscales below following thresholds: pain <math>\leq 86.1</math>, symptoms <math>\leq 85.7</math>, ADL <math>\leq 86.8</math>, sport/rec <math>\leq 85.0</math>.</li> <li>▶ International Knee Documentation Committee Subjective Knee Evaluation Form 2000: (1) scores below the 15th percentile of uninjured people<sup>33</sup>; (2) less than 'Patient Acceptable Symptom State' (PASS) threshold (ie, <math>&lt;75.9</math>)<sup>34</sup></li> </ul>
Success with ACL deficiency (successful with non-operative treatment)	<ul style="list-style-type: none"> <li>▶ Instability or giving way episodes or</li> <li>▶ Subsequent ACLR</li> </ul>
Subsequent knee injury after ACLR	<ul style="list-style-type: none"> <li>▶ Injury to the ipsilateral or contralateral knee</li> </ul>
Knee osteoarthritis	<ul style="list-style-type: none"> <li>▶ Presence of structural features on imaging which reach an established expert or consensus threshold of magnitude and characteristics to be termed osteoarthritis (eg, Kellgren and Lawrence grade)</li> </ul>

.ACL, anterior cruciate ligament; ACLR, anterior cruciate ligament reconstruction; ADL, activities of daily living; QOL, quality of life.

meta-analysis was performed (Stata, V.17.0) if the OR and 95% CI could be calculated. The OR represents the likelihood of a higher LSI relating to a successful outcome. We used random-effects models as differences in population, functional test administration and outcome definition were expected. When ORs were not reported in the included studies, or could not be calculated from the number of individuals achieving a successful outcome, the standard mean difference (of functional performance between participants with and without a successful outcome) was calculated and then transformed to an OR; this was done using the method described in the Cochrane Handbook.<sup>29</sup> Data from adjusted analyses were extracted wherever possible. An OR  $>1$  indicates a higher LSI being associated with successful future outcomes for return to sport rates, symptoms and function and success with ACL deficiency, while an OR  $<1$  indicates a higher LSI being associated with successful future outcomes for a subsequent knee injury and knee OA (ie, no further knee injury and no knee OA). Unplanned sensitivity analysis was completed to compare outcomes from The Knee injury and Osteoarthritis Outcome Score (KOOS; Englund symptomatic knee criteria)<sup>32</sup> and International Knee Document Committee Subjective Knee Form (IKDC; not scoring a 'PASS' or scores below the 15th percentile of uninjured)<sup>33,34</sup> as they were combined under the single self-reported symptoms and function construct (detailed definitions in table 1). Fourteen studies used a functional test battery and 13 were able to be included in meta-analysis. Where a battery of tests was used, further unplanned sensitivity analysis was performed to compare the batteries consisting of only hop tests to the batteries that consisted of hop tests and strength tests to ascertain how these measures contributed to prognostic value.

Heterogeneity between studies was evaluated for each outcome measure using standard Q-tests, and was calculated as  $I^2$  statistics, describing the percentage of the variability in effect estimates that is due to heterogeneity rather than chance.<sup>29</sup> Data that could not be included in meta-analyses were summarised using qualitative narrative synthesis.

## RESULTS

### Study characteristics

From an initial yield of 10 272 studies, 42 were included in this review consisting of a total 13 150 adolescents and adults (44% female) (figure 1, table 2). Of the 42 studies, 38 (95%) assessed functional performance up to 1 year after ACL reconstruction (ACLR) (hop tests conducted at 19 months in one study,<sup>35</sup> 3 years in another<sup>15</sup> and 4 years post-ACL injury in two studies<sup>36,37</sup> with follow-up assessment of outcomes ranging between 1 and

37 years following injury (see online supplemental appendix 3 for study outcome definitions). All but two studies<sup>38,39</sup> reported functional test results as an LSI, of which one<sup>38</sup> could not be estimated and was reported separately. All studies included the single-forward hop and 32 studies (78%) included a repeated forward hop (crossover hop;  $n=22$ , triple hop;  $n=17$ ; 6m-timed hop;  $n=15$ ) (online supplemental appendix 4). It was not feasible to analyse subgroups due to insufficient data for stratification.

### Risk of bias and certainty of evidence

Our risk of bias assessment had excellent interrater reliability between the two independent assessors ( $k=97\%$ ). Only four studies conducted power analysis, 10 had acceptable loss to follow-up (mean lost to follow-up for all included studies was 19%) and 16 adjusted for potential confounders (online supplemental appendix 5). The overall certainty of evidence for all estimates was rated as very low using the GRADE tool. Evidence was downgraded based on study limitations, study design (observational studies), risk of bias, inconsistency, indirectness and publication bias (online supplemental appendices 6, 7).


### Synthesis of results for limb symmetry as a continuous score

#### Return to sport

Meta-analysis of nine studies (1020 participants) revealed that higher single-forward and repeated-forward hop test scores reported as an LSI up to 1 year following ACLR were associated with higher return to sport rates 1.5–3 years post-ACLR (OR 2.15; 95% CI 1.30 to 3.54;  $I^2$  70.6% and OR 2.11; 95% CI 1.23 to 3.60;  $I^2$  72.7%, respectively) (figures 2 and 3). One study not included in meta-analysis reported no association between single-forward hop test scores 3 years post-ACL injury or reconstruction, and level of sports participation at 5 years.<sup>15</sup>

#### Self-reported symptoms and function

Meta-analysis of 10 studies (3107 participants) indicated that higher LSI scores on the single-forward and repeated-forward hop tests up to 4 years post-ACL injury were associated with better self-reported knee symptoms and function 1–37 years post-ACL injury (OR 2.51; 95% CI 1.62 to 3.88;  $I^2$  69.2% and OR 4.28; 95% CI 1.65 to 11.08;  $I^2$  69.2%, respectively) (figures 2 and 3). Sensitivity analysis separating studies reporting data from the KOOS (Englund symptomatic knee criteria)<sup>32</sup> and IKDC (not scoring a 'PASS' or scores below the 15th percentile of uninjured),<sup>33,34</sup> indicated that higher LSI scores for the single-forward hop were associated with better KOOS (OR 1.44;



**Figure 1** PRISMA flow diagram of study selection. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

95% CI 1.21 to 1.71;  $I^2$  4.0%) and IKDC scores independently (OR 6.47; 95% CI 3.45 to 12.15;  $I^2$  0.0%) (online supplemental appendix 8). Higher LSI scores on the one-leg rise test within 1 year post-ACLR were also associated with better self-reported symptoms and function at 3–5 years post-ACLR (OR 3.04; 95% CI 1.56 to 5.92;  $I^2$  0.0%) (online supplemental appendix 9).

#### Success with ACL deficiency

Meta-analysis of seven studies (419 participants) showed no association between single-forward hop LSI up to 2 years post-ACL injury and success for individuals with an ACL deficient knee 1–8 years later (OR 1.32; 95% CI 0.75 to 2.32;  $I^2$  54.1%) (figure 2). Higher scores on a repeated-forward hop test were not associated with increased odds of successful outcome (ie, no ACLR) 1–8 years after ACL injury (OR 1.51; 95% CI 0.94 to 2.44;  $I^2$  27.4%) (figure 3). One study's results reporting on the vertical hop, were not included in meta-analyses and found that higher scores were prognostic for success with ACL deficiency.<sup>18</sup>

#### Subsequent knee injury after ACLR

Meta-analysis of seven studies (7549 participants) found that a higher LSI on single- or repeated-forward hop tests at 6–12 months post-ACLR was not associated with reduced odds of subsequent knee injury to either knee 2–5 years later (OR 1.24; 95% CI 0.90 to 1.71;  $I^2$  47.2% and OR 1.03; 95% CI 0.74 to 1.44;  $I^2$  0.0%, respectively) (figures 2 and 3).

#### Knee osteoarthritis

Meta-analysis of four studies (444 participants) revealed no association between single-forward and repeated-forward hop tests and developing knee OA between 1 and 37 years after ACL injury (OR 0.60; 95% CI 0.25 to 1.43;  $I^2$  75.8% and OR 0.39; 95% CI 0.13 to 1.20;  $I^2$  76.7%, respectively) (figures 2 and 3). Sensitivity analysis indicated that higher scores on the crossover hop (but not other repeated-forward hop tests) may be prognostic for structural knee OA (OR 0.14 95% CI 0.05 to 0.34) (online supplemental appendix 10).

#### Other outcomes

Single studies reported the association between functional tests and outcomes of knee biomechanics and self-efficacy. Higher scores on a single-leg step-down test at 3 months post-ACLR significantly correlated with greater knee flexion excursion ( $r=0.65$ ,  $p=0.001$ ) and knee extensor moment ( $r=0.54$ ,  $p=0.001$ ) during running 6 months post-ACLR.<sup>38</sup> LSI for the single-forward hop, within the first year after ACL injury, was moderately correlated with worse knee-related self-efficacy after 1 year of rehabilitation alone or ACLR ( $r=0.31$ ,  $p=0.001$ ).<sup>40</sup>

#### Synthesis of results for dichotomised limb symmetry scores

Meta-analyses of studies using a dichotomous threshold of  $\geq 90\%$  limb symmetry demonstrated consistent findings compared with using continuous LSI measures for outcomes of



Table 2 Characteristics of included studies grouped by outcome

Study	Study design	Participants*†	Female‡ (%)	Time of functional test§¶	Age±SD, years**	BMI±SD, kg/m <sup>2</sup> **	Functional tests	Mean follow-up¶¶
<b>Return to sport</b>								
Ardern et al, 2015 <sup>65</sup> †† Australia	Prospective	ACLR: 94	38%	1 year	29±8	NR	Single hop Triple hop	2 years
Ebert and Annear, 2019 <sup>66</sup> †††† Australia	Prospective	ACLR: 50	36%	1 year	26±10	24.8±4.0	Single hop Crossover hop 6m-timed hop Triple hop	2 years
Faleide et al, 2021 <sup>67</sup> Norway	Prospective	ACLR: 103	47%	9 months	29±10	NR	Test battery only§§	2 years
Kitaguchi et al, 2020 <sup>68</sup> Japan	Prospective	ACLR: 124	59%	6 months	17±2¶¶¶	NR	Single hop	1 year
Moksnes and Risberg, 2009 <sup>69</sup> *** Norway	Prospective	ACLD: 52 ACLR: 50	45%	6 months	28±9	NR	Single hop Crossover hop Triple hop 6m-timed hop	1.5 years
Nawasreh et al, 2017 <sup>73</sup> †† USA	Prospective	ACLR: 95	34%	6 months	27±10	24.9±3.6	Single hop Crossover hop Triple hop 6m-timed hop Test battery	1–2 years
Toole et al, 2017 USA <sup>70</sup>	Prospective	ACLR: 115	77%	8 months	17±2	NR	Single hop Crossover hop Triple hop 6m-timed hop Test battery	1.5 years
Webster et al, 2019 Australia <sup>71</sup> ††	Prospective	ACLR: 222	40%	1 year	26±9	NR	Single hop Crossover hop	3 years
Welling et al, 2020 <sup>72</sup> †††† Netherlands	Prospective	ACLR: 64	31%	10 months	27±8	NR	Single hop Triple hop Side hop Test battery	2 years
<b>Symptoms and function</b>								
Cristiani et al, 2020 <sup>73</sup> †† Sweden	Prospective	ACLR: 2335	49%	6 months	29±10	NR	Single hop	2 years
Culvenor et al, 2016 <sup>74</sup> †† Australia	Prospective	ACLR: 93	40%	1 year	27 (13) (median and IQR)	NR	Single hop Triple hop One-leg rise	3 years
Ericsson et al, 2013 <sup>18</sup> †††† Denmark	Prospective	ACLD: 42 Delayed ACLR: 20 Early ACLR: 45	26%	37 weeks (median)	26±5	23.9±2.6	Single hop Square hop Vertical hop One-leg rise Test battery	5 years
Filbay et al, 2021 <sup>36</sup> †† Sweden	Prospective	ACLR or ACLD: 109	28%	4 years	24±6†††	27 (24–29) (median and IQR)¶¶¶	Single hop	32–37 years
Logerstedt et al, 2012 <sup>12</sup> †† USA	Prospective	ACLR: 85	45%	6 months	26 (15–54) (median and range)	NR	Single hop Crossover hop Triple hop 6m-timed hop	1 year
Månsson et al, 2013 <sup>75</sup> †† Sweden	Prospective	ACLR: 73	30%	Pre-ACLR	24 (14–40) (median and range)	NR	Single hop	2 years
McGrath et al, 2017 <sup>76</sup> †† Australia	Prospective	ACLR: 64	31%	24 weeks	28	25	Single hop 6m-timed hop Test battery	2 years
Øiestad et al, 2012 <sup>77</sup> †††† Norway	Prospective	ACLR: 181	42%	1 year	27±9	NR	Single hop Triple hop	15 years
Stropnik et al, 2020 <sup>78</sup> Slovenia	Prospective	ACLR: 60	31%	Pre-ACLR	32±11§§§	25.7±3.5§§§	Single hop	6 months
<b>Success with ACL deficiency</b>								
Button et al, 2006 <sup>39</sup> †† Wales	Prospective	ACLD: 42¶¶¶¶	40%	5 months	28±7	NR	Single hop	2–3 years
Eitzen et al, 2010 <sup>79</sup> Norway/USA	Prospective	ACLD: 145	52%	60 days	26 (14–47) (median and range)	NR	Single hop Crossover hop Triple hop 6m-timed hop	15 months
Ekås et al, 2019 <sup>80</sup> Norway	Prospective	ACLD: 44	34%	2 years	11±1	24.7 (range: 16.6–40.8)	Single hop Crossover hop Triple hop 6 m-timed hop	8 years
Fitzgerald et al, 2000 <sup>81</sup> USA	Prospective	ACLD: 28	25%****	4 weeks (median)	29±11		Single hop Crossover hop Triple hop 6m-timed hop	8 months
Grindem et al, 2018 <sup>82</sup> Norway/USA	Prospective	ACLD 118	51%	2 months	28±10	24.2±3.8	Single hop Cross-over hop Triple hop 6m-timed hop	2 years
<b>Subsequent injury after ACLR</b>								
Beischer et al, 2020 <sup>83</sup> Sweden	Prospective	ACLR: 159	44%	9 months	20±3	NR	Test battery only§§	2 years
Cristiani et al, 2021 <sup>84</sup> Sweden	Prospective	ACLR: 6510	44%	6 months	28±10	24.2±3.4	Single hop	2 years
Fältström et al, 2021 <sup>35</sup> Sweden	Prospective	ACLR: 117	100%	19 months	20±2	NR	Single hop Side hop Test battery	2 years

Continued

Table 2 Continued


Study	Study design	Participants*†	Female‡ (%)	Time of functional tests§¶	Age±SD, years**	BMI±SD, kg/m <sup>2</sup> **	Functional tests	Mean follow-up¶
Grindem et al, 2016 <sup>25</sup> †† Norway/USA	Prospective	ACL: 106	51%	6 months††††	25±7	NR	Single hop Crossover hop Triple hop 6m-timed hop Test battery	2 years
King et al, 2021 <sup>85</sup> †††† Ireland	Prospective	ACL: 115	0%	9 months	21±4	NR	Single hop Test battery	2 years
Kyritsis et al, 2016 <sup>26</sup> ††††† Qatar	Prospective	ACL: 158	0%	247 days	21±4	NR	Single hop Crossover hop Triple hop Test battery	2 years
Nawasreh et al, 2016 <sup>86</sup> USA	Prospective	ACL: 95	32%	6 months	27±10	25±3.8	Test battery only§§	2 years
Sousa et al, 2017 <sup>42</sup> USA	Retrospective	ACL: 223	59%	6 months	26±11	25.8±4.5	Test battery only§§	2 years
van Melick et al, 2021 <sup>87</sup> Netherlands	Prospective	ACL: 144	31%	1 year	24±7	NR	Single hop Vertical hop Side hop Test battery	2 years
Webster et al, 2019 <sup>88</sup> Australia	Prospective	ACL: 329	39%	1 year	17±2	NR	Single hop Crossover hop	5 years
Wellsandt et al, 2017 <sup>51</sup> USA	Prospective	ACL: 70	33%	6 months	26±10	24.9±3.8	Single hop Crossover hop Triple hop 6m-timed hop Test battery	2 years
Knee osteoarthritis								
Filbay et al, 2021 <sup>37</sup> †† Sweden	Prospective	ACL or ACLD: 127	28%	4 years	23 (19–28) (median and IQR)†††	27 (24–29) (median and IQR)¶¶	Single hop	32–37 years
Janssen et al, 2013 <sup>89</sup> †† Netherlands	Prospective	ACL: 86	34%	Pre-ACL	31±8§§§	24.5±3.1§§§	Single hop	10 years
Patterson et al, 2018 <sup>16</sup> †† Australia	Prospective	ACL: 78	38%	1 year	28±14	25.7±4.2	Single hop Crossover hop Side hop One-leg rise	5 years
Pinczewski et al, 2007 <sup>90</sup> †† Australia	Prospective	ACL: 149	47%****	1 year	25 (13–52) (median and range)****	NR	Single hop	10 years
Wellsandt et al, 2018 <sup>14</sup> †† USA	Prospective	ACL or ACLD: 58	35%****	6 months	28±11	NR	Single hop 6m-timed hop	5 years
Other (studies not included in meta-analysis)								
Flosadottir et al, 2016 <sup>15</sup> Sweden	Prospective	ACL or ACLD: 54	28%	3 years	29±5	24.6±3.4	Single hop Square hop Vertical hop	5 years
Flosadottir et al, 2018 <sup>40</sup> §§§§ Sweden (Knee self-efficacy)	Prospective	ACLD: 20 ACL: 69	28%	37 weeks (median)	25±5	23.8±2.8	Single hop Square hop Vertical hop	5 years
Kline et al, 2016 <sup>38</sup> ¶¶¶¶ USA (Knee biomechanics)	Prospective	ACL: 30	53%	3 months	21±7	NR	Single leg step down	6 months

\*As listed in the original paper (may differ with extra data provided by the author).  
†Studies were of mixed sporting populations apart from those indicated otherwise.  
‡Most studies reported on percentage of male vs female sex though used female/women and men/male terminology interchangeably.  
§Mean, unless indicated otherwise.  
¶Test or follow-up was number of years from the time of injury (deficient cohorts) or ACLR (reconstructed cohorts) unless indicated otherwise.  
\*\* Assessed at time of functional assessment unless indicated otherwise.  
††Extra data provided by authors to enable synthesis of results and dichotomous outcome comparison.  
‡‡Hop test results also compared to self-reported symptoms and function.  
§§Only results from test battery able to be included for analysis.  
¶¶Assessed at time of follow-up.  
\*\*\*Hop test results also compared to success with ACL deficiency.  
†††Assessed at time of injury.  
‡‡‡Hop test results also compared with knee osteoarthritis.  
§§§Assessed at time of primary ACLR.  
¶¶¶¶Only 32 participants included in analysis of hopping data because 10 were unable to hop on ACLD limb.  
\*\*\*\* Estimated from data describing total cohort.  
††††Functional testing was taken at 6 and 12 months post-ACL injury.  
‡‡‡‡Studies included professional male athletes (all other studies were of a general population).  
§§§§Outcome used was knee self-efficacy.  
¶¶¶¶Outcome used was biomechanical variables.  
ACLD, anterior cruciate ligament deficient; ACLR, anterior cruciate ligament reconstruction; BMI, body mass index; NR, not reported.

return to sport (online supplemental appendices 11, 12). Effect sizes for self-reported symptoms and function and success with ACL deficiency outcomes were attenuated where dichotomous LSI data were used (online supplemental appendices 11, 12). Meta-analysis of two studies also found no association with side hop 2 years post-ACLR and subsequent knee injury (online supplemental appendix 13). The notable exception was knee OA, with meta-analysis of six studies (679 participants) demonstrating that achieving LSI  $\geq 90\%$  on the single-forward hop up to 4 years post-ACL injury was associated with reduced odds of radiographic knee OA at 5–37 years after ACL injury/reconstruction (LSI  $\geq 90\%$ : OR 0.46; 95% CI 0.23 to 0.94;  $I^2$  54.5%) (online supplemental appendix 11).

### Synthesis of results for a test battery

Passing a test battery (ie, achieving LSI  $\geq 90\%$  on all tests) was associated with greater odds of returning to sport (OR 3.33; 95% CI 1.05 to 10.53;  $I^2$  55.3%) (online supplemental appendix 14). Passing a battery of tests was not associated with reduced odds of subsequent knee injury after ACLR (OR 0.62; 95% CI 0.26 to 1.48;  $I^2$  67.3%) (online supplemental appendix 14). Sensitivity analysis revealed that passing test batteries made up exclusively of hop tests was associated with avoiding subsequent knee injury (OR 0.29, 95% CI 0.10 to 0.85;  $I^2$  44.3%). Conversely, passing test batteries that included strength measures along with hops tests was not associated with avoiding



**Figure 2** Forest plot displaying the association between single-forward hop results (continuous Limb Symmetry Index scores) and future outcomes.


subsequent knee injury (1.31; 95% CI 0.63 to 2.71  $I^2$  22.4%) (online supplemental appendix 15).

### Sensitivity analysis of associations found between hop test and knee-related outcomes

Sensitivity analysis separating each repeated-forward hop test demonstrated that triple hop, crossover hop and 6m-timed hop had similar effect sizes for each outcome, with no significant between-test heterogeneity observed: RTS ( $I^2$  71.8%,  $p=0.645$ ), self-reported symptoms and function ( $I^2$  51.9%,  $p=0.978$ ), success with ACL deficiency ( $I^2$  19.6%,  $p=0.843$ ) and subsequent injury after ACLR ( $I^2$  1.2%,  $p=0.175$ ). Between test heterogeneity was significantly different with individual repeated-forward hop tests and knee OA, with the crossover hop demonstrating a significantly larger association with knee OA than the 6m-timed hop or triple hop test ( $I^2$  76.7%,  $p=0.014$ ) (online supplemental appendix 10). Meta-regression indicated that follow-up time and proportion of females did not influence the relationship between single-forward and repeated-forward hop tests and all outcomes (online supplemental appendix 16).

## DISCUSSION

This systematic review of 42 studies and 13 150 adolescents and adults following ACL injury, found an increased odds of poor knee-related outcomes after 1–37 follow-up years in individuals not achieving adequate LSI on unilateral functional tests. These findings highlight the need to optimise knee function in the first 1–2 years post-ACL injury and reconstruction to reduce the long-term burden. Caution is warranted when interpreting these findings given the very low certainty of evidence associated with the results. The low certainty of evidence points to the need for high quality longitudinal data that is well controlled for confounders existing (and/or changing) during the long duration between exposure (hop tests) and outcomes (eg, physical activity, muscle strength).



**Figure 3** Forest plot displaying the association between repeated-forward hop results (continuous Limb Symmetry Index scores) and future outcomes. In brackets are the specific repeated-forward hop used in analysis.

### What is the prognostic value of hop tests for return to sport, reinjury and self-reported outcomes?

Achieving a higher LSI on the singleforward and repeated-forward hop tests was associated with twice the odds of future return to sport (OR 2.15; 95% CI 1.30 to 3.54;  $I^2$  70.6% and OR 2.11; 95% CI 1.23 to 3.60;  $I^2$  72.7%, respectively) (figures 2 and 3) with 9/10 studies in this analysis using pre-injury level of sport as the measure of successful return. While this indicates the potential importance of achieving adequate functional performance to facilitate a return to preinjury sport, those who return to high-impact and pivoting sports are known to be at higher risk of reinjury.<sup>25</sup> Our pooled results suggest a slightly elevated risk of reinjury in those with a higher hop score, which may reflect higher return to sport rates in those with better hop performance. There is much debate regarding the importance of achieving common return to sport criteria (ie, LSI  $\geq 90\%$  on a functional test battery including hop tests) to reduce reinjury risk.<sup>20–41</sup> For example, some original studies and systematic reviews report that meeting functional criteria increases reinjury risk,<sup>20–42</sup> while others report a reduced risk of reinjury on passing return to sport criteria.<sup>21–25</sup> Inconsistencies are likely explained by whether sport participation (both level and type of sport) exposure was accounted for and whether some participants who may initially fail test batteries continued rehabilitation until passing.<sup>41–43</sup> Individuals following ACL injury and reconstruction should be educated about the risk of reinjury when returning to sport regardless of adequate functional performance and consider other factors known to increase re-injury risk (eg, time postsurgery, younger age, symmetry of quadriceps strength, fear).<sup>25–44–45</sup>

We also observed that higher LSI on single-forward and repeated-forward hop tests was associated with more than twofold and fourfold higher odds of better self-reported symptoms and function, respectively (OR 2.51; 95% CI 1.62 to

3.88;  $I^2$  69.2% and OR 4.28; 95% CI 1.65 to 11.08;  $I^2$  69.2%) (figures 2 and 3). These findings suggest that hop tests provide important prognostic value for longer-term symptomatic and functional outcomes and may provide targets to reduce the burden of symptoms following ACL injury. Original studies did not account for baseline symptoms and function that may have altered results. Self-reported symptoms and function at follow-up may reflect higher scores at baseline rather than improvement over time, although other studies have shown only fair cross-sectional correlation between functional tests and self-reported symptoms and function.<sup>46</sup> Nevertheless, our results demonstrate the prognostic value of these functional tests, which may aid clinicians in identifying those at higher risk of longer-term burden.

### Can hop tests tell us who is going to get post-traumatic structural OA?

Knee OA is a burdensome sequelae of traumatic knee injury with its risk not reduced with reconstruction.<sup>47</sup> We found achieving LSI  $\geq 90\%$  on the single-forward hop was associated with reduced odds of developing structural knee OA (LSI  $\geq 90\%$ : OR 0.46; 95% CI 0.23 to 0.94;  $I^2$  54.5%) (online supplemental appendix 11), though this was attenuated when analysing hop test scores as continuous variables. The stronger association with OA when using a dichotomous hop score, may be due to including different studies in each meta-analysis ( $n=2$  not included in both meta-analyses). In sensitivity analysis data from a single study<sup>16</sup> indicated that higher scores on the cross-over hop may be prognostic for structural knee OA (OR 0.14; 95% CI 0.05 to 0.34) (online supplemental appendix 10). Structural knee OA was defined differently across included studies (eg, joint space narrowing, Kellgren and Lawrence classification and adding symptoms to definition) and there was no observable trend for the impact of hop test performance and structural OA based on length of follow-up. In the context of OA prevention, addressing impairments in hop tests (ie, quadriceps weakness) may provide a target for intervention though RCTs are required to confirm this.<sup>48 49</sup>

### Is one test better than another or is a battery of tests needed?

Generally, results were similar between single- and repeated-forward hop tests for most outcomes. As studies often included multiple tests, we created a hierarchy for reporting repeated-forward hop tests (ie, crossover hop, triple hop, 6m-timed hop). Despite tests measuring somewhat different constructs (ie, crossover hop includes lateral and forward movement and 6m-timed hop measuring speed of completion) each individual repeated-forward hop tests found similar associations (online supplemental appendix 10). Current clinical trial evidence suggests that trialling rehabilitation without surgery is not inferior to having early ACLR.<sup>50</sup> In this context, our results highlight the importance of conducting hop tests during the initial rehabilitation period to monitor function, which might help to determine an individual's ability to cope with an ACL deficient knee, alongside other factors such as desired level of activity (eg, return to cutting sports).

A battery of tests is recommended to determine readiness to return to sport, although it is difficult to compare batteries given the inconsistency in tests included across different studies.<sup>23</sup> The common definition for 'passing' a test battery is LSI  $\geq 90\%$  on all tests included.<sup>25 26 51</sup> Results from our review suggest that both single-forward and repeated-forward hops are similar to a test

battery as prognostic indicators of return to sport. Passing a test battery (consisting only of hop tests) was associated with reduced odds of further knee injury more than threefold compared with not passing (OR 0.29; 95% CI 0.10 to 0.85) (online supplemental appendix 15). In contrast, we found that a higher LSI on individual hop tests (figures 2 and 3) or passing test batteries (including those with hop tests and strength measures) was not associated with subsequent knee injury (online supplemental appendix 14). There is no agreement across our analyses reflective of findings from various recent systematic reviews.<sup>20-22 46</sup> Compared with these reviews, we included different studies and we also pooled knee injuries under one outcome (eg, ipsilateral and contralateral reinjury, other knee injury). Given our results and those from previous reviews, we suggest that a test battery is likely needed to identify those at higher risk of future knee injury. Unfortunately for clinicians, exactly how many and what particular combination of tests to use remains unclear but current guidelines recommend using test batteries that include hop tests and measures of strength.<sup>3 52</sup>

### The prognostic capacity of less common functional tests

Less common tests of functional performance included in this review were the one-leg rise, side hop and vertical hop. We found that performance on the one-leg rise was associated with future self-reported symptoms and function. The one-leg rise test may be a surrogate of quadriceps (and lower-limb) strength<sup>53</sup> and is becoming more popular in clinical settings due to the lack of equipment required compared with isokinetic dynamometry. Higher side hop performance was not associated with subsequent knee injury after ACLR (online supplemental appendix 13). Higher LSI on the vertical hop was associated with success with an ACL deficient knee<sup>18</sup> and may be a better alternative to forward hop tests to assess knee function because the knee contributes more to vertical height than horizontal distance (which is more a function of hip/ankle power).<sup>54-56</sup>

In our review, we only compared the quantitative achievement of a functional test and results do not account for movement quality or other aspects of how tests were completed. Sixty-five per cent of included studies were deemed high risk of bias in terms of how the hop tests were assessed or completed, identifying the need for greater consistency to strengthen comparisons. It is also becoming more apparent that poor movement quality (eg, measured using two-dimensional or three-dimensional kinematics) is another important construct to assess given its association to poor outcome despite restoration of quantitative function (eg, LSI  $\geq 90\%$ ).<sup>57</sup> The importance of qualitative assessment also highlights the limitation of the LSI as a measure of recovered function. The contralateral (reference) limb may deteriorate after ACL injury with disuse and central inhibition<sup>58 59</sup> and thus not be an accurate preinjury reference standard. For this reason, other criteria, such as an estimated preinjury capacity (ie, contralateral hop test result immediately after ACL injury) have been proposed as a more suitable measure of preinjury function as well as normative values from well-matched controls.<sup>51 58</sup>

### Limitations

Results from our review are limited by the very low certainty of evidence. The level of evidence (GRADE assessment) required downgrading due to observational studies and large  $I^2$  values. These values indicated heterogeneity among pooled studies that was largely unexplained as our sensitivity analyses were unable to determine a source of this. Most studies had unacceptable lost to follow-up (eg,  $>15\%$ ), which may indicate attrition bias



skewing results. While eight of the 42 included studies accounted for baseline differences (eg, age, sex, body mass index), the majority did not account for these factors and their potential impact on results. Indirectness was also found with time between test and follow-up differing across studies and though this enhances generalisability, it also reduces confidence in the estimated effects. Graft type used for ACLR was also not able to be included as part of our analysis to compare surgery types as the data were not available. Finally, we were unable to provide an estimate of the relationship between functional test results and future psychological recovery (eg, confidence, fear) as no studies had available data evaluating this. However, we recognise the growing awareness of the importance of restoring confidence and reducing fear after ACL injury<sup>60–64</sup> and this should be investigated further.

## CONCLUSION

In this systematic review, we found unilateral tests of lower-limb function can be prognostic for future knee-related outcomes in individuals after ACL injury, though with very low certainty evidence. Early in rehabilitation, hop testing may be used as part of criteria to consider non-operative management and along with other tests may provide insight into potential future symptoms and function over the long term, including the development of post-traumatic knee OA. Hop tests are also recommended to be used as part of return to sport testing, though achieving adequate performance may not reduce subsequent knee injuries.

**Twitter** Thomas J West @tsewmot, Andrea M Bruder @AndreaBruder, Kay M Crossley @kaymrossley and Adam G Culvenor @agculvenor

**Contributors** TJW, AGC, AMB and KMC designed the study. TJW and AGC completed all searches, study selection (including inclusion and exclusion of abstracts). TJW and AGC completed all data extraction. TJW and AMB completed all risk of bias assessment. TJW and AGC and KMC planned the analyses, TJW completed the meta-analyses and all authors interpreted the data. TJW wrote the initial draft and all authors critically revised the manuscript for important intellectual content approved the final version of the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

## ORCID iDs

Thomas J West <http://orcid.org/0000-0002-6297-1094>  
 Andrea M Bruder <http://orcid.org/0000-0001-5422-5756>  
 Kay M Crossley <http://orcid.org/0000-0001-5892-129X>  
 Adam G Culvenor <http://orcid.org/0000-0001-9491-0264>

## REFERENCES

- Culvenor AG, Collins NJ, Guermazi A, *et al.* Early knee osteoarthritis is evident one year following anterior cruciate ligament reconstruction: a magnetic resonance imaging evaluation. *Arthritis Rheumatol* 2015;67:946–55.
- Keays SL, Bullock-Saxton JE, Keays AC, *et al.* A 6-year follow-up of the effect of graft site on strength, stability, range of motion, function, and joint degeneration after anterior cruciate ligament reconstruction: Patellar tendon versus semitendinosus and gracilis tendon graft. *Am J Sports Med* 2007;35:729–39.
- van Melick N, van Cingel REH, Brooijmans F, *et al.* Evidence-based clinical practice update: practice guidelines for anterior cruciate ligament rehabilitation

based on a systematic review and multidisciplinary consensus. *Br J Sports Med* 2016;50:1506–15.

- Diermeier TA, Rothrauff BB, Engebretsen L, *et al.* Treatment after ACL injury: Panther symposium ACL treatment consensus group. *Br J Sports Med* 2021;55:14.
- Rambaud AJM, Ardern CL, Thoreux P, *et al.* Criteria for return to running after anterior cruciate ligament reconstruction: a scoping review. *Br J Sports Med* 2018;52:1437–44.
- Rambaud AJM, Rossi J, Neri T, *et al.* Evolution of functional recovery using hop test assessment after ACL reconstruction. *Int J Sports Med* 2020;41:696–704.
- Grindem H, Risberg MA, Eitzen I. Two factors that may underpin outstanding outcomes after ACL rehabilitation. *Br J Sports Med* 2015;49:1425.
- Davies WT, Myer GD, Read PJ. Is it time we better understood the tests we are using for return to sport decision making following ACL reconstruction? A critical review of the hop tests. *Sports Med* 2020;50:485–95.
- Reid A, Birmingham TB, Stratford PW, *et al.* Hop testing provides a reliable and valid outcome measure during rehabilitation after anterior cruciate ligament reconstruction. *Phys Ther* 2007;87:337–49.
- Gustavsson A, Neeter C, Thomeé P, *et al.* A test battery for evaluating hop performance in patients with an ACL injury and patients who have undergone ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2006;14:778–88.
- Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. *Am J Sports Med* 1991;19:513–8.
- Logerstedt D, Grindem H, Lynch A, *et al.* Single-legged hop tests as predictors of self-reported knee function after anterior cruciate ligament reconstruction: the Delaware-Oslo ACL cohort study. *Am J Sports Med* 2012;40:2348–56.
- Nawasreh Z, Logerstedt D, Cummer K, *et al.* Functional performance 6 months after ACL reconstruction can predict return to participation in the same preinjury activity level 12 and 24 months after surgery. *Br J Sports Med* 2018;52:375.
- Wellsandt E, Axe MJ, Snyder-Mackler L. Poor performance on single-legged hop tests associated with development of posttraumatic knee osteoarthritis after anterior cruciate ligament injury. *Orthop J Sports Med* 2018;6:2325967118810775.
- Flosadottir V, Roos EM, Ageberg E. Muscle function is associated with future patient-reported outcomes in young adults with ACL injury. *BMJ Open Sport Exerc Med* 2016;2:e000154.
- Patterson B, Culvenor AG, Barton CJ, *et al.* Poor functional performance 1 year after ACL reconstruction increases the risk of early osteoarthritis progression. *Br J Sports Med* 2020;54:546–55.
- Ithurburn MP, Longfellow MA, Thomas S, *et al.* Knee function, strength, and resumption of Preinjury sports participation in young athletes following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2019;49:145–53.
- Ericsson YB, Roos EM, Frobell RB. Lower extremity performance following ACL rehabilitation in the KANON-trial: impact of reconstruction and predictive value at 2 and 5 years. *Br J Sports Med* 2013;47:980–5.
- Wellsandt E, Gardinier ES, Manal K, *et al.* Decreased knee joint loading associated with early knee osteoarthritis after anterior cruciate ligament injury. *Am J Sports Med* 2016;44:143–51.
- Webster KE, Hewett TE. What is the evidence for and validity of Return-to-Sport testing after anterior cruciate ligament reconstruction surgery? A systematic review and meta-analysis. *Sports Med* 2019;49:917–29.
- Ashigbi EYK, Banzer W, Niederer D. Return to sport tests' prognostic value for reinjury risk after anterior cruciate ligament reconstruction: a systematic review. *Med Sci Sports Exerc* 2020;52:1263–71.
- Losciale JM, Zdeb RM, Ledbetter L, *et al.* The association between passing Return-to-Sport criteria and second anterior cruciate ligament injury risk: a systematic review with meta-analysis. *J Orthop Sports Phys Ther* 2019;49:43–54.
- Roe C, Jacobs C, Hoch J, *et al.* Test batteries after primary anterior cruciate ligament reconstruction: a systematic review. *Sports Health* 2022;14:205–15.
- Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- Grindem H, Snyder-Mackler L, Moksnes H, *et al.* Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. *Br J Sports Med* 2016;50:804–8.
- Kyritsis P, Bahr R, Landreau P, *et al.* Likelihood of ACL graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture. *Br J Sports Med* 2016;50:946.
- Culvenor AG, Øiestad BE, Hart HF, *et al.* Prevalence of knee osteoarthritis features on magnetic resonance imaging in asymptomatic uninjured adults: a systematic review and meta-analysis. *Br J Sports Med* 2019;53:1268–78.
- Wells G, Shea B, O'Connell D, *et al.* The Newcastle–Ottawa Scale (NOS) for Assessing the Quality of Non-Randomized Studies in Meta-Analysis The Ottawa Hospital. [Internet]. 2000. The Ottawa Hospital. Available: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp) [Accessed 2022].
- Akl E, Altman D, Aluko P, *et al.* *Cochrane Handbook for systematic reviews of interventions*, 2019.
- Atkins D, Best D, Briss PA, *et al.* Grading quality of evidence and strength of recommendations. *BMJ* 2004;328:1490.

- 31 Hegedus EJ, McDonough S, Bleakley C, *et al.* Clinician-friendly lower extremity physical performance measures in athletes: a systematic review of measurement properties and correlation with injury, part 1. The tests for knee function including the hop tests. *Br J Sports Med* 2015;49:642.
- 32 Englund M, Roos EM, Lohmander LS. Impact of type of meniscal tear on radiographic and symptomatic knee osteoarthritis: a sixteen-year followup of meniscectomy with matched controls. *Arthritis Rheum* 2003;48:2178–87.
- 33 Anderson AF, Irrgang JJ, Kocher MS, *et al.* The International knee documentation Committee subjective knee evaluation form: normative data. *Am J Sports Med* 2006;34:128–35.
- 34 Muller B, Yabroudi MA, Lynch A, *et al.* Defining thresholds for the patient acceptable symptom state for the IKDC subjective knee form and KOOS for patients who underwent ACL reconstruction. *Am J Sports Med* 2016;44:2820–6.
- 35 Fältström A, Häggglund M, Hedevik H, *et al.* Poor validity of functional performance tests to predict knee injury in female soccer players with or without anterior cruciate ligament reconstruction. *Am J Sports Med* 2021;49:1441–50.
- 36 Filbay S, Andersson C, Gauffin H, *et al.* Prognostic factors for patient-reported outcomes at 32 to 37 years after surgical or nonsurgical management of anterior cruciate ligament injury. *Orthop J Sports Med* 2021;9:23259671211021592.
- 37 Filbay S, Gauffin H, Andersson C, *et al.* Prognostic factors for tibiofemoral and patellofemoral osteoarthritis 32–37 years after anterior cruciate ligament injury managed with early surgical repair or rehabilitation alone. *Osteoarthr Cartil* 2021.
- 38 Kline PW, Johnson DL, Ireland ML, *et al.* Clinical predictors of knee mechanics at return to sport after ACL reconstruction. *Med Sci Sports Exerc* 2016;48:790–5.
- 39 Button K, van Deursen R, Price P. Classification of functional recovery of anterior cruciate ligament copers, non-copers, and adapters. *Br J Sports Med* 2006;40:853–9.
- 40 Flosadottir V, Frobell R, Roos EM, *et al.* Impact of treatment strategy and physical performance on future knee-related self-efficacy in individuals with ACL injury. *BMC Musculoskelet Disord* 2018;19:50.
- 41 Capin JJ, Snyder-Mackler L, Risberg MA, *et al.* Keep calm and carry on testing: a substantive reanalysis and critique of 'what is the evidence for and validity of return-to-sport testing after anterior cruciate ligament reconstruction surgery? A systematic review and meta-analysis'. *Br J Sports Med* 2019;53:1444.
- 42 Sousa PL, Krych AJ, Cates RA, *et al.* Return to sport: does excellent 6-month strength and function following ACL reconstruction predict midterm outcomes? *Knee Surg Sports Traumatol Arthrosc* 2017;25:1356–63.
- 43 Webster KE, Hewett TE. Return-to-sport testing following ACL reconstruction revisited. *Br J Sports Med* 2020;54:2–3.
- 44 Paterno MV, Huang B, Thomas S, *et al.* Clinical factors that predict a second ACL injury after ACL reconstruction and return to sport: preliminary development of a clinical decision algorithm. *Orthop J Sports Med* 2017;5:2325967117745279.
- 45 Paterno MV, Rauh MJ, Schmitt LC, *et al.* Incidence of second ACL injuries 2 years after primary ACL reconstruction and return to sport. *Am J Sports Med* 2014;42:1567–73.
- 46 Losciale JM, Bullock G, Cromwell C, *et al.* Hop testing lacks strong association with key outcome variables after primary anterior cruciate ligament reconstruction: a systematic review. *Am J Sports Med* 2020;48:511–22.
- 47 Meuffels DE, Favejee MM, Vissers MM, *et al.* Ten year follow-up study comparing conservative versus operative treatment of anterior cruciate ligament ruptures. A matched-pair analysis of high level athletes. *Br J Sports Med* 2009;43:347–51.
- 48 Culvenor AG, West TJ, Bruder AM, *et al.* Supervised exercise-therapy and patient education rehabilitation (super) versus minimal intervention for young adults at risk of knee osteoarthritis after ACL reconstruction: SUPER-Knee randomised controlled trial protocol 2022:(under review).
- 49 Culvenor AG, Girdwood MA, Juhl CB, *et al.* Rehabilitation after anterior cruciate ligament and meniscal injuries: a best-evidence synthesis of systematic reviews for the OPTIKNEE consensus. *Br J Sports Med* 2022;56:1445–53.
- 50 Frobell RB, Roos EM, Roos HP, *et al.* A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med* 2010;363:331–42.
- 51 Wellsandt E, Failla MJ, Snyder-Mackler L. Limb symmetry indexes can overestimate knee function after anterior cruciate ligament injury. *J Orthop Sports Phys Ther* 2017;47:334–8.
- 52 Meredith SJ, Rauer T, Chmielewski TL, *et al.* Return to sport after anterior cruciate ligament injury: Panther symposium ACL injury return to sport consensus group. *Orthop J Sports Med* 2020;8:2325967120930829–29.
- 53 Woon E-L, Low J, Sng Y-L, *et al.* Feasibility, correlates, and validity of the one-leg sit-to-stand test in individuals following anterior cruciate ligament reconstruction. *Phys Ther Sport* 2021;52:280–6.
- 54 Kotsifaki A, Korakakis V, Graham-Smith P, *et al.* Vertical and horizontal hop performance: contributions of the hip, knee, and ankle. *Sports Health* 2021;13:128–35.
- 55 Kotsifaki A, Van Rossom S, Whiteley R, *et al.* Single leg vertical jump performance identifies knee function deficits at return to sport after ACL reconstruction in male athletes. *Br J Sports Med* 2022;56:490.
- 56 Kotsifaki A, Korakakis V, Whiteley R, *et al.* Measuring only hop distance during single leg hop testing is insufficient to detect deficits in knee function after ACL reconstruction: a systematic review and meta-analysis. *Br J Sports Med* 2020;54:139–53.
- 57 King E, Richter C, Daniels KAJ, *et al.* Biomechanical but not strength or performance measures differentiate male athletes who experience ACL reinjury on return to level 1 sports. *Am J Sports Med* 2021;49:918–27.
- 58 Patterson BE, Crossley KM, Perraton LG, *et al.* Limb symmetry index on a functional test battery improves between one and five years after anterior cruciate ligament reconstruction, primarily due to worsening contralateral limb function. *Phys Ther Sport* 2020;44:67–74.
- 59 Luc-Harkey BA, Harkey MS, Pamukoff DN, *et al.* Greater intracortical inhibition associates with lower quadriceps voluntary activation in individuals with ACL reconstruction. *Exp Brain Res* 2017;235:1129–37.
- 60 George SZ, Lentz TA, Zeppieri G, *et al.* Analysis of shortened versions of the Tampa scale for kinesophobia and pain catastrophizing scale for patients after anterior cruciate ligament reconstruction. *Clin J Pain* 2012;28:73–80.
- 61 Kvist J, Ek A, Sporrstedt K, *et al.* Fear of re-injury: a hindrance for returning to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2005;13:393–7.
- 62 Paterno MV, Flynn K, Thomas S, *et al.* Self-reported fear predicts functional performance and second ACL injury after ACL reconstruction and return to sport: a pilot study. *Sports Health* 2018;10:228–33.
- 63 Ardern CL, Taylor NF, Feller JA, *et al.* A systematic review of the psychological factors associated with returning to sport following injury. *Br J Sports Med* 2013;47:1120–6.
- 64 Hart HF, Culvenor AG, Guermazi A, *et al.* Worse knee confidence, fear of movement, psychological readiness to return-to-sport and pain are associated with worse function after ACL reconstruction. *Phys Ther Sport* 2020;41:1–8.
- 65 Ardern CL, Taylor NF, Feller JA, *et al.* Sports participation 2 years after anterior cruciate ligament reconstruction in athletes who had not returned to sport at 1 year: a prospective follow-up of physical function and psychological factors in 122 athletes. *Am J Sports Med* 2015;43:848–56.
- 66 Ebert JR, Annear PT. ACL reconstruction using autologous Hamstrings augmented with the ligament augmentation and reconstruction system provides good clinical scores, high levels of satisfaction and return to sport, and a low Retear rate at 2 years. *Orthop J Sports Med* 2019;7:2325967119879079–79.
- 67 Faleide AGH, Magnussen LH, Strand T, *et al.* The role of psychological readiness in return to sport assessment after anterior cruciate ligament reconstruction. *Am J Sports Med* 2021;49:1236–43.
- 68 Kitaguchi T, Tanaka Y, Takeshita S, *et al.* Importance of functional performance and psychological readiness for return to preinjury level of sports 1 year after ACL reconstruction in competitive athletes. *Knee Surg Sports Traumatol Arthrosc* 2020;28:2203–12.
- 69 Moksnes H, Risberg MA. Performance-based functional evaluation of non-operative and operative treatment after anterior cruciate ligament injury. *Scand J Med Sci Sports* 2009;19:345–55.
- 70 Toole AR, Ithurburn MP, Rauh MJ, *et al.* Young athletes cleared for sports participation after anterior cruciate ligament reconstruction: how many actually meet recommended Return-to-Sport criterion cutoffs? *J Orthop Sports Phys Ther* 2017;47:825–33.
- 71 Webster KE, McPherson AL, Hewett TE, *et al.* Factors associated with a return to Preinjury level of sport performance after anterior cruciate ligament reconstruction surgery. *Am J Sports Med* 2019;47:2557–62.
- 72 Welling W, Benjaminse A, Lemmink K, *et al.* Passing return to sports tests after ACL reconstruction is associated with greater likelihood for return to sport but fail to identify second injury risk. *Knee* 2020;27:949–57.
- 73 Cristiani R, Mikkelsen C, Edman G, *et al.* Age, gender, quadriceps strength and hop test performance are the most important factors affecting the achievement of a patient-acceptable symptom state after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2020;28:369–80.
- 74 Culvenor AG, Collins NJ, Guermazi A, *et al.* Early Patellofemoral osteoarthritis features one year after anterior cruciate ligament reconstruction: symptoms and quality of life at three years. *Arthritis Care Res* 2016;68:784–92.
- 75 Månsson O, Kartus J, Sernert N. Pre-operative factors predicting good outcome in terms of health-related quality of life after ACL reconstruction. *Scand J Med Sci Sports* 2013;23:15–22.
- 76 McGrath TM, Waddington G, Scarvell JM, *et al.* An ecological study of anterior cruciate ligament reconstruction, part 2: functional performance tests correlate with Return-to-Sport outcomes. *Orthop J Sports Med* 2017;5:2325967116688443.
- 77 Øiestad BE, Holm I, Engebretsen L, *et al.* The prevalence of patellofemoral osteoarthritis 12 years after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2013;21:942–9.
- 78 Stropnik D, Sajovic M, Kacin A, *et al.* Early clinical and neuromuscular properties in patients with normal or sub-normal subjective knee function after anterior cruciate ligament reconstruction. *Arch Orthop Trauma Surg* 2020;140:1231–9.
- 79 Eitzen I, Moksnes H, Snyder-Mackler L, *et al.* Functional tests should be accentuated more in the decision for ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2010;18:1517–25.
- 80 Ekås GR, Moksnes H, Grindem H, *et al.* Coping with anterior cruciate ligament injury from childhood to maturation: a prospective case series of 44 patients with mean 8 years' follow-up. *Am J Sports Med* 2019;47:22–30.

- 81 Fitzgerald GK, Axe MJ, Snyder-Mackler L. A decision-making scheme for returning patients to high-level activity with nonoperative treatment after anterior cruciate ligament rupture. *Knee Surg Sports Traumatol Arthrosc* 2000;8:76–82.
- 82 Grindem H, Wellsandt E, Failla M, *et al.* Anterior cruciate ligament injury—who succeeds without reconstructive surgery? The Delaware-Oslo ACL cohort study. *Orthop J Sports Med* 2018;6:2325967118774255.
- 83 Beischer S, Gustavsson L, Senorski EH, *et al.* Young athletes who return to sport before 9 months after anterior cruciate ligament reconstruction have a rate of new injury 7 times that of those who delay return. *J Orthop Sports Phys Ther* 2020;50:83–90.
- 84 Cristiani R, Forssblad M, Edman G, *et al.* Age, time from injury to surgery and quadriceps strength affect the risk of revision surgery after primary ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2021;29:4154–4162.
- 85 King E, Richter C, Daniels KAJ, *et al.* Can biomechanical testing after anterior cruciate ligament reconstruction identify athletes at risk for subsequent ACL injury to the contralateral uninjured limb? *Am J Sports Med* 2021;49:609–19.
- 86 Nawasreh Z, Logerstedt D, Cummer K, *et al.* Do patients failing Return-to-Activity criteria at 6 months after anterior cruciate ligament reconstruction continue demonstrating deficits at 2 years? *Am J Sports Med* 2017;45:1037–48.
- 87 van Melick N, Pronk Y, Nijhuis-van der Sanden M, *et al.* Meeting movement quantity or quality return to sport criteria is associated with reduced second ACL injury rate. *J Orthop Res* 2022;40:117–28.
- 88 Webster KE, Feller JA. Clinical tests can be used to screen for second anterior cruciate ligament injury in younger patients who return to sport. *Orthop J Sports Med* 2019;7:2325967119863003.
- 89 Janssen RPA, du Mée AWF, van Valkenburg J, *et al.* Anterior cruciate ligament reconstruction with 4-strand hamstring autograft and accelerated rehabilitation: a 10-year prospective study on clinical results, knee osteoarthritis and its predictors. *Knee Surg Sports Traumatol Arthrosc* 2013;21:1977–88.
- 90 Pinczewski LA, Lyman J, Salmon LJ, *et al.* A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and Patellar tendon autograft: a controlled, prospective trial. *Am J Sports Med* 2007;35:564–74.

## Appendix 1.

## Systematic search strategy

<b>MEDLINE</b>	Anterior Cruciate Ligament [MeSH] OR Anterior Cruciate Ligament Injuries [Mesh] OR Anterior Cruciate Ligament Reconstruction [Mesh] OR "anterior cruciate ligament" [mp] ACL [mp] <b>AND</b> Leg adj4 function* [mp] OR leg adj4 performance [mp] OR "lower extremity" adj4 function* [mp] OR "lower extremity" adj4 performance [mp] OR "lower-extremity" adj4 function* [mp] OR "lower-extremity" adj4 performance [mp] OR "lower limb" adj4 function* [mp] OR "lower limb" adj4 performance [mp] OR "lower-limb" adj4 function* [mp] OR "lower-limb" adj4 performance [mp] OR hop* [mp] OR squat* [mp] OR "one-leg rise" [mp] OR "one leg rise" [mp] OR "single leg rise" [mp] OR jump* [mp] OR "knee bend" [mp] OR battery [mp] OR stair* [mp]
<b>Embase</b>	Anterior Cruciate Ligament [MeSH] OR Anterior Cruciate Ligament Injury [Mesh] OR Anterior Cruciate Ligament Reconstruction [MeSH] OR Anterior Cruciate Ligament Rupture [MeSH] OR "anterior cruciate ligament" [mp] ACL [mp] <b>AND</b> Leg adj4 function* [mp] OR leg adj4 performance [mp] OR "lower extremity" adj4 function* [mp] OR "lower extremity" adj4 performance [mp] OR "lower-extremity" adj4 function* [mp] OR "lower-extremity" adj4 performance [mp] OR "lower limb" adj4 function* [mp] OR "lower limb" adj4 performance [mp] OR "lower-limb" adj4 function* [mp] OR "lower-limb" adj4 performance [mp] OR hop* [mp] OR squat* [mp] OR "one-leg rise" [mp] OR "one leg rise" [mp] OR "single leg rise" [mp] OR jump* [mp] OR "knee bend" [mp] OR battery [mp] OR stair* [mp]
<b>CINAHL</b>	Anterior Cruciate Ligament Injuries [MeSH] OR Anterior Cruciate Ligament Reconstruction [MeSH] OR ACL [tiab] OR "anterior cruciate ligament" [tiab] <b>AND</b> Leg n4 function* [tiab] OR leg n4 performance [tiab] OR "lower extremity" n4 function* [tiab] OR "lower extremity" n4 performance [tiab] OR "lower-extremity" n4 function* [tiab] OR "lower-extremity" n4 performance [tiab] OR "lower limb" n4 function* [tiab] OR "lower limb" n4 performance [tiab] OR "lower-limb" n4 function* [tiab] OR "lower-limb" n4 performance [tiab] OR hop* [tiab] OR squat* [tiab] OR "one-leg rise" [tiab] OR "one leg rise" [tiab] OR "single leg rise" [tiab] OR jump* [tiab] OR "knee bend*" [tiab] OR battery [tiab] OR stair* [tiab]
<b>Scopus</b>	ACL [tiab] OR "anterior cruciate ligament" [tiab] <b>AND</b>
<b>Web of Science</b>	Leg W/4 function* [tiab] OR leg W/4 performance [tiab] OR "lower extremity" W/4 function* [tiab] OR "lower extremity" W/4 performance [tiab] OR "lower-extremity" W/4 function* [tiab] OR "lower-extremity" W/4 performance [tiab] OR "lower limb" W/4 function* [tiab] OR "lower limb" W/4 performance [tiab] OR "lower-limb" W/4 function* [tiab] OR "lower-limb" W/4 performance [tiab] OR hop* [tiab] OR squat* [tiab] OR "one-leg rise" [tiab] OR "one leg rise" [tiab] OR "single leg rise" [tiab] OR jump* [tiab] OR "knee bend*" [tiab] OR battery [tiab] OR stair* [tiab]
<b>SPORTDiscus</b>	



## Appendix 2. Risk of Bias Tool

Modified Newcastle-Ottawa Scale from Wells, G. et al, The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp) 2013;

Selection	
<p>1. Definition of ACL injured population</p> <p>Low risk of bias = a High risk of bias = b</p>	<p>a) Clearly described if the inclusion/exclusion criteria of an ACL injured person stated <b>both</b> of the following criteria:</p> <ul style="list-style-type: none"> <li>i) Diagnosed ACL injury with clinical/imaging or surgical confirmation (e.g. Lachman's or pivot shift test ± MRI/arthroscopic confirmation),</li> <li>ii) Reports of surgical or non-surgical management</li> </ul> <p>b) Not described OR used minimal criteria for inclusion/exclusion.</p>
<p>2. Source population</p> <p>Low risk of bias = a High risk if = b, c</p>	<p>a) A consecutive sample or random selection from a source population that is well described and representative of the condition under study (e.g. surgeon's clinic, outpatient clinic).</p> <p>b) A consecutive sample or random selection from a population that is not highly representative of the condition under study.</p> <p>c) Cannot be defined or enumerated (i.e. volunteering or self-recruitment).</p>
<p>3. Typical of the average ACL injured population (representativeness of cohort)</p> <p>Low risk of bias = a High risk if = b</p>	<p>a) Truly representative of the average ACL injured person in the community if <b>all</b> of the following criteria are present:</p> <ul style="list-style-type: none"> <li>i) Including men and women,</li> <li>ii) Typical age range at time of ACL injury/surgery (mean age = 16-35),</li> <li>iii) If surgery, then 'typical' surgical procedure (arthroscopic and not synthetic graft*)</li> </ul> <p><i>*If non-surgical management then N/A for this point</i></p> <p>b) Above criteria are not present then not truly representative of the average ACL injured population.</p>
<p>4. Sample size</p> <p>Low risk of bias = a High risk if = b, c</p>	<p>a) Power analysis completed and sample size adequate to detect meaningful difference.</p> <p>b) Power analysis completed but sample size not adequate to detect meaningful difference.</p> <p>c) No power analysis completed.</p>

Exposure	
<p>5. Methods for assessment of functional performance (i.e. ascertainment of exposure)</p> <p>Low risk of bias = a High risk if = b, c</p>	<p>a) Well described methods for functional tests - including an appropriately trained or appropriate profession as assessor <b>AND</b> describes or cites reliability.</p> <p>b) Well described methods for functional tests including an appropriately trained or appropriate profession as assessor) <b>OR</b> describes or cites reliability.</p> <p>c) Not described.</p>
<p>6. Demonstration that outcome of interest was not present at ascertainment of exposure (i.e. outcome that is compared to exposure)</p> <p>Low risk = a High risk if = b</p>	<p>a) <b>True</b> if baseline score of outcome of interest for both exposed/non-exposed (poor/good functional performance) has been accounted for (for example as a covariate or change in score or not present at ascertainment of exposure).</p> <p>b) No demonstration that the baseline score of outcome of interest has been accounted for.</p>
Comparability	
<p>7. Comparability of cohorts on the basis of the design or analysis</p> <p>Low risk = a High risk if = b</p>	<p>a) Comparability exists if study cohort (exposed/non-exposed) was a priori matched for <b>at least one covariate</b>, or confounding controlled for in statistical analysis.</p> <p>Covariate examples:</p> <ul style="list-style-type: none"> <li>i) Age,</li> <li>ii) BMI,</li> <li>iii) Sex</li> </ul> <p>b) Study not controlled in design or analysis and no confounders acknowledged.</p> <p>(Statements of no differences between groups or that differences were not statistically significant are not sufficient for establishing comparability)</p>

Outcome	
<p>8. Validity and reliability of outcome(s) of interest</p> <p>Low risk = a High risk if = b</p>	<p>a) Outcome measure(s) of interest are clearly described, and references other article(s) which found outcome measure to be valid &amp; reliable <b>OR</b> demonstrates the outcome measure(s) of interest are valid and reliable. (note all outcome(s) of interest must be valid and reliable for (a))</p> <p>b) If outcome measure(s) of interest were not explained in reproducible detail, or validity and reliability not proven/reported.</p>
<p>9. Assessment of outcome(s) of interest</p> <p>Low risk = a High risk if = b</p>	<p>a) Assessor has suitable qualification to interpret findings (e.g. musculoskeletal radiologist) <b>AND</b> blind to participant baseline exposure/non-exposure. <i>*N/A: Blinding not needed for self-reported outcomes</i></p> <p>b) Poor or no description.</p>
<p>10. Adequacy of follow-up of cohorts</p> <p>Low risk = a High risk if = b</p>	<p>a) Adequacy of follow-up if <b>either</b> of the following are satisfied:</p> <ul style="list-style-type: none"> <li>i) &lt;15% lost to follow up + description of those lost,</li> <li>ii) &lt;5% lost to follow up with no description</li> </ul> <p>b) &gt;15% lost to follow up or not explicitly stated with number of participants lost to follow-up <b>OR</b> characteristics of those lost to follow-up were not described.</p>

\*risk of bias assessed from published paper, not considering extra data if provided by the authors

**Appendix 3. Measures used for outcomes presented**

Paper	Measure used
<b>Return to Sport</b>	
Ardern et al, 2015*	Are you currently playing sport at the same level that you played pre-injury?
Ebert and Annear, 2019*	Tegner activity scale – Pre-injury level
Faleide et al, 2021	At what level do you perform your main sport/activity now? (pre-injury)
Kitaguchi et al, 2020	Tegner activity scale – Pre-injury level
McGrath et al, 2017*	Tegner activity scale – Pre-injury level
Moksnes and Risberg, 2009*	IKDC - pre-injury activity level
Nawasreh et al, 2017*	Global Rating scale – pre-injury level
Toole et al, 2017*	Tegner activity scale – maintain or improve
Webster et al, 2019*	Marx Scale – Pre-injury level
Welling et al, 2020*	Did you return to the pre-injury level of sport?
<b>Symptoms and function</b>	
Cristiani et al, 2020*	KOOS – Meeting Englund criteria
Culvenor et al, 2016*	KOOS – Meeting Englund criteria
Ebert and Annear, 2019*	IKDC – PASS (>75.9)
Ericsson et al, 2013*	KOOS – Meeting Englund criteria
Filbay et al, 2021*	KOOS – Meeting Englund criteria
Logerstedt et al, 2012*	IKDC – Below 15th percentile
Mansson et al, 2013*	KOOS – Meeting Englund criteria
Oiestad et al, 2012*	KOOS – Meeting Englund criteria
Stropnik et al, 2020	IKDC – Below 15th percentile
Welling et al, 2020*	IKDC – PASS (>75.9)
<b>Success with ACL deficiency</b>	
Button et al, 2006*	Return to pre-injury activity level (phone questionnaire)
Eitzen et al, 2010	Not having delayed ACLR up to 15/12 post-injury
Ekas et al, 2019	Not having delayed ACLR up to 8-years post-injury
Ericsson et al, 2013*	Not having delayed ACLR up to 5-years post-injury
Fitzgerald et al, 2000	Ability to return to preinjury levels of activity without experiencing an episode of giving-way at the knee
Grindem et al, 2018*	Delayed ACLR up to 2-years post-injury
<b>Subsequent knee injury</b>	



Beischer et al, 2020	ACL re-rupture or contralateral injury up to 46-months post-surgery
Cristiani et al, 2021	Revision ACLR within 2-years post-ACLR
Faleide et al, 2021	ACL re-rupture or contralateral ACL rupture up to 2-years post-surgery
Faltstrom et al, 2021	ACL re-rupture, contralateral re-rupture or "severe" injury (absence from soccer play $\geq 28$ days) up to 2-years post-surgery
Grindem et al, 2016*	ACL re-rupture and other injuries to ACLR knee or contralateral rupture/injury up to 2-years post-surgery
King et al, 2021	Contralateral ACL injury within 2-years post-ACLR
Kyritsis et al, 2016*	ACL graft re-rupture up to 2.5 years after surgery
Nawasreh et al, 2016	ACL re-rupture or contralateral injury up to 2-years post-surgery
Sousa et al, 2017	ACL re-rupture or contralateral injury 4-years post-surgery
van Melick et al, 2021*	ACL re-rupture or contralateral ACL injury up to 2-years post-surgery
Webster et al, 2019	ACL re-rupture or contralateral ACL injury up to 5-years post-surgery
Wellsandt et al, 2017*	ACL re-rupture or contralateral injury $\geq 2$ -years post-surgery
<b>Knee Osteoarthritis</b>	
Filbay et al, 2021*	Kellgren & Lawrence OA grading $\geq 2$ and defined symptoms
Janssen et al, 2013*	A combination of Ahlbäck grade 1 and Kellgren & Lawrence (K&L) grade 3 was defined as 'radiographic signs of knee OA'
Oiestad et al, 2012*	Kellgren & Lawrence OA grading score $\geq 2$ (Extra data received from author for this analysis)
Patterson et al, 2018*	OARSI criteria; i) JSN of grade 2 or higher, ii) Sum of osteophyte grades $\geq 2$ , iii) Grade 1 JSN in combination with a grade 1 osteophyte
Pinczewski et al, 2007*	IKDC OA grading
Wellsandt et al, 2018*	Kellgren & Lawrence OA grading $\geq 2$

**Studies not included in meta-analysis**

Flosadottir et al, 2016	Tegner activity scale
Flossadottir et al, 2018	Knee – self efficacy scale
Kline et al, 2016	Biomechanical variables

KOOS, The Knee Injury and Osteoarthritis Outcome Score, IKDC, International Knee Documentation Committee score, ACLR, anterior cruciate ligament reconstruction, OA, osteoarthritis, JSN, joint space narrowing,

\*Extra data received from author to report outcomes in this format for our analysis

**Appendix 4.** Tests used in each study

Study	Single-forward hop (n=42)	Repeated forward hops			Side hop (n=5)	One leg rise (n=4)	Vertical or square hop (n=2)	Battery of tests (n=15)
		Triple hop (n=18)	Crossover hop (n=23)	6-m timed hop (n=15)				
<b>Return to Sport</b>								
Ardern et al, 2015								
Ebert et al, 2019								
Faleide et al, 2019								
Kitaguchi et al, 2020								
McGrath et al, 2017								
Moksnes et al, 2009								
Nawasreh et al, 2017								
Toole et al, 2017								
Webster et al, 2019								
Welling et al, 2020								
<b>Patient-reported symptoms and function</b>								
Cristiani et al, 2020								
Culvenor et al, 2016								
Ericsson et al, 2013								
Filbay et al, 2021								
Logerstedt et al, 2012								
Mansson et al, 2013								

Oiestad et al, 2012								
Stropnik et al, 2020								
<b>Success with ACL deficiency</b>								
Button et al, 2006								
Eitzen et al, 2010								
Ekas et al, 2019								
Fitzgerald et al, 2000								
Grindem et al, 2018								
<b>Subsequent knee injury after ACLR</b>								
Beischer et al, 2020								
Cristiani et al, 2021								
Faltstrom et al, 2021								
Grindem et al, 2016								
King et al, 2021								
Kyritsis et al, 2016								
Nawasreh et al, 2016								
Sousa et al, 2017								
van Melick et al, 2021								
Webster et al, 2019								
Wellsandt et al, 2017								
<b>Knee osteoarthritis</b>								
Filbay et al, 2021								
Janssen et al, 2013								

Patterson et al, 2018								
Pinczewski et al, 2007								
Wellsandt et al, 2018								

**Studies not included in meta-analysis**

Flosadottir et al, 2016								
Flosadottir et al, 2018								
Kline et al, 2016								

ACLR, anterior cruciate ligament reconstruction

Green (darker) shaded cells indicate where test was used within study



## Appendix 5. Risk of bias using a Modified Newcastle Ottawa Scale

Article	Selection bias				Exposure bias		Comparability bias	Outcome bias		
	Definition of ACL injured population	Source population	Typical of average ACL injured population	Sample Size	How performance using a functional test was decided	Demonstrates outcome was not present at time of functional test	Comparability of cohorts on the basis of the design or analysis	Validity and reliability of outcomes	Assessment of outcome*	Adequacy of follow up
<i>Return to Sport</i>										
Ardern et al, 2015	Low	Low	Low	High	Low	Low	High	High	Low	High
Ebert et al, 2019	Low	High	High	High	High	Low	High	Low	Low	High
Faleide et al, 2019	Low	Low	Low	High	High	Low	Low	High	Low	High
Kitaguchi et al, 2020	Low	Low	Low	High	Low	Low	High	High	Low	High
McGrath 2017	Low	High	High	Low	Low	Low	Low	Low	Low	High
Moksnes et al, 2009	Low	Low	Low	High	Low	Low	High	Low	Low	High
Nawasreh et al, 2017	Low	High	Low	High	High	High	Low	High	High	High
Toole et al, 2017	Low	High	Low	High	High	Low	High	Low	Low	Low
Webster et al, 2019	Low	Low	Low	High	High	Low	High	Low	Low	High
Welling et al, 2020	Low	Low	Low	High	High	High	High	High	Low	High
<i>Patient-reported symptoms and function</i>										
Cristiani et al, 2020	Low	Low	Low	Low	High	High	Low	Low	Low	High
Culvenor et al, 2016	Low	Low	Low	High	High	Low	Low	Low	Low	Low
Ericsson et al, 2013	Low	Low	Low	High	Low	Low	Low	Low	Low	High
Filbay et al, 2021	Low	Low	Low	Low	High	Low	Low	Low	Low	High
Logerstedt et al, 2012	Low	Low	Low	High	Low	Low	High	Low	Low	High
Mansson et al, 2013	Low	High	Low	High	High	High	High	Low	Low	High
Oiestad et al, 2012	Low	High	Low	High	High	High	Low	Low	Low	High

Stropnik et al, 2020	Low	Low	Low	High	High	Low	High	Low	Low	Low
<i>Success with ACL deficiency</i>										
Button et al, 2006	Low	High	Low	High	High	Low	High	High	Low	High
Eitzen et al, 2010	Low	Low	Low	High	Low	Low	High	High	High	Low
Ekas et al, 2019	Low	Low	High	High	High	Low	High	Low	Low	Low
Fitzgerald et al, 2000	Low	Low	Low	High	High	Low	High	High	High	Low
Grindem et al, 2018	Low	Low	Low	High	High	Low	Low	Low	High	Low
<i>Subsequent knee injury after ACLR</i>										
Beischer et al, 2020	Low	Low	Low	High	Low	Low	High	High	High	High
Cristiani et al, 2021	Low	Low	Low	High	Low	Low	Low	Low	Low	High
Faltstrom et al, 2021	Low	High	High	High	Low	Low	Low	Low	High	Low
Grindem et al, 2016	Low	Low	Low	High	High	Low	Low	Low	Low	High
King et al, 2021	Low	Low	High	High	High	Low	Low	Low	High	High
Kyritsis et al, 2016	Low	Low	High	High	High	Low	Low	High	High	Low
Nawasreh et al, 2016	Low	High	Low	High	High	Low	High	High	High	High
Sousa et al, 2017	Low	Low	Low	Low	High	Low	High	High	High	High
van Melick et al, 2021	Low	Low	Low	High	Low	Low	High	High	Low	High
Webster et al, 2019	Low	Low	Low	High	Low	Low	High	High	Low	High
Wellsandt et al, 2017	Low	High	Low	High	High	Low	High	High	High	High
<i>Knee Osteoarthritis</i>										
Filbay et al, 2021	Low	Low	Low	High	High	High	Low	Low	Low	High
Janssen et al, 2013	Low	Low	Low	High	Low	Low	High	Low	Low	Low
Patterson et al, 2018	Low	High	Low	High	High	Low	Low	Low	Low	High
Pinczewski et al, 2007	Low	High	Low	High	High	High	High	High	High	High

Wellsandt et al, 2018	Low	High	Low	High	High	High	High	Low	High	High
<i>Studies not included in meta-analysis</i>										
Flosadottir et al, 2016	Low	Low	Low	High	Low	Low	Low	Low	Low	High
Flosadottir et al, 2018	Low	Low	Low	High	Low	High	Low	Low	Low	Low
Kline et al, 2016	Low	High	Low	High	High	High	High	High	High	High

## Appendix 6. GRADE assessment for the single leg hop test applied to all outcomes\*

	Study design	Risk of bias	Inconsistency (I <sup>2</sup> )	Indirectness	Imprecision	Publication bias	Odds Ratio	GRADE
<i>Return to Sport</i>								
Ardern et al, 2015 Ebert et al, 2019 Moksnes et al, 2009 Nawasreh et al, 2017 Toole et al, 2017 Faleide et al, 2021 Kitaguchi et al, 2020 Webster et al, 2019 Welling et al, 2020	(-2) Prospective	(-1) Serious: Most did not control for confounders, had inadequate sample size and poor follow-up	(-1) Not serious: I <sup>2</sup> = 70.6% (P= 0.001) Overlapping confidence intervals from 8/10 studies. Odds ratio between 0.94 to 16.44 without large discrepancies in weighting	(0) Not serious Similar populations, timepoints of testing and outcomes	(0) Total n= 638 CI (1.30, 3.54)	N/A	(0) 1.97 (1.24, 3.13)	Very low
<i>Patient-reported symptoms and function</i>								
Cristiani et al, 2020 Culvenor et al, 2016 Ebert et al, 2019 Ericsson et al, 2013 Filbay et al, 2021 Logerstedt et al, 2012 Mansson et al, 2013 McGrath et al, 2017 Oiestad et al, 2012 Stropnik et al, 2020 Welling et al, 2020	(-2) Prospective	(-1) Serious: Most did not control for confounders, had inadequate sample size and poor follow-up	(-1) Serious: I <sup>2</sup> = 69.2% (P= 0.001) Overlapping confidence intervals of most studies with two outlying studies. Odds ratio between 1.28 to 18.95. Discrepancies in weighting.	(-1) Serious Differing populations (age), differing timepoints of outcome, and outcomes used (IKDC and KOOS)	(0) Total n=1737 CI (1.62, 3.88)	(-1) Serious P = 0.002 Eggers test for small study effects (as there were 10 studies)	(+1) 2.51 (1.62, 3.88)	Very low
<i>Success with ACL deficiency</i>								
Button et al, 2006 Eitzen et al, 2010 Ekas et al, 2019 Ericsson et al, 2013 Fitzgerald et al, 2000 Grindem et al, 2018 Mosknes et al, 2009	(-2) Prospective	(-1) Serious: Most did not control for confounders, had inadequate sample size and poor follow-up	(0) Not serious: I <sup>2</sup> = 54.1% (P= 0.042) Overlapping confidence intervals from 5/7 studies. Odds ratio between 0.59 to 6.65 without large discrepancies in weighting	(-1) Serious Differing outcome definitions	(0) Total n= 228 CI (0.75, 2.32)	N/A	(0) 1.32 (0.75, 2.32)	Very low
<i>Subsequent knee-injury after ACLR</i>								
Cristiani et al, 2021 Falstrom et al, 2021 Grindem et al, 2016 King et al, 2021 Kyritsis et al, 2016 Webster et al, 2019 Wellstandt et al, 2017	(-2) Prospective	(-1) Serious: Most did not control for confounders, had inadequate sample size and poor follow-up	(0) Not serious: I <sup>2</sup> = 47.2% (P= 0.078)	(-1) Serious Different outcome definition for knee-injury or re-injury	(0) Total n= 6970 CI (0.58, 1.11)	N/A	(0) 0.81 (0.58, 1.11)	Very low


Knee Osteoarthritis								
Filbay et al, 2021 Janssen et al, 2013 Patterson et al, 2018 Pinczewski et al, 2007 Wellstandt et al, 2018	<b>(-2)</b> Prospective	<b>(-1) Serious:</b> Most did not control for confounders, had inadequate sample size and poor follow-up	<b>(-1) Serious:</b> I <sup>2</sup> = 75.8% (P= 0.006)	<b>(-1)</b> ? maybe high due to timing of outcome (and therefore population) and outcome definition	<b>(-1)</b> Total n= 222 CI (0.70, 3.98)* wide	N/A	<b>(0)</b> 1.67 (0.70, 3.98)	Very low

Grade of evidence was assigned using the GRADE system, which has 4 categories HIGH, MODERATE, LOW or VERY LOW. Evidence is initially assigned as HIGH from randomised trials. The grade of evidence was then reduced if there was serious (-1) or very serious (-2) limitations to study quality or uncertainties about directness of association; important inconsistency (-1), imprecise or sparse data (-1) or a high probability of reporting bias (-1). Grade of evidence was increased if strong evidence of association was seen (e.g., RR >2 or <0.5) from ≥2 observational studies with no plausible confounders (+1) or very strong direct evidence (RR >5 or <0.2) with no major threats to validity (+2); if there was evidence of a dose-response gradient (+1) or if all plausible confounders would have reduced the effect/association seen (+1). The interpretation of GRADE evidence assessments is that for HIGH certainty evidence further research is very unlikely to change our confidence in the estimate of effect; for MODERATE certainty evidence further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate; for LOW certainty evidence further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate; and for VERY LOW certainty evidence any estimate of effect is very uncertain.

\*Only level of evidence of the single-forward hop test was assessed as this was used across all outcomes and studies. The level of evidence was found to be very low across all outcomes using this test and so we decided there was not need to complete this assessment for other tests as they would likely yield the same result and have less data from which to draw conclusions of evidence certainty.

KOOS, Knee osteoarthritis outcome score, IKDC, international knee documentation committee score

## Funnel plot with pseudo 95% confidence limits



Number of studies = 10



Root MSE = 1.042

Std_Eff	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
slope	.095743	.1061769	0.90	0.394	-.1491014	.3405874
bias	2.014122	.4626747	4.35	0.002	.9471918	3.081052



Test of H0: no small-study effects

P = 0.002







**Better self-reported symptoms and function**




**Appendix 10.** Sensitivity analysis of repeated forward hop tests (i.e., crossover, triple, 6m-timed) - stratified meta-analysis


Outcomes (Individual hop test subgroup totals)	Odds ratio (95%CI)	Study (n)	Participants (n)	Between hop test heterogeneity (I <sup>2</sup> & p-value)
<b>Return to sport</b>				I <sup>2</sup> = 71.8% p=0.645
Crossover	1.77 (1.05, 2.96)	7	781	
Triple	2.54 (1.09, 5.92)	6	443	
6m-timed	2.55 (1.2, 5.54)	6	529	
<b>Self-reported symptoms and function</b>				I <sup>2</sup> = 51.9% p=0.978
Crossover	4.96 (1.55, 15.85)	3	228	
Triple	4.48 (1.5, 13.33)	4	380	
6m-timed	5.2 (2.16, 12.53)	2	135	
<b>Success with ACL deficiency</b>				I <sup>2</sup> = 19.6% p=0.843
Crossover	1.51 (0.94, 2.44)	5	437	
Triple	1.27 (0.87, 1.86)	5	337	
6m-timed	1.43 (0.78, 2.63)	5	385	
<b>Subsequent injury after ACL reconstruction</b>				I <sup>2</sup> = 1.2% p=0.175
Crossover	1.03 (0.74, 1.44)	4	663	
Triple	0.85 (0.47, 1.54)	3	334	
6m-timed		2	176	
<b>Knee osteoarthritis</b>				I <sup>2</sup> = 76.7% p=0.014
Crossover	0.141 (0.05, 0.34)	1	78	
Triple	0.737 (0.38, 1.42)	1	181	
6m-timed	0.584 (0.15, 2.27)	1	58	















## Appendix 16. Meta-regression results for studies reporting continuous hop scores and association to each outcome

### Single-forward hop scores

Outcome	Coefficient (95%CI)	P-value	Adjusted r <sup>2</sup>
<b>Follow-up time</b>			
Return to sport	-0.01 (-0.07, 0.04)	0.63	20.74%
Self-reported symptoms and function	-0.00 (-0.003, 0.001)	0.22	3.79%
Success with ACL deficiency	0.00 (-0.003, 0.001)	0.90	52.48%
Subsequent injury after ACL reconstruction	-0.01 (-0.01, 0.01)	0.11	68.43%
Knee osteoarthritis	-0.00 (-0.006, 0.009)	0.48	8.08%
<b>Proportion of females</b>			
Return to sport	-0.49 (-3.48, 2.48)	0.70	23.22%
Self-reported symptoms and function	-0.84 (-5.12, 3.43)	0.66	23.97%
Success with ACL deficiency	-1.63 (-6.09, 2.82)	0.39	9.15%
Subsequent injury after ACL reconstruction	-0.14 (-0.96, 0.67)	0.67	20.15%
Knee osteoarthritis	-2.14 (-26.73, 22.43)	0.74	60.84%

### Repeated-forward hop scores

Outcome	Coefficient (95%CI)	P-value	Adjusted r <sup>2</sup>
<b>Follow-up time</b>			
Return to sport	0.00 (-0.08, 0.09)	0.90	28.73%
Self-reported symptoms and function	0.00 (-0.01, 0.005)	0.21	39.40%
Success with ACL deficiency	0.00 (-0.01, 0.02)	0.57	-*
Subsequent injury after ACL reconstruction	0.00 (-0.02, 0.01)	0.61	-*
Knee osteoarthritis	0.00 (-0.06, 0.07)	0.52	1.72%
<b>Proportion of females</b>			
Return to sport	-1.77 (-4.77, 1.21)	0.20	15.36%
Self-reported symptoms and function	-6.73 (-26.14, 12.68)	0.35	2.81%
Success with ACL deficiency	-2.89 (-7.33, 1.53)	0.13	-*
Subsequent injury after ACL reconstruction	0.21 (-2.22, 2.65)	0.74	-*
Knee osteoarthritis	3.93 (-170.94, 178.81)	0.82	95.31%

ACL, anterior cruciate ligament

\*Indicates values that were not able to be calculated