

**Supplementary file 3.** Proposed cluster symptoms and relative rationale for selection.

<b>Main symptoms</b>	<b>Question</b>	
Urinary incontinence (Any type)	Do you usually experience urine leakage?	PFDI-20, section for urinary symptoms related to UDI-6. Summary of #16,17,18.
Overactive bladder syndrome	Do you usually experience urinary urgency (that is a strong sensation of needing to go to the bathroom) usually accompanied by frequent urination and nocturia?	OAB-V8 Overactive Bladder-Validated 8-question Screener. Adaptation from OAB-V8 and definition of OAB: Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. <i>Neurourol Urodyn.</i> 2002;21:167-178
Pelvic Organ Prolapse	Do you usually have a bulge or something falling out that you can see or feel in your vaginal area?	National Guideline Alliance (UK). Evidence review for assessing pelvic organ prolapse: Urinary incontinence and pelvic organ prolapse in women: management. London: National Institute for Health and Care Excellence (NICE); April 2019.
	Have you ever had to push in the perineal area with your fingers to start or complete a bowel movement or to start or complete urination?	National Guideline Alliance (UK). Evidence review for assessing pelvic organ prolapse: Urinary incontinence and pelvic organ prolapse in women: management. London: National Institute for Health and Care Excellence (NICE); April 2019.
Anal incontinence	Do you usually lose stool or gas beyond your control?	PFDI-20, section of colorectal-anal symptoms related to CRAD-8, integration of #10,11.
Pelvic pain	Do you usually experience pain or discomfort in the lower abdomen or genital region?	PFDI-20, section for urinary symptoms related to UDI-6, #20