



OPEN ACCESS

When ACL reconstruction does not help: risk factors associated with not achieving the minimal important change for the KOOS Sport/Rec and QoL

Janina Kaarre ,^{1,2,3} Rebecca Simonson ,^{2,4} Viktor Ris,^{1,2} Thorkell Snaebjörnsson,^{1,2} James J Irrgang,^{5,6} Volker Musahl,^{3,6} Kristian Samuelsson,^{1,2,7} Eric Hamrin Senorski ^{2,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2022-106191>).

¹Department of Orthopaedics, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

²Sahlgrenska Sports Medicine Center, Gothenburg, Sweden

³Department of Orthopaedic Surgery, UPMC Freddie Fu Sports Medicine Center, University of Pittsburgh Medical Center, Pittsburgh, PA, USA

⁴Unit of Physiotherapy, Department of Health and Rehabilitation, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

⁵Department of Physical Therapy, University of Pittsburgh School of Health and Rehabilitation Sciences, Pittsburgh, PA, USA

⁶Department of Orthopaedic Surgery, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

⁷Department of Orthopaedics, Sahlgrenska University Hospital, Mölndal, Sweden

Correspondence to

Dr Janina Kaarre, Department of Orthopaedics, University of Gothenburg, Mölndal 431 80, Sweden; janina.kaarre@gu.se

Accepted 20 February 2023
Published Online First
1 March 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Kaarre J, Simonson R, Ris V, et al. *Br J Sports Med* 2023;**57**:528–535.

ABSTRACT

Objectives To determine factors associated with not achieving a minimal important change (MIC) in the Knee injury and Osteoarthritis Outcome Score (KOOS) Function in Sport and Recreation (Sport/Rec), and Knee-Related Quality of Life (QoL) subscales 1 year after anterior cruciate ligament reconstruction (ACL-R).

Methods This study used data from the Swedish National Knee Ligament Registry. Multivariable logistic regression models were used to identify factors associated with not achieving a MIC. The change in the preoperative and postoperative KOOS Sport/Rec and QoL subscale scores were dichotomised based on not achieving MIC for both subscales versus achieving MIC for either one or both subscales. The MICs for the Sport/Rec and QoL subscales were 12.1 and 18.3, respectively, and were used to combine both subscales into a single variable (Sport & QoL).

Results Of 16 131 included patients, 44% did not achieve the MIC for the combined Sport/Rec and QoL subscales 1 year after ACL-R. From the multivariable stepwise logistic regression, older patients (OR 0.91, 95% CI 0.88 to 0.94; $p<0.0001$), males (OR 0.93, 95% CI 0.87 to 0.99; $p=0.034$) and patients receiving hamstring tendon autograft ACL-R (OR 0.70, 95% CI 0.60 to 0.81; $p<0.0001$) had lower odds of not achieving the MIC 1 year after ACL-R compared with younger patients, females and patients receiving patellar tendon autograft. Furthermore, patients with cartilage injuries (OR 1.17, 95% CI 1.09 to 1.27; $p<0.0001$) and higher pre-operative KOOS Sport/Rec and QoL scores (OR 1.34, 95% CI 1.31 to 1.36; $p<0.0001$) had higher odds of not achieving the MIC.

Conclusion Younger patients, females and patients with cartilage injuries and higher pre-operative Sport/Rec and QoL KOOS scores are less likely to benefit from ACL-R and subsequently, have a lower probability for improved Sport/Rec and QoL scores after ACL-R. Furthermore, graft choice may also affect the risk of not achieving the MIC.

Level of evidence Retrospective cohort study, level III.

INTRODUCTION

Injury to the anterior cruciate ligament (ACL) is one of the most common severe injuries in the knee leading to decreased knee function, activity level and quality of life (QoL).^{1 2} Currently, ACL injuries can be treated with either rehabilitation alone or combined with surgical reconstruction ACL

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Patient-reported outcomes (PROs) are necessary to evaluate outcomes following anterior cruciate ligament reconstruction (ACL-R).
- ⇒ The minimal important change (MIC) for different PROs has recently received increased interest in orthopaedic research and has been subsequently used to evaluate effects of patient characteristics on post-operative outcomes.
- ⇒ Although improvements in the Knee injury and Osteoarthritis Outcome Score subscale scores have been associated with ACL-R, there are still patients who do not achieve the MIC after ACL-R.

WHAT THIS STUDY ADDS

- ⇒ The increased understanding of the patients who do not benefit from the surgical treatment have a high clinical relevance and subsequently, can help tailor treatment for individual patients.
- ⇒ Achievement of the MIC after ACL-R may be multifactorial and therefore the identified possible risk factors should be considered in a treatment plan to establish realistic expectations for the patient.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Clinicians should be aware of both modifiable and non-modifiable risk factors for not achieving an MIC after ACL-R and thus, should adjust the treatment approach and establish realistic expectations for the patient.
- ⇒ The MIC values may be different for patients with different baseline PRO scores requiring further research.

(ACL-R), where the surgical treatment is reported to improve the likelihood of returning to more knee-strenuous sports (RTS).^{3 4} During the past decades, there has been an increasing interest in using patient-reported outcomes (PROs), such as the Knee injury and Osteoarthritis Outcome Score (KOOS), to evaluate outcomes related to different treatment options for patients with ACL injuries.^{4 5} In addition, different patient characteristics (sports participation, age), injury characteristics (concomitant injuries) as well as surgical factors (graft diameter, graft fixation, time from injury to surgery)

Table 1 Baseline characteristics of patients

Variable	Total (n=16 131)
Age at time of injury (years)	27.1±10.3 24 (1–73)
Age at time of surgery (years)	28.9±10.7 26 (16–74)
Sex (male), n (%)	8274 (51.3)
BMI (kg/m ²)	24.6±3.3 24.2 (15.1–45.4)
Smoking (yes), n (%)	525 (4.4)
Time from injury to surgery (months)	20.1±37.2 7.9 (0–551)
ACL graft, n (%)	
Patellar tendon autograft	768 (4.8)
Hamstring tendon autograft	14835 (93.3)
Quadriceps tendon autograft	237 (1.5)
Allograft	32 (0.2)
Direct suture/synthetic/other	32 (0.2)
Concomitant injury (yes), n (%)	9240 (57.3)
Meniscus injury (yes), (%)	7086 (43.9)
Lateral meniscus injury	4016 (24.9)
Medial meniscus injury	4234 (26.2)
Cartilage injury (yes), (%)	4356 (27.0)
Lateral femoral condyle	886 (20.4)
Medial femoral condyle	2932 (67.6)
Lateral patella	537 (12.4)
Medial patella	871 (20.1)
Lateral tibial plateau	1083 (25.0)
Medial tibial plateau	826 (19.0)
Trochlea	530 (12.2)
Collateral ligament injury (yes), n (%)	653 (4.0)
LCL	107 (0.7)
MCL	562 (3.5)
PLC injury (yes), n (%)	19 (0.1)
Values are given as n (%) and mean±SD or median (minimum–maximum) for categorical and continuous variables, respectively. The sums may vary due to missing values. The variables with missing values, n (%) of the total sample were age at time of injury 366 (2.3); BMI 4490 (27.8); smoking 4305 (26.7); time from injury to surgery 382 (2.4); ACL graft 227 (1.4); cartilage injury to lateral and medial femoral condyle, lateral and medial patella, lateral and medial tibial plateau as well as trochlea 11 795 (73.1).	
ACL, anterior cruciate ligament; BMI, body mass index; LCL, lateral collateral ligament; MCL, medial collateral ligament; PLC, posterolateral corner.	

have been demonstrated to affect post-operative outcomes,^{6–12} and subsequently, contribute to variation in outcomes following ACL-R. A growing interest has recently developed in estimating the minimal important change (MIC), defined as the smallest change in outcome that a patient perceives as important,⁶ for different PROs. The MICs have been used to identify the effects of demographic and surgical factors on post-operative outcomes for individuals undergoing ACL-R to evaluate.^{6 13}

While a large part of previous studies investigating improvement in knee function following ACL-R primarily focus on evaluating changes in PRO scores, less is known about the patients not achieving an MIC for specific combined KOOS subscales.^{4 14 15} An increased understanding of the patients who do not benefit from ACL-R can have a high clinical relevance and help tailor treatment for individual patients. Therefore, the purpose of this study was to determine patient and surgical risk factors for not achieving the MIC in both the KOOS Function in Sport and

Recreation (Sport/Rec), and Knee-Related QoL subscales 1 year after ACL-R.

METHODS

The study is presented according to the Strengthening the Reporting of Observational studies in Epidemiology guidelines.¹⁶

This registry study was based on data obtained from the Swedish National Knee Ligament Registry (SNKLR). The purpose of the SNKLR is to collect data on patients undergoing treatment for ACL injury and uses a web-based protocol for data registration consisting of patient-reported and surgeon-reported sections.¹⁷ Surgeons report on patient demographic characteristics, injury-related and surgery-related factors, while patients answer questionnaires regarding knee function, including the KOOS Sport/Rec and QoL subscales. The SNKLR has previously been described in detail.¹⁸

Study population

Data on patients ≥15 years of age at time of primary ACL-R with pre-operative and 1-year post-operative KOOS data in the SNKLR between 1 January 2006 and 30 October 2019 were extracted for this study. The study population was divided into two groups: (1) those that did not achieve the MIC for the KOOS Sport/Rec and QoL subscales and (2) those that achieved the MIC for both or only one of the subscales 1 year after ACL-R.

Patients with any prior knee surgery, concomitant fracture, nerve, or vascular injury, concomitant posterior cruciate ligament (PCL) injury or surgically treated medial collateral ligament (MCL) or lateral collateral ligament injuries were excluded. Patients treated with double bundle ACL-R or revision ACL-R were also excluded from the analysis. Furthermore, patients with either high preoperative KOOS Sport/Rec (>87.9) or QoL (>81.7) scores were excluded since the achievement of the MIC could not be observed in these patients.¹⁹

Data regarding patient demographics (sex, age, body mass index (BMI), smoking status), injury characteristics (activity at time of injury, concomitant injury) and surgical factors (time from injury to surgery, graft type) were also retrieved from the SNKLR. The KOOS Sport/Rec and QoL subscale scores before surgery and at 1-year follow-up were also extracted from the SNKLR. Activities at the time of injury were further divided into sports-related (alpine/skiing, pivoting sport, non-pivoting sport, other physical activity) and non-sports-related activities (other, traffic-related activities).

Outcome measures

The KOOS consists of 5 subscales including Pain, Other Symptoms, Sport/Rec, QoL and Activities of Daily Living, each of which are scaled from 0 to 100, where a higher score indicates a better outcome.²⁰ The KOOS has been previously validated and specifically described to be a reliable assessment tool to measure function and QoL in patients undergoing ACL-R,^{19 21} with intra-class correlation coefficients reported to be 0.75 and 0.89 for KOOS Sport/Rec and QoL, respectively.²¹ The threshold scores for achievement of the MIC have been previously calculated using an anchor-based approach that made use of a 7-point global rating of change question as the external anchor for an important change and use of predictive modelling methods and receiver operating characteristic (ROC) curve analyses. The MICs were found to be 12.1 and 18.3 for Sport/Rec and QoL, respectively.¹⁹

The main outcomes of interest for this study were the KOOS Sport/Rec and QoL subscale scores, which were dichotomised

Table 2 Activities at time of injury for patients undergoing ACL-R

Variable	Total (n=16 131)
Activity at time of injury, n (%)	
Alpine/skiing	2797 (17.4)
Pivoting sport	10237 (63.6)
Non-pivoting sport	684 (4.2)
Other physical activity	668 (4.1)
Traffic related	257 (1.6)
Other	1456 (9.0)
Missing	32

Values are given as n (%).
ACL-R, anterior cruciate ligament reconstruction.

into those not achieving versus achieving the MIC for each subscale. Subsequently, achievement of the MIC for each subscale was combined into a single variable that represented failure to achieve the MIC for both the Sport/Rec and QoL subscales (MIC KOOS Sport & QoL). Thus, the primary outcome for this study was the MIC KOOS Sport & QoL that consisted of patients who failed to achieve the MIC on both the KOOS Sport/Rec and QoL subscales versus those that achieved the MIC in either one or both of the KOOS subscales.¹⁹

Equity, diversity and inclusion

This study included both female and male patients undergoing ACL-R. Our multidisciplinary research team involved both female and male researchers from different medical areas (orthopaedics and physiotherapy) and age categories (junior researchers, professors). Data were collected from the SNKLR and thereby, data collection did not differ between the included patients. All patients undergoing ACL-R, including different ethnicities, culture, socioeconomic level, sex and gender are included in the SNKLR.

Statistical analyses

The statistical analyses were performed with the SAS System for Windows (V.9, SAS Institute). Continuous and ordinal data were presented by using the mean and SD as well as median with minimum and maximum. Count (n) and proportion (%) were used to summarise dichotomous variables. In addition, univariable and stepwise multivariable logistic regression analyses were used to determine demographic and surgical factors associated with not achieving the MIC for both the Sport/Rec and QoL subscales and presented as OR, p value and 95% CI. Variables

Table 3 Pre-operative and post-operative KOOS

Variable	Total (n=16 131)
Pre-operative KOOS	
KOOS-Sport/Rec	36.8±24.7 35 (0–85)
KOOS-QoL	32.0±16.5 31.3 (0–81.3)
Post-operative KOOS	
KOOS-Sport/Rec	63.7±26.7 70 (0–100)
KOOS-QoL	57.9±23.0 62.5 (0–100)

Values are given as mean±SD or median (minimum-maximum) for variables.
KOOS, Knee injury and Osteoarthritis Outcome Score; QoL, quality of life; Sport/Rec, function in sport and recreation.

included in the stepwise multivariable logistic regression analyses were age at time of surgery, sex, meniscal and cartilage injury as well as ACL graft type and pre-operative KOOS Sport/Rec and QoL scores. The OR for patient age and pre-operative KOOS scores, calculated in the univariable and stepwise multivariable logistic regression analysis, were based on a 10-year and 10-point increase, respectively. Furthermore, the area under the curve (AUC) with 95% CI was used to present the results from the univariable analyses. The AUC varies between 0.5 and 1, where a greater AUC represents better predictive capacity. Further, the AUC between 0.5 and 0.7 indicates poor predictive capacity, while the AUC between 0.7 and 0.9 or between 0.9 and 1.00 represents acceptable/good and excellent predictive capacity, respectively. All significance tests were two sided and conducted at the 5% significance level.

Supplemental analyses

Supplemental analyses were performed separately to identify factors associated with not achieving the MIC for either the KOOS Sport/Rec or QoL subscales. Univariable and stepwise multivariable logistic regression analyses were performed to determine demographic and surgical factors associated with not achieving the MIC for the KOOS Sport/Rec or QoL subscales 1 year after ACL-R. Variables included in these stepwise multivariable logistic regression analyses were the same predictor variables that were included in the primary analysis

RESULTS

Baseline characteristics

Of the 16 131 patients included in the study, 8274 were males (51%) and the mean age at time of surgery was 29 years±11. The median time from injury to surgery was 7.9 (0–551) months. A hamstring tendon autograft was found to be the most commonly used autograft (93%), while patellar tendon autograft was used in 4.8% of the cases. Additionally, concomitant injuries were found to be common among the patients (57%), where 27% and 44% of the patients had concomitant cartilage and meniscus injuries, respectively (table 1). Most of the ACL injuries were the result of a pivoting sport activity (64%) (table 2). The average scores for the pre-operative Sport/Rec and QoL subscales for the KOOS were found to be 37±25 and 32±17, respectively. Post-operatively, the average scores for the KOOS Sport/Rec and QoL subscales were 64±27 and 58±23, respectively (table 3).

Minimal important change

In total, 7109 (44%) of patients included in the study did not achieve the MIC for both the Sport/Rec and QoL scales 1 year after ACL-R. Results from univariable logistic regression are presented in the figure 1. From the multivariable stepwise logistic regression, older patients (OR 0.91, 95% CI 0.88 to 0.94; p<0.0001), males (OR 0.93, 95% CI 0.87 to 0.99; p=0.034) and patients undergoing ACL-R with hamstring tendon autograft (OR 0.70, 95% CI 0.60 to 0.81; p<0.0001) had lower odds of not achieving the MIC for both the Sport/Rec and QoL subscales compared with younger patients, females and patients undergoing ACL-R with a patellar tendon autograft. Furthermore, patients with concomitant cartilage injuries (OR 1.17, 95% CI 1.09 to 1.27; p<0.0001) had higher odds for not achieving the MIC for both the Sport/Rec and QoL subscales 1 year after ACL-R compared with patients without concomitant cartilage injuries. In addition, patients with higher pre-operative KOOS Sport/Rec and QoL scores had higher odds for not achieving the MIC for both the Sports/Rec and QoL subscales 1 year after




Figure 1 Forest plot illustrating the OR and 95% CI for predicting not achieving MIC for Combined Sport/Rec and QoL KOOS subscales 1 year after ACL reconstruction. OR>1 = not achieving MIC; OR<1 = achieving MIC. ACL, anterior cruciate ligament; BMI, body mass index; KOOS, Knee Injury and Osteoarthritis Score; MIC, minimal important change; Pre-op, pre-operative; QoL, quality of life; Sport/Rec, function in sport and recreation.

ACL-R (OR 1.34, 95% CI 1.31 to 1.36; $p<0.0001$) (figure 2). Neither activity at time of injury (OR 0.95, 95% CI 0.87 to 1.04; $p=0.30$) nor BMI (OR 1.00, 95% CI 0.99 to 1.01; $p=0.90$) were found to be associated with not achieving the MIC for both the Sport/Rec and QoL subscales 1 year after ACL-R (table 4). The predictive capacity of the multivariable model was relatively poor as the AUC was 0.65 (95% CI 0.64 to 0.66).

Supplemental analyses

In total, 4959 (31%) and 5490 (34%) of included patients did not achieve the MIC for the KOOS Sport/Rec and QoL, respectively. Results from the supplemental univariable and multivariable stepwise logistic regression models were comparable with the results from the univariable and multivariable analyses for the combined KOOS Sport/Rec and QoL scales. Demographic and surgical factors associated with not achieving MIC for the KOOS Sport/Rec or QoL subscales 1 year after ACL-R are presented in online supplemental material tables 5–7.




Figure 2 Diagrammatic overview of the factors associated with achieving and not achieving MIC for Combined Sport/Rec and QoL KOOS subscales (KOOS Sport/QoL) 1 year after ACL reconstruction. ACL, anterior cruciate ligament; KOOS, Knee Injury and Osteoarthritis Score; MIC, minimal important change; QoL, quality of life; Sport/Rec, function in sport and recreation; Sport/QoL, combined Sport/Rec and QoL KOOS subscales.

DISCUSSION

This large, registry-based study identified several factors associated with the risk for not achieving the MIC for both the Sport/Rec and QoL subscales 1 year after ACL-R. Both modifiable and non-modifiable factors were found to be associated with not achieving the MIC for both the Sport/Rec and QoL KOOS subscales 1 year after ACL-R. Approximately 44% of the patients included in this analysis did not achieve a change in both the Sport/Rec and QoL KOOS subscales larger than the MIC 1 year after ACL-R, suggesting that they failed to perceive an important change in both Sport/Rec and QoL.

The results of this study demonstrate that older individuals at the time of ACL-R, males and patients undergoing reconstruction with a hamstring tendon autograft had lower odds of not achieving the MIC for both the Sport/Rec and QoL KOOS subscales 1 year after ACL-R. Furthermore, patients with concomitant injuries to the cartilage and patients with higher pre-operative KOOS Sport/Rec and QoL scores had higher odds of not achieving the MIC for both the Sport/Rec and QoL KOOS subscales 1 year after ACL-R. As such, our results suggest that the association between injury-related, surgical-related and patient-related factors and not achieving the MIC for the Sport/Rec and knee-related QoL is multifactorial.

There are several factors that might contribute to our results suggesting lower odds for not achieving the MIC in older patients, where, for instance, older individuals tend to be less physically active and this group of patients usually includes a smaller proportion of athletes compared with a younger population.²² The lower odds of not achieving the MIC for both the Sport/Rec and QoL subscales 1 year after ACL-R in older population, may be explained by the difference in expected knee function following ACL-R, where older individuals might have lower expectations on future knee function compared with younger individuals.²³

In agreement with previous research, males were found to have better outcomes following ACL-R.^{24–26} This was supported by our results that found that males had lower odds of not achieving the MIC for both the Sport/Rec and QoL KOOS subscales 1 year after ACL-R. This association between male sex and better post-operative outcomes following ACL-R may

Table 4 Results from univariable and multivariable logistics regression analysis. Prediction for not achieving the MIC for Combined Sport/Rec and QoL KOOS Scores.

Variable	n missing	Value	n (%) of event	Univariable		Multivariable	
				OR (95% CI)	P value	Area under ROC-curve (95% CI)	OR (95% CI) MIC KOOS Sport and QoL
Age at time of surgery	0	16 to <22	2439 (46.9)	0.86 (0.83 to 0.88)	<0.0001	0.54 (0.53 to 0.55)	0.91 (0.88 to 0.94)
		22 to <33	2601 (46.5)				
		33 to 74	2069 (38.7)				
Sex	0	Female	3438 (43.8)	1.03 (0.96 to 1.09)	0.43	0.50 (0.50 to 0.51)	0.93 (0.87 to 0.99)
		Male	3671 (44.4)				
BMI	4490	15.1 to <23.1	1674 (43.1)	1.00 (0.99 to 1.01)	0.90	0.50 (0.49 to 0.51)	
		23.1 to <25.5	1634 (42.2)				
		25.5 to 45.4	1654 (42.6)				
Smoking	4305	No	4802 (42.5)	1.15 (0.96 to 1.37)	0.12	0.50 (0.50 to 0.51)	
		Yes	241 (45.9)				
Meniscal injury	0	No	4098 (45.3)	0.89 (0.84 to 0.95)	0.0004	0.51 (0.51 to 0.52)	0.97 (0.90 to 1.04)
		Yes	3011 (42.5)				
Cartilage injury	0	No	5219 (44.3)	0.96 (0.90 to 1.03)	0.29	0.50 (0.50 to 0.51)	1.17 (1.09 to 1.27)
		Yes	1890 (43.4)				
Collateral ligament injury (MCL/LCL)	0	No	6829 (44.1)	0.95 (0.81 to 1.11)	0.53	0.50 (0.50 to 0.50)	
		Yes	280 (42.9)				
ACL graft	528	Patellar	396 (51.6)	0.72 (0.62 to 0.83)	<0.0001	0.51 (0.50 to 0.51)	0.70 (0.60 to 0.81)
		Hamstring	6445 (43.4)				
Sports-related injury	32	No	1073 (45.1)	0.95 (0.87 to 1.04)	0.30	0.50 (0.50 to 0.51)	
		Yes	6025 (43.9)				

Continued

Table 4 Continued

Variable	n missing	Value	n (%) of event	Univariable		Multivariable		P value
				OR (95% CI)	MIC KOOS Sport and QoL	Area under ROC-curve (95% CI)	OR (95% CI) MIC KOOS Sport and QoL	
KOOS pre-operative Sport/Rec and QoL	0	0 to <25	1696 (31.7)	1.35 (1.32 to 1.37)	<0.0001	0.64 (0.64 to 0.66)	1.34 (1.31 to 1.36)	<0.0001
		25 to <45	2394 (40.9)					
		45 to 83.125	3019 (61.3)					

Univariable and multivariable logistic regression analysis was used for prediction analyses. All tests were performed with univariable logistic regression, while multivariable logistic regression models were only performed for age at time of surgery, sex, cartilage injury, graft type and KOOS preoperative Sports & QoL. Area under ROC-curve with 95% CI for multivariable model=0.65 (0.64 to 0.66). Bold font indicates significant significance.

BMI, body mass index; KOOS, Knee injury and Osteoarthritis Outcome Score; LCL, lateral collateral ligament; MCL, medial collateral ligament; MIC, minimal important change; QoL, quality of life; ROC, receiver operating characteristics; Sport & QoL, combined Sport/Rec and QoL KOOS subscales; Sport/Rec, function in sport and recreation.

be explained by the different rehabilitation strategies and psychosocial barriers related to self-efficacy compared with females, where males have been reported to maintain a more positive mindset compared with females.^{26 27} Thus, the eventual discrepancy in post-operative care between females and males might have had an impact on the post-operative outcomes 1 year after ACL-R. This discrepancy in outcomes followed by an ACL injury between males and females has previously been discussed to be potentially influenced by 'gender environment' of ACL rehabilitation, where gender has been explained to function as an extrinsic determinant over the course of treatment for ACL injury.²⁶ Furthermore, self-efficacy has previously been identified as an important predictor of the KOOS Sport/Rec and QoL scores at 1 year follow up²⁸ and subsequently, could partly explain our study results reporting better post-operative outcomes in males.

This study also found that patients undergoing ACL-R with hamstring tendon autograft had lower odds for not achieving the MIC for both the Sport/Rec and QoL KOOS subscales compared with patients undergoing patellar tendon autograft ACL-R. Less post-operative kneeling pain has been previously reported in the patients undergoing ACL-R with hamstring tendon autograft compared with patients undergoing patellar tendon autograft ACL-R²⁹⁻³¹ and thus, could partly explain the differences found in early post-operative outcomes. In addition, a relatively small proportion of patients in our cohort underwent ACL-R using patellar tendon autograft, which could have affected the study results since a small sample size may lead to overestimates of the effect size. Consequently, these findings should be interpreted with caution since use of hamstring tendon and patellar tendon autografts for ACL-R might result in similar short-term and long-term outcomes.

Poorer outcomes have previously been reported following ACL-R with concomitant cartilage injuries,^{11 12 32 33} and may partly be explained by the more advanced surgical procedures required followed by eventually more restricted post-operative rehabilitation for patients with concomitant cartilage injuries compared those without a cartilage injury.³⁴ In addition, a higher degree of persistent symptoms and pain after ACL-R have been previously reported in patients with concomitant cartilage injuries compared with patients without cartilage injuries,³⁵ which may result in reduced activity and decreased QoL.

Adding to previous studies, this study found that patients with higher pre-operative KOOS Sport/Rec and QoL scores had higher odds of not achieving an MIC for both the Sport/Rec and QoL subscales 1 year after ACL-R compared with patients with lower pre-operative KOOS Sport/Rec and QoL scores. Higher pre-operative PROs have previously been associated with an increased likelihood of better outcome scores after ACL-R.^{36 37} The higher odds of not achieving the MIC 1 year after ACL-R in patients with higher pre-operative KOOS may be explained by the fact that these patients experience less improvement in KOOS Sports/Rec and QoL scores,³⁷ because pre-operatively they already had good/better knee function and QoL. In addition, patients with lower pre-operative KOOS Sport/Rec and QoL scores in our cohort appeared to benefit more, in terms of improvement in KOOS from undergoing ACL-R compared with patients with higher pre-operative KOOS Sport/Rec and QoL scores. Furthermore, patients with higher pre-operative scores have less room for improvement and achievement of the MIC and as such, the MIC value, which was determined on all patients, may not be an appropriate MIC for patients that have higher pre-operative scores. Hence, it may be necessary to have different MIC values based on the patients baseline PRO score.

The modifiable (graft choice, concomitant cartilage injuries, preoperative KOOS scores) and non-modifiable risk factors (age, patient sex) for not achieving the MIC found in this study should be considered prior to surgery. Importantly, the results suggest that younger patients, females and patients with higher pre-operative KOOS Sport/Rec and QoL scores may benefit less from the surgical treatment compared with older patients, males and patients with lower pre-operative KOOS Sport/Rec and QoL scores. Additionally, the findings of this study suggest that treatment of concomitant cartilage injuries may be necessary to increase the likelihood of achieving a clinically important benefit from surgery. Achievement of the MIC after ACL-R may be multifactorial, and therefore, the identified possible risk factors for the failure to achieve the MIC should be considered in a treatment plan to establish realistic expectations for the patient.

This study had several strengths and limitations. One of the main strengths of this study was the large sample size including both baseline and follow-up data on 16 131 patients undergoing ACL-R. In addition, the SNKLR has previously been reported to cover more than 90% of all ACL-Rs performed in Sweden³⁵ and, therefore, has good generalisability to the Swedish population undergoing ACL-R. One limitation of this study was the relatively poor predictive capacity of the multivariable models leading to some potential limitations when interpreting the study findings. Other variables need to be explored to better understand which patients who do not benefit from ACL-R. However, the multivariable models of this study can help to correctly estimate which individuals will fail to achieve the MIC for both the Sport/Rec and QoL subscales 1 year after ACL-R in approximately 64% of cases in expectation of that the variables included in this current multivariable model are known. Additionally, data in regard to treatment of concomitant injuries was not obtained from the registry and thus, could not be analysed. The response rate for the KOOS at 1-year follow-up has also been relatively low in the SNKLR, varying between 58% and 76%.³⁸ Finally, the MIC values, used in this study, might have not identified all the patients achieving the MIC after ACL-R since the MIC may vary based on the magnitude of the baseline Sport/Rec and QoL scores.

CONCLUSION

Older patients, males and patients treated with hamstring tendon autograft ACL-R had lower odds of not achieving the MIC for both the Sport/Rec and QoL KOOS subscales 1 year after ACL-R. In addition, patients with concomitant cartilage injuries and higher pre-operative KOOS Sport/Rec and QoL scores had higher odds for not achieving the MIC 1 year after ACL-R. Our study results suggest that younger patients, females and patients with concomitant cartilage injuries and higher pre-operative KOOS Sport/Rec and QoL scores are less likely to benefit from ACL-R, and subsequently, have a lower probability of improved knee function after ACL-R. Furthermore, graft choice may also affect the risk of not achieving the MIC. However, results should be interpreted with caution as the predictive capacity of the multivariable models was relatively poor and thus, validation on a second independent sample is needed before the results can be applied clinically.

Twitter Eric Hamrin Senorski @senorski

Acknowledgements The authors would like to thank statisticians Bengt Bengtsson and Nils-Gunnar Pehrsson from the Statistiska Konsultgruppen for help

with performing the statistical analyses and giving advice regarding interpretation of the data.

Contributors All listed authors have contributed substantially to this work: literature search, and primary manuscript preparation were performed by JK, RS, VR, and TS. JJI, VM, KS and EHS assisted with study design, interpretation of the results, initial drafting of the manuscript, as well as editing and final manuscript preparation. All authors have read and approved the final manuscript to be submitted and published. EHS acted as the study guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests VM reports educational grants, consulting fees and speaking fees from Smith & Nephew plc, educational grants from Arthrex and DePuy/Synthes, is a board member of the International Society of Arthroscopy, Knee Surgery and Orthopaedic Sports Medicine (ISAKOS), and deputy editor-in-chief of Knee Surgery, Sports Traumatology, Arthroscopy (KSSTA). KS is a member of the Board of Directors in Getinge AB. JJI will become President-Elect of the Board of Directors for the Journal of Orthopaedic and Sports Physical Therapy on 27 February 2023.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study was approved by the Regional Ethical Board in Stockholm, Sweden (2011/337-31/3), and the Swedish Ethical Review Authority (2022-00913-01), and was performed in accordance with the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Janina Kaarre <http://orcid.org/0000-0003-2559-8283>

Rebecca Simonson <http://orcid.org/0000-0001-7025-7619>

Eric Hamrin Senorski <http://orcid.org/0000-0002-9340-0147>

REFERENCES

- 1 Lohmander LS, Englund PM, Dahl LL, *et al.* The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med* 2007;35:1756–69.
- 2 Kaur M, Ribeiro DC, Lamb P, *et al.* Low knee-related quality of life and persistent physical asymmetries in participants up to 10 years post-ACL reconstruction—a cross-sectional study. *Phys Ther Sport* 2021;48:35–42.
- 3 Fabricant PD, Lakomkin N, Cruz Al Jr, *et al.* ACL reconstruction in youth athletes results in an improved rate of return to athletic activity when compared with non-operative treatment: a systematic review of the literature. *Journal of ISAKOS* 2016;1:62–9.
- 4 Persson K, Bergerson E, Svantesson E, *et al.* Greater proportion of patients report an acceptable symptom state after ACL reconstruction compared with non-surgical treatment: a 10-year follow-up from the Swedish national knee ligament registry. *Br J Sports Med* 2022;56:862–9.
- 5 Hamrin Senorski E, Svantesson E, Beischer S, *et al.* Low 1-year return-to-sport rate after anterior cruciate ligament reconstruction regardless of patient and surgical factors: a prospective cohort study of 272 patients. *Am J Sports Med* 2018;46:1551–8.
- 6 Nwachukwu BU, Sullivan SW, Rauck RC, *et al.* Patient-reported outcomes and factors associated with achieving the minimal clinically important difference after ACL reconstruction: results at a mean 7.7-year follow-up. *JB JS Open Access* 2021;6:e21.00056.
- 7 Mai HT, Chun DS, Schneider AD, *et al.* Performance-based outcomes after anterior cruciate ligament reconstruction in professional athletes differ between sports. *Am J Sports Med* 2017;45:2226–32.

- 8 MARS Group, Allen CR, Anderson AF, *et al.* Surgical predictors of clinical outcomes after revision anterior cruciate ligament reconstruction. *Am J Sports Med* 2017;45:2586–94.
- 9 Snaebjörnsson T, Hamrin Senorski E, Ayeni OR, *et al.* Graft diameter as a predictor for revision anterior cruciate ligament reconstruction and KOOS and EQ-5D values: a cohort study from the Swedish national knee ligament register based on 2240 patients. *Am J Sports Med* 2017;45:2092–7.
- 10 Hamrin Senorski E, Sundemo D, Svantesson E, *et al.* Preoperative and intraoperative predictors of long-term acceptable knee function and osteoarthritis after anterior cruciate ligament reconstruction: an analysis based on 2 randomized controlled trials. *Arthroscopy* 2019;35:489–99.
- 11 Hamrin Senorski E, Svantesson E, Baldari A, *et al.* Factors that affect patient reported outcome after anterior cruciate ligament reconstruction—a systematic review of the scandinavian knee ligament registers. *Br J Sports Med* 2019;53:410–7.
- 12 Randsborg P-H, Cepeda N, Adamec D, *et al.* Patient-reported outcome, return to sport, and revision rates 7–9 years after anterior cruciate ligament reconstruction: results from a cohort of 2042 patients. *Am J Sports Med* 2022;50:423–32.
- 13 Beletsky A, Naami E, Lu Y, *et al.* The minimally clinically important difference and substantial clinical benefit in anterior cruciate ligament reconstruction: a time-to-achievement analysis. *Orthopedics* 2021;44:299–305.
- 14 Heijne A, Hagströmer M, Werner S. A two- and five-year follow-up of clinical outcome after ACL reconstruction using BPTB or hamstring tendon grafts: a prospective intervention outcome study. *Knee Surg Sports Traumatol Arthrosc* 2015;23:799–807.
- 15 Magnitskaya N, Mouton C, Gokeler A, *et al.* Younger age and hamstring tendon graft are associated with higher IKDC 2000 and KOOS scores during the first year after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2020;28:823–32.
- 16 von Elm E, Altman DG, Egger M, *et al.* The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008;61:344–9.
- 17 Kvist J, Kartus J, Karlsson J, *et al.* Results from the swedish national anterior cruciate ligament register. *Arthroscopy* 2014;30:803–10.
- 18 Hamrin Senorski E, Svantesson E, Engebretsen L, *et al.* 15 years of the Scandinavian knee ligament registries: lessons, limitations and likely prospects. *Br J Sports Med* 2019;53:1259–60.
- 19 Ingelsrud LH, Terwee CB, Terluin B, *et al.* Meaningful change scores in the knee injury and osteoarthritis outcome score in patients undergoing anterior cruciate ligament reconstruction. *Am J Sports Med* 2018;46:1120–8.
- 20 Roos EM, Lohmander LS. The knee injury and osteoarthritis outcome score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes* 2003;1:64.
- 21 Salavati M, Akhbari B, Mohammadi F, *et al.* Knee injury and osteoarthritis outcome score (KOOS): reliability and validity in competitive athletes after anterior cruciate ligament reconstruction. *Osteoarthritis Cartilage* 2011;19:406–10.
- 22 Beischer S, Hamrin Senorski E, Thomeé C, *et al.* Young athletes return too early to knee-strenuous sport, without acceptable knee function after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2018;26:1966–74.
- 23 Feucht MJ, Cotic M, Saier T, *et al.* Patient expectations of primary and revision anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2016;24:201–7.
- 24 Barenius B, Forssblad M, Engström B, *et al.* Functional recovery after anterior cruciate ligament reconstruction, a study of health-related quality of life based on the swedish national knee ligament register. *Knee Surg Sports Traumatol Arthrosc* 2013;21:914–27.
- 25 Ageberg E, Forssblad M, Herbertsson P, *et al.* Sex differences in patient-reported outcomes after anterior cruciate ligament reconstruction: data from the Swedish knee ligament register. *Am J Sports Med* 2010;38:1334–42.
- 26 Parsons JL, Coen SE, Bekker S. Anterior cruciate ligament injury: towards a gendered environmental approach. *Br J Sports Med* 2021;55:984–90.
- 27 Lisee CM, DiSanti JS, Chan M, *et al.* Gender differences in psychological responses to recovery after anterior cruciate ligament reconstruction before return to sport. *J Athl Train* 2020;55:1098–105.
- 28 Thomeé P, Währborg P, Börjesson M, *et al.* Self-efficacy of knee function as a pre-operative predictor of outcome 1 year after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2008;16:118–27.
- 29 Zhao L, Lu M, Deng M, *et al.* Outcome of bone-patellar tendon-bone vs hamstring tendon autograft for anterior cruciate ligament reconstruction: a meta-analysis of randomized controlled trials with a 5-year minimum follow-up. *Medicine (Baltimore)* 2020;99:e23476.
- 30 Mastrokalos DS, Springer J, Siebold R, *et al.* Donor site morbidity and return to the preinjury activity level after anterior cruciate ligament reconstruction using ipsilateral and contralateral patellar tendon autograft: a retrospective, nonrandomized study. *Am J Sports Med* 2005;33:85–93.
- 31 Dai W, Leng X, Wang J, *et al.* Quadriceps tendon autograft versus bone-patellar tendon-bone and hamstring tendon autografts for anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med* 2022;50:3425–39.
- 32 Hamrin Senorski E, Alentorn-Geli E, Musahl V, *et al.* Increased odds of patient-reported success at 2 years after anterior cruciate ligament reconstruction in patients without cartilage lesions: a cohort study from the Swedish national knee ligament register. *Knee Surg Sports Traumatol Arthrosc* 2018;26:1086–95.
- 33 Ulstein S, Årøen A, Engebretsen L, *et al.* Effect of concomitant cartilage lesions on patient-reported outcomes after anterior cruciate ligament reconstruction: a nationwide cohort study from norway and sweden of 8470 patients with 5-year follow-up. *Orthop J Sports Med* 2018;6:2325967118786219.
- 34 Thrush C, Porter TJ, Devitt BM. No evidence for the most appropriate postoperative rehabilitation protocol following anterior cruciate ligament reconstruction with concomitant articular cartilage lesions: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2018;26:1065–73.
- 35 Ahlén M, Samuelsson K, Sernert N, *et al.* The Swedish national anterior cruciate ligament register: a report on baseline variables and outcomes of surgery for almost 18,000 patients. *Am J Sports Med* 2012;40:2230–5.
- 36 MARS Group, Wright RW, Huston LJ, *et al.* Predictors of patient-reported outcomes at 2 years after revision anterior cruciate ligament reconstruction. *Am J Sports Med* 2019;47:2394–401.
- 37 Beletsky A, Naami E, Lu Y, *et al.* The patient acceptable symptomatic state in primary anterior cruciate ligament reconstruction: predictors of achievement. *Arthroscopy* 2021;37:600–5.
- 38 XBase The Swedish National Knee Ligament Registry. The swedish knee ligament registry annual report; 2020. Available: www.acregister.se

Supplemental material

Table 5. Achievement of the MIC for the KOOS Sport/Rec and QoL subscales.

MIC KOOS Sport/Rec (MIC KOOS Sport/Rec change)		MIC KOOS QoL (MIC KOOS QoL Change)	
Frequency	Not achieving the MIC QoL	Achieving the MIC QoL	Total
Not achieving the MIC Sport/Rec	3,340	1,619	4,959
Achieving the MIC Sport/Rec	2,150	9,022	11,172
Total	5,490	10,641	16,131
KOOS=Knee injury and osteoarthritis outcome score; MIC=minimal important change; QoL=quality of life; Sport/Rec= function in sport and recreation			

Table 6. Results from univariable and multivariable logistics regression analysis. Prediction for not achieving the MIC for Sport/Rec KOOS Score.

Univariable					Multivariable			
Variable	n	Value	n (%) of	OR (95% CI) MIC KOOS	P	Area under ROC-Curve	OR (95% CI) MIC KOOS	P
	missing		event	Sport/Rec		(95% CI)	Sport/Rec	
Age at time of surgery	0	16-<22	1,594 (30.7)					
		22-<33	1,819 (32.5)					
		33-74	1,546 (28.9)	0.95 (0.92-0.98)	0.0012	0.51 (0.50-0.52)	1.05 (1.01-1.08)	0.015
Sex	0	Female	2,438 (31.0)					
		Male	2,521 (30.5)	0.97 (0.91-1.04)	0.44	0.50 (0.49-0.51)	0.81 (0.75-0.87)	<0.0001
BMI	4,490	15.1-<23.1	1,101 (28.3)					
		23.1-<25.5	1,117 (28.8)					
		25.5-45.4	1,190 (30.7)	1.02 (1.00-1.03)	0.0092	0.51 (0.50-0.52)		
Smoking	4,305	No	3,295 (29.2)					
		Yes	169 (32.2)	1.15 (0.96-1.39)	0.14	0.50 (0.50-0.51)		
Meniscal injury	0	No	2,910 (32.2)					
		Yes	2,049 (28.9)	0.86 (0.80-0.92)	<0.0001	0.52 (0.51-0.53)	0.93 (0.86-1.00)	0.045
Cartilage injury	0	No	3,583 (30.4)					
		Yes	1,376 (31.6)	1.06 (0.98-1.14)	0.15	0.51 (0.50-0.51)	1.27 (1.17-1.36)	<0.0001

Collateral ligament injury	0	No	4,767 (30.8)					
(MCL/LCL)		Yes	192 (29.4)	0.94 (0.79-1.11)	0.45	0.50 (0.50-0.50)		
ACL graft	528	Patellar	308 (40.1)					
		Hamstring	4,458 (30.1)	0.64 (0.55-0.74)	<0.0001	0.51 (0.50-0.52)	0.59 (0.51-0.70)	<0.0001
Sports-related injury	32	No	808 (33.9)					
		Yes	4,141 (30.2)	0.84 (0.77-0.92)	0.0003	0.51 (0.51-0.52)		
KOOS pre-op Sport/Rec	0	0.0-<25.0	921 (17.4)					
		25.0-<50.0	1,280 (24.9)	1.34 (1.32-1.36)	<0.0001	0.69 (0.68-0.70)	1.36 (1.31-1.38)	<0.0001
		50.0-85.0	2,758 (48.4)					

Univariable and multivariable logistic regression analysis were used for prediction analyses. All tests were performed with univariable logistic regression, while multivariable logistic regression models were only performed for age at time of surgery, sex, cartilage injury, graft type and KOOS pre-op Sports/Rec. Area under ROC-curve with 95% CI for multivariable model = 0.70 (0.69-0.70).

BMI=body mass index; CI=confidence interval; KOOS=Knee injury and osteoarthritis outcome score; LCL=lateral collateral ligament; MCL=medial collateral ligament; MIC=minimal important change; OR=odds ratio; Pre-op=pre-operative; ROC=receiver operating characteristics; Sport/Rec=function in sport and recreation

Table 7. Results from univariable and multivariable logistics regression analysis. Prediction for not achieving the MIC for QoL KOOS Score.

Univariable				Multivariable				
Variable	n	Value	n (%) of	OR (95% CI) MIC	P	Area under ROC-Curve	OR (95% CI) MIC	P
	missing		event	KOOS QoL		(95% CI)	KOOS QoL	
Age at time of surgery	0	16-<22	1,998 (38.4)					
		22-<33	2,015 (36.0)					
		33-74	1,477 (27.7)	0.81 (0.78-0.83)	<0.0001	0.56 (0.55-0.57)	0.85 (0.82-0.88)	<0.0001
Sex	0	Female	2,617 (33.3)					
		Male	2,873 (34.7)	1.07 (1.00-1.14)	0.058	0.51 (0.50-0.52)	1.02 (0.95-1.09)	0.53
BMI	4,490	15.1-<23.1	1,293 (33.3)					
		23.1-<25.5	1,244 (32.1)					
		25.5-45.4	1,266 (32.6)	1.00 (0.99-1.01)	0.86	0.49 (0.48-0.51)		
Smoking	4,305	No	3,679 (32.6)					
		Yes	195 (37.1)	1.22 (1.02-1.47)	0.029	0.50 (0.50-0.51)		
Meniscal injury	0	No	3,159 (34.9)					
		Yes	2,331 (32.9)	0.91 (0.86-0.98)	0.0069	0.51 (0.50-0.52)	0.99 (0.92-1.06)	0.73
Cartilage injury	0	No	4,058 (34.5)					
		Yes	1,432 (32.9)	0.93 (0.87-1.00)	0.059	0.51 (0.50-0.51)	1.15 (1.06-1.25)	0.0006

Collateral ligament injury	0	No	5,283 (34.1)					
(MCL/LCL)		Yes	207 (31.7)	0.90 (0.76-1.06)	0.20	0.50 (0.50-0.51)		
ACL graft	528	Patellar	286 (37.2)					
		Hamstring	5,002 (33.7)	0.86 (0.74-1.00)	0.045	0.50 (0.50-0.51)	0.85 (0.73-1.00)	0.047
Sports-related injury	32	No	739 (33.3)					
		Yes	4,691 (34.2)	1.04 (0.95-1.14)	0.40	0.50 (0.50-0.51)		
KOOS pre-op QoL	0	0.0-<25	969 (21.7)					
		25.0-<50	2,873 (32.7)					
		50.0-81.3	1,648 (57.5)	1.42 (1.39-1.45)	<0.0001	0.65 (0.64-0.66)	1.40 (1.37-1.43)	<0.0001
Univariable and multivariable logistic regression analysis were used for prediction analyses. All tests were performed with univariable logistic regression, while multivariable logistic regression models were only performed for age at time of surgery, sex, cartilage injury, graft type and KOOS pre-op QoL. Area under ROC-curve with 95% CI for multivariable model = 0.66 (0.65-0.67).								
BMI=body mass index; CI=confidence interval; KOOS=Knee injury and osteoarthritis outcome score; LCL=lateral collateral ligament; MCL=medial collateral ligament; MIC=minimal important change; OR=odds ratio; Pre-op=pre-operative; ROC=receiver operating characteristics; Sport/Rec=function in sport and recreation								