

# Clinical and exercise professional opinion of return-to-running readiness after childbirth: an international Delphi study and consensus statement

Shefali Mathur Christopher <sup>1,2</sup>, Gráinne Donnelly,<sup>3</sup> Emma Brockwell,<sup>4</sup> Kari Bo,<sup>5,6</sup> Margie H Davenport <sup>7</sup>, Marlize De Vivo <sup>8,9</sup>, Sinead Dufour,<sup>10</sup> Lori Forner <sup>11</sup>, Hayley Mills <sup>8</sup>, Isabel S Moore <sup>3</sup>, Amanda Olson,<sup>12</sup> Rita E Deering <sup>13,14</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2023-107489>).

For numbered affiliations see end of article.

## Correspondence to

Dr Shefali Mathur Christopher, Department of Rehabilitation Sciences, Doctor of Physical Therapy, Tufts University, Seattle, Washington, USA; [Shefalichristopher@gmail.com](mailto:Shefalichristopher@gmail.com)

Accepted 21 November 2023  
Published Online First  
26 December 2023

## ABSTRACT

Female athletes have identified a lack of guidance as a barrier to successfully returning to running postpartum, and existing guidelines are vague. Our aim was to define the current practice of determining postpartum run-readiness through a consensus survey of international clinicians and exercise professionals in postpartum exercise to assist clinicians and inform sport policy changes.

A three-round Delphi approach was used to gain international consensus from clinicians and exercise professionals on run-readiness postpartum. Professionals who work with postpartum runners participated in an online survey to answer open-ended questions about the following postpartum return-to-running topics: definitions (runner and postpartum), key biopsychosocial milestones that runners need to meet, recommended screening, timeline to initiate running, support items, education topics and factors that contribute to advising against running. Consensus was defined as  $\geq 75\%$  participant agreement.

One hundred and eighteen professionals participated in round I, 107 participated in round II (response rate 90.6%) and 95 participated in round III (response rate 80.5%). Responses indicated that, following a minimum 3-week period of rest and recovery, an individualised timeline and gradual return to running progression can be considered. Screening for medical and psychological concerns, current physical capacity, and prior training history is recommended prior to a return to running. This study proposes recommendations for the initial guidance on return-to-running postpartum, framed in the context of current research and consensus from professionals. Future research is needed to strengthen and validate specific recommendations and develop guidelines for best practice when returning-to-running after childbirth.

## INTRODUCTION

Females experience key transitions across the lifespan—including puberty, pregnancy and menopause—where significant changes in hormones and body morphology may influence exercise participation and performance.<sup>1–3</sup> The perinatal period is one such transition that profoundly affects a female's physiology and biomechanics, with lasting implications that may challenge future exercise participation.<sup>4–6</sup> Running is a popular form of exercise for the perinatal population,<sup>7–10</sup> but a recent

study reported only 31% of pregnant or postpartum runners received advice on returning to running after childbirth.<sup>11</sup> As a result, runners often self-determine how to continue running during and after pregnancy, and approximately 46% of runners stop running during pregnancy and 25% do not return-to-running after childbirth.<sup>12</sup> This lack of information on safe participation in running during the perinatal period is a significant barrier to gender and sex equity in sports.<sup>13</sup>

After major surgery or injury, most athletes undergo rehabilitation before returning to sport. During rehabilitation, the athlete must meet key milestones to progress through rehabilitation stages and, at a minimum, be screened for mental and physical readiness to fully participate in sport.<sup>14–15</sup> A similar approach has been proposed for return-to-running postpartum<sup>16–19</sup> but the high-quality evidence needed to confirm and optimise these approaches is still lacking. Due to this lack of evidence, runners and the clinicians who work with them have to rely on expert opinion, which extrapolates findings from the general research on return-to-sport (ie, following an athletic injury), postpartum populations (non-athletes) and running-related injury (RRI) research.<sup>16–22</sup> While the existing frameworks have many similarities, there are some conflicting theories (ie, timeline for return). Therefore, the current study employed a Delphi technique to determine consensus from many experienced clinical and exercise professionals on current practice of determining run-readiness after childbirth. Expert opinion consensus on the rehabilitation programme and running programme design is presented in another publication.<sup>23</sup>

## METHODS

The Delphi technique (three rounds)—which is commonly used for decision making and forecasting studies—was used to determine consensus of clinical and exercise professionals on postpartum return-to-run topics.<sup>24–31</sup> Experienced professionals (respondent group) were asked their opinion on key musculoskeletal assessments, milestones and screening that should be used when determining run-readiness postpartum.


## Participants

The respondent group were experienced professionals recruited through personal networks, social



© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Christopher SM, Donnelly G, Brockwell E, et al. *Br J Sports Med* 2024;**58**:299–312.



**Figure 1** Methods.

media (ie, Twitter, Instagram and Facebook) and word-of-mouth via a purposeful and snowball sampling approach. All prospective participants completed an online recruitment survey in which they reported demographic information, profession, number of years working with postpartum runners, and percentage of caseload consisting of postpartum runners. From this online recruitment survey, respondents were eligible to participate (ie, considered experienced professionals) if they were health, rehabilitation and/or fitness professionals with either (a)  $\geq 5$  years' experience treating postpartum runners OR (b) if  $< 5$  years' experience, their caseload is primarily postpartum runners ( $\geq 50\%$ ).

The workgroup (authors) consisted of investigators that had an average of 10 years of experience working with perinatal runners and represented a variety of disciplines (exercise physiology, biomechanics, psychology and physiotherapy). All authors reviewed the Delphi results and current literature, then participated in a discussion to finalise recommendations.

### Instrument development and piloting

All authors contributed to the development of a pilot survey consisting of open-ended questions with free-text responses (figure 1). Eleven practitioners, who were either retired professionals in the field or were no longer working with this population, were identified by the authors as pilot participants for round 1 of the survey. Pilot participants provided feedback (eg, question clarity), and necessary changes were made before distribution of round I of the survey to study participants. These data were separate from the Delphi survey and used only for development and piloting round I of the survey.

### Procedure

A narrative literature review on postpartum physiology/biomechanics, running, RRI and existing run-readiness frameworks

(including grey literature) informed the questions chosen for round I of the survey (online supplemental appendix A). For each round, Qualtrics (Seattle, USA) distributed surveys via a personalised email link. Informed consent was obtained prior to entering the survey questions. The definition of 'consensus' was established a priori as 75% and it was decided to limit voting to three rounds for participant retention.<sup>32</sup> All identified experienced practitioners from the recruitment survey were sent a link to the round I survey. All participants who completed round I were sent the survey for rounds II and III. Each round was live for 3–4 weeks with weekly email reminders sent to respondents who had not completed the survey. Four authors (SMC, MHD, SD and RED) with experience in Delphi studies or similar mixed-methods research undertook thematic coding of the survey free text responses in rounds I and II.

After completion of all rounds of the Delphi survey, all authors contributed to an additional literature search to summarise the current scientific evidence and determine if respondent consensus was in line with current research. Search topics were determined by the themes identified by respondents and a narrative review was conducted. Due to limited evidence in the postpartum running population, searches were not limited to postpartum running-related literature or to systematic reviews or randomised controlled trials. When appropriate, grey literature was included. The level of evidence for each topic, based on the Sackett scale of scientific evidence (figure 2), is provided at the end of each evidence summary section.<sup>33</sup> A table indicating the level of evidence for each article cited is provided in supplemental digital content.

### Round I survey

The first round included demographic questions about the respondents. There were also five open-ended questions about screening for run-readiness and three open-ended questions

Level of Evidence	Description of evidence
I	Meta-analyses; systematic reviews; large, high-quality RCTs
II	Small or low-quality RCTs
III	Non-randomized trials; observational studies (including survey studies); cohort studies
IV	Non-randomized trials with historical controls
V	Case series; expert opinion; textbooks; formal consensus; narrative reviews

**Figure 2** Levels of evidence. RCTs, randomised controlled trial.

about return-to-running considerations (key milestones, factors to stop running, items that can aid running). In addition, respondents were asked to define ‘postpartum’ and ‘runner’ (online supplemental appendix A).

### Round II survey

Thematic coding of round I responses led to the development of the round II survey, which was primarily statements with Likert-scale choices (strongly agree, agree, disagree, strongly disagree).

### Round III survey

Round III of the survey was designed to establish consensus on the Likert-scale statements from round II. According to Delphi methodology, the same survey questions from round II were presented to the participants with the addition of graphs representing participant responses from round II (percentage of votes for strongly agree, agree, disagree and strongly disagree) in lieu of in-person discussion.<sup>31</sup> Participants were again asked to choose their level of agreement (as per round II) with each statement.

### Author recommendations

After reviewing the survey results and completing a narrative literature review, recommendations were proposed based on author discussion and synthesis of the Delphi data and current evidence. An anonymous survey was then sent out to all authors to determine author consensus on the recommendations. Authors completed three rounds of voting: Vote 1 consisted of the original recommendation for each section from the group meeting along with free-text options to indicate dissenting opinions. Vote 2 presented all author-suggested recommendations for each section. Vote 3 again presented all author-suggested recommendations along with the results of round 2 voting.

### Diversity, equity and inclusion statement

The all-female author group, representing five countries across three continents, were primarily Caucasian with one woman of colour. Experienced practitioners (respondent group) were included based on number of years working with postpartum runners and thus junior, mid-career and senior level practitioners from a variety of professional backgrounds were included. Only two men participated in the Delphi survey as respondents. In discussing generalisability of our results and limitations in our findings, we recognise that these results may exclude professionals of a low socioeconomic status, where advanced education is unavailable, or from marginalised communities as perinatal care is not part of basic training in many professions. While efforts to recruit diverse respondents with sociocultural differences were made (through personal networks, social media (ie, Twitter, Instagram and Facebook) and word-of-mouth), the

recommendations made in this consensus statement may not be reflective of every culture.

## RESULTS

Two hundred and twenty-two professionals met the inclusion criteria and were sent the link for round I. 118 participants completed round I. Those 118 participants were sent invitations to complete rounds II and III. 107 completed round II and 95 completed round III. Participants had an average of 8.9 (range 2–37) years’ experience working with postpartum runners and represented seven different professions, 12 countries and 4 continents (North America, Europe, Australia and Africa). Most of the participating professionals identified as women (97%) (table 1).

### Definitions of ‘runner’ and ‘postpartum’

#### Consensus

Consensus was reached that ‘runner’ was defined as ‘anyone who runs, regardless of frequency or mileage’ (90.6%) and/or ‘anyone who self-identifies as a runner’ (92.9%). No true consensus was reached on the definition of ‘postpartum’, though respondents agreed (78.8%) that it does not refer only to the first 12 weeks after childbirth.

#### Current evidence

Various definitions of ‘runner’ exist. Some studies identify runners by a certain number of miles per week.<sup>34</sup> Experience level is usually reported (eg, novice, competitive), but standard terminology has not been used, meaning different terms may be used to describe similar cohorts (eg, beginner and novice).<sup>35</sup> The definition of ‘postpartum’ also varies, focusing on length of time since giving birth (eg, 12 weeks to 2 years).<sup>36–38</sup> The consensus that ‘postpartum’ does not refer to only the first 12 weeks after childbirth is supported by several studies using time frames > 12 weeks to define their postpartum population<sup>7–9 11 12 38–43</sup> and by evidence that postpartum mental health symptoms can still be present up to 3 years postpartum.<sup>44</sup> The inconsistencies in the literature of how long the postpartum phase persists appear to be reflected in several time frames being identified by respondents in free-text responses and inability to reach consensus on one specific time frame.

(No summary of the level of evidence is provided, as consistent definitions are non-existent.)

#### Recommendation (12/12 authors assent)

This Delphi recommends that someone who self-identifies as a runner should be evaluated and treated as one, regardless of mileage, frequency or skill level. Due to the lack of longitudinal evidence investigating perinatal runners, an individualised

**Table 1** Participant demographics

	Round 1	Round 2	Round 3
Total no of surveys started (n)	144	108	96
Total no of surveys completed (n)	118	107	95
Physical therapist/physiotherapist	96	88	80
Occupational therapist	1	1	1
Personal trainer	8	7	6
Chiropractor	1	1	0
Exercise physiologist	5	4	4
Physician	5	4	3
Run coach	1	2	1
Completion rate (%)	53	91	81
Years in current profession (n)			
0–4 years	10	10	8
5–9 years	27	24	22
10–14 years	36	31	28
15–19 years	20	18	15
20+ years	25	24	22
Years working with postpartum runners (years)			
Mean	8.85	8.99	8.93
Range	1–30	1–30	1–30
Percentage of caseload consisting of postpartum runners (n)			
0%–24%	65	57	52
25%–49%	37	35	31
50%–74%	15	14	11
75%–100%	1	1	1
Gender identity of respondents (n)			
Woman	116	105	93
Man	2	2	2
Age (years)			
Mean	38.9	39.0	39.2
Range	23–63	23–63	23–63
Race/ethnicity of respondents (n)			
White	114	103	92
Black/African American	2	2	1
Asian	3	3	3
Other	1	1	1
Respondents who identify as a runner (n)			
Yes	86	79	70
No	32	28	25
Have the respondents themselves given birth? (n)			
Yes	65	60	51
No	21	19	19
Preferred not to answer	32	28	25
Trained in internal pelvic floor muscle assessment? (n)			
Yes		72	75
No, refers to pelvic floor trained provider		20	20
No, relies on symptom reports from patient		6	0
No response		9	0

approach should be taken to determine if the runner is still recovering from pregnancy-related and childbirth-related changes or not. For example, if someone is returning to running at 2 years postpartum, they should still be evaluated or screened for postpartum run-readiness, as pregnancy and childbirth-related impairments may still be present.

**Table 2** Key milestones to assess for return to running and suggested metrics for meeting milestones

Key milestones to assess for return to running	Agree/strongly agree in round II (%)	Agree/strongly agree in round III (%)
Pelvic floor strength	<b>91.5</b>	<b>95.3</b>
Pelvic floor endurance	<b>89.4</b>	<b>94.1</b>
Pelvic floor coordination	<b>95.7</b>	<b>98.8</b>
Pelvic organ prolapse	<b>93.6</b>	<b>97.7</b>
Urinary incontinence	<b>97.8</b>	<b>97.7</b>
Anal incontinence	<b>97.9</b>	<b>97.7</b>
Lumbopelvic strength	<b>96.8</b>	<b>95.3</b>
Lower extremity strength	<b>95</b>	<b>98.8</b>
Inter-recti distance	62.8	55.3
Balance/ proprioception	<b>93.6</b>	<b>95.3</b>
Gait analysis	<b>75.5</b>	<b>78.8</b>

Bold text indicates meets consensus (>75%).

### Key milestones that need to be addressed before postpartum return-to-running

#### Consensus

From round I, 11 themes were identified as key milestones that need to be addressed before return-to-running (table 2), including: pelvic floor muscle (PFM) strength, endurance and coordination; symptoms of urinary incontinence (UI); symptoms of anal incontinence (AI); symptoms of pelvic organ prolapse (POP); lumbopelvic strength; inter-recti distance (IRD); balance and proprioception; lower extremity strength; and running gait analysis. Ten milestones met consensus, with IRD being the only milestone that did not. To note, specific cut-offs or benchmarks were not identified; rather, respondents identified key areas for evaluation.

#### Current evidence

Symptoms of pelvic floor dysfunction (PFD) are widely reported in nulliparous and parous female runners,<sup>6 9 45–59</sup> and pregnancy and childbirth increase the general population risk of PFD.<sup>60</sup> Reported frequency of UI in postpartum runners ranges from 8% to 57%,<sup>8 9 59</sup> AI was reported in 39% of postpartum runners and 19% reported symptoms of POP.<sup>59</sup> However, no studies have identified specific PFM function (strength, endurance, coordination) parameters that indicate definitive resolution and/or prevention of PFD symptoms in runners.<sup>61–63</sup> There is, however, strong evidence in the general postpartum population that PFM training is effective for treating PFD.<sup>64</sup>

Lower extremity strength has only been investigated in a small cohort of postpartum runners (N=9), which showed significantly lower hip abduction and adduction strength compared with nulliparous controls.<sup>43</sup> When considering the general running population, systematic reviews have reported that musculoskeletal measures (eg, strength) and biomechanical measures (eg, kinematics) are not stand-alone risk factors for RRI.<sup>65</sup>

Current literature on IRD has reported correlations with abdominal muscle strength and fatigability,<sup>39 41 42 66</sup> abdominal pain and quality of life,<sup>67</sup> and no correlation between IRD and low back pain, pelvic girdle pain or UI.<sup>67 68</sup> Increased IRD can also lead to fear-avoidance behaviours, which may be a barrier to return-to-exercise and running.<sup>69 70</sup> One small study showed decreased IRD with exercise in postpartum runners.<sup>38</sup> While there is insufficient evidence to support reduction in IRD with exercise training,<sup>71</sup> abdominal muscle training can influence muscular strength and endurance,<sup>72 73</sup> both of which are shown

**Table 3** Consensus on load and impact screen for return to running

Screening activity	Agree/strongly agree in round II (%)	Agree/strongly agree in round III (%)
Walking for 30 min	<b>97.9</b>	<b>97.7</b>
Single leg balance for 10s each leg	<b>89.4</b>	<b>92.9</b>
Single leg squats×10 repetitions each leg	<b>86.2</b>	<b>89.4</b>
Jogging on the spot for 1 min	<b>92.6</b>	<b>98.8</b>
Forward bounds×10 repetitions	<b>79.8</b>	<b>87.1</b>
Hopping in place×10 repetitions each leg	<b>92.6</b>	<b>95.3</b>
Single leg 'running man' (opposite arm and hip flexion/extension with knee bent)×10 repetitions each side	<b>85.1</b>	<b>84.7</b>
Calf raises×20 repetitions	<b>91.5</b>	<b>90.6</b>
Single leg bridge×20 repetitions each leg	<b>86</b>	<b>87.1</b>
Single leg sit to stand×20 repetitions each leg	<b>76.3</b>	<b>80</b>
Load and impact screening activities should be performed without exacerbation of musculoskeletal or pelvic health symptoms. Bold text indicates meets consensus (>75%).		

to be impaired in the general postpartum population and in postpartum females with diastasis recti abdominus (DRA).<sup>39 41 42 66</sup>

An initial biomechanical investigation in a small cohort of postpartum runners showed no difference in kinematic and kinetic (except braking loading rate) measures in postpartum running gait when compared with nulliparous controls.<sup>43</sup> Lastly, literature on balance and proprioception is non-existent in the postpartum running population. In the general perinatal population, evidence on changes in static balance is conflicting, with some reporting increased postural sway and others reporting no changes.<sup>74–76</sup> Expert opinions on rehabilitation of postpartum runners have included exercises to improve balance and proprioception.<sup>16 18</sup> Balance and proprioception are recommended assessments for run-readiness following knee and ankle injuries in the general population.<sup>77–80</sup> (Level of evidence: III)

### Recommendation (12/12 authors assent)

As incontinence and prolapse symptoms are well documented in both nulliparous and postpartum female runners, as well as in the general postpartum female population, a postpartum runner should ideally be evaluated for these pelvic health-related symptoms prior to initiating running. Runners with PFD should be referred to an appropriate and specialised professional. As RRI and pain are multifactorial, it is recommended to include PFM, lower extremity and lumbopelvic strength as well as balance assessments in the physical examination to aid successful return-to-running; however, due to lack of evidence, no recommendation can be made on PFD (eg, prolapse, incontinence) severity scores, objective strength or balance measurement minimums that would indicate return-to-running readiness. While IRD did not reach consensus as a milestone, runners with abdominal pain or who exhibit fear avoidance behaviours may benefit from assessment.

### Load and impact screening

#### Consensus

Consensus was reached in both rounds that a runner should be able to complete the screening tasks in [table 3](#) without musculoskeletal or pelvic health symptoms before initiating running.

#### Current evidence

No evidence exists assessing which load and impact screening tasks are ideal for identifying postpartum run-readiness. Several expert opinions recommend being able to walk for 30 min without eliciting/exacerbating cardiorespiratory, pelvic health or other musculoskeletal symptoms prior to engaging in running postpartum.<sup>16–18 20</sup> Two screens have been proposed to evaluate run-readiness, one specifically for postpartum runners. The Running Readiness Scale, which consists of five tasks (hopping, planks, step-ups, single leg squats and wall sits), was proposed to identify injury risk due to movement patterns. An initial study of this scale, validating it against three-dimensional (3D) running biomechanics in asymptomatic novice runners, showed reliability

**Table 4** Support items for return to running

Support items	Agree/strongly agree in Round II (%)	Agree/strongly agree in Round III (%)
A runner requiring support items (such as sacroiliac joint belts, taping, compression shorts, etc) is a reason to recommend that someone not resume/participate in/continue running postpartum.	3.2	1.2
Runners should be educated on appropriate breast support before returning to running after childbirth.	<b>96.8</b>	<b>97.7</b>
Footwear should be assessed for proper fit and compatibility with running goals and current musculoskeletal profile before returning to running after childbirth.	<b>92.5</b>	<b>96.5</b>
Compression garments (eg, compression shorts/leggings that go over the abdomen, compression socks) can be helpful for some postpartum runners.	<b>95.7</b>	<b>96.5</b>
Intravaginal support items (eg, pessary, Poise Impressa, tampons, menstrual cups, etc) can be helpful for postpartum runners with prolapse symptoms.	<b>97.9</b>	<b>100</b>
Intravaginal support items (eg, pessary, Poise Impressa, tampons, menstrual cups, etc) can be helpful for postpartum runners with incontinence symptoms.	<b>95.7</b>	<b>100</b>
Abdominal and/or low back taping techniques can be helpful for some runners.	<b>83.9</b>	<b>91.8</b>
Sacroiliac joint belts can be helpful for some runners.	21.5	7.1
Abdominal braces can be helpful for some runners	63	71
If a runner plans to run with their child, the stroller/pram/buggy that they intend to use should be assessed for appropriateness.	<b>89.4</b>	<b>94.1</b>
Incontinence products (eg, pads, incontinence underwear, etc) can be helpful for some runners.	<b>92.6</b>	<b>98.8</b>
Runners should not be encouraged to utilise support items (such as sacroiliac joint belts, taping, compression shorts, etc); rather, they should be encouraged to build functional strength so that these items are not necessary.	50.5	42.3
Bold text indicates meets consensus (>75%).		

**Table 5** Consensus on when to advise against running

Themes for recommending abstaining from running	Agree/strongly agree (%) Round II	Agree/strongly agree (%) Round III
<b>Pelvic health</b>		
One cannot return to running with symptoms of pelvic organ prolapse	12.7	1.2
One can return to running with mild symptoms of pelvic organ prolapse	<b>88</b>	<b>100</b>
Presence of severe/significant POP is a reason to recommend that someone not resume/participate in/continue running postpartum	69.9	<b>80</b>
One cannot return to running with symptoms of urinary incontinence	10.3	2.4
One can return to running with mild symptoms of urinary incontinence	<b>92.1</b>	<b>98.8</b>
Presence of severe/significant urinary incontinence is a reason to recommend that someone not resume/participate in/continue running postpartum	67.7	<b>75.3</b>
One cannot return to running with symptoms of anal incontinence	26.6	5.9
One can return to running with mild symptoms of anal incontinence	<b>81.7</b>	<b>96.5</b>
Presence of severe/significant urinary incontinence is a reason to recommend that someone not resume/participate in/continue running postpartum	71	<b>82.4</b>
Presence of severe/significant structural pelvic floor muscle injury (eg, levator ani avulsion, anal sphincter injury, etc) is a reason to recommend that someone not resume/participate in/continue running postpartum.	64.5	68.2
Presence of lochia (postbirth vaginal bleeding) is a reason to recommend that someone not resume/participate in/continue running postpartum.	<b>86.2</b>	<b>92.9</b>
Presence of birth complications/delayed recovery from childbirth is a reason to recommend that someone not resume/participate in/continue running postpartum.	71	<b>80</b>
<b>Musculoskeletal</b>		
Presence of musculoskeletal injuries is a reason to recommend that someone not resume/participate in/continue running postpartum.	61.3	50.6
Presence of consistent musculoskeletal pain is a reason to recommend that someone not resume/participate in/continue running postpartum.	59.1	50.6
Inter-recti distance of three finger widths or more without doming is a reason to recommend that someone not resume/participate in/continue running postpartum.	0	3.5
Inter-recti distance of three finger widths or more with doming is a reason to recommend that someone not resume/participate in/continue running postpartum.	41.5	28.2
Presence of diastasis recti abdominis with a hernia is a reason to recommend that someone not resume/participate in/continue running postpartum.	44.6	40
Poor biomechanics with day-to-day mobility (walking, stair negotiation, squats, etc) is a reason to recommend that someone not resume/participate in/continue running postpartum.	57.5	56.5
Poor bone health is a reason to recommend that someone not resume/participate in/continue running postpartum.	39.8	27.1
A runner requiring support items (such as sacroiliac joint belts, taping, compression shorts, etc) is a reason to recommend that someone not resume/participate in/continue running postpartum.	3.2	1.2
<b>Biopsychosocial</b>		
Poor sleep habits (less than 6 hours accumulated sleep/night; no stretches of sleep longer than 4 hours; etc) are a reason to recommend that someone not resume/participate in/continue running postpartum.	44.7	31.8
Poor mental health status that may be worsened by running is a reason to recommend that someone not resume/participate in/continue running postpartum.	<b>83</b>	<b>90.6</b>
High risk for REDs (ie, poor nutritional intake, history of disordered eating, rapid and drastic weight loss, etc.) is a reason to recommend that someone not resume/participate in/continue running postpartum.	<b>86</b>	<b>89.4</b>
<b>Other</b>		
Runners with pre-existing medical conditions (ie, present before pregnancy) should receive medical clearance before returning to running.	<b>85.9</b>	<b>91.8</b>
Runners who wish to run despite symptoms should not be told that they cannot run; rather, running habits may need to be modified (eg, decrease mileage) while the runner is treated for identified impairments.	<b>94.7</b>	<b>100</b>

Bold text indicates meets consensus (>75%).

and validity with the screen and knee abduction angles.<sup>81</sup> Goom *et al*<sup>19</sup> proposed that a postpartum runner should be able to walk (30 min), and perform exercises (single leg balance, single leg squats, jog, perform forward bounds, hops and single leg running man) to evaluate postpartum load and impact management in regard to provocation of pelvic floor symptoms or pain. To our knowledge, this screen has not been further investigated. A recent study of common running drills in healthy runners included three of the screening tasks (hopping in place (jump rope), jogging on the spot and forward bounds) had 76%, 87% and 104% of the vertical reaction forces of fast running, respectively, indicating that these tasks may closely mimic loads associated with running. Therefore, these drills could be used to screen or progress asymptomatic or symptomatic runners (pain,

incontinence, etc) as high impact activities have been associated with incontinence in parous and nulligravid females.<sup>50 82–84</sup> (Level of evidence: V)

**Recommendation (12/12 authors assent)**

While no studies have examined the influence of ground reaction forces on symptoms in the postpartum runner, high-impact activities have been associated with incontinence in both nulligravid and parous females. As such, it is recommended that, prior to initiating running after childbirth, a series of gradual and progressive load and impact challenges be administered to assess provocation or exacerbation of symptoms.

**Table 6** Consensus On Timeline for Return to Running.

Theme for return-to-run timeline	agree/Strongly Agree (%)Round II	agree/Strongly Agree (%)Round III
One cannot return to running before 3 weeks postpartum	<b>76.1</b>	<b>85.5</b>
One cannot return to running before 6 weeks postpartum	59.8	55.3
One cannot return to running before 12 weeks postpartum	33	20
Timeline post-birth to return to running is person-specific	<b>94.7</b>	<b>100</b>
Any acute birth injuries (such as perineal tearing, episiotomy, caesarean incision, etc.) should be completely healed before returning to running	<b>97.9</b>	<b>97.6</b>
Bold text indicates meets consensus (>75%).		

### Screening for biopsychosocial milestones

#### Consensus

Unanimous consensus was reached that it is important to assess sleep quality and habits, screen for pre-existing conditions (ie, musculoskeletal or pelvic floor symptoms) and evaluate mental health and fatigue when determining postpartum run-readiness. The importance of screening for energy availability (EA)/relative energy deficiency in sport (97.7%); whether milk supply has been sufficiently established (if desired) (98.8%) and hydration status (98.8%) also reached consensus in both rounds.

#### Current evidence

Several qualitative studies and expert opinions on readiness for return-to-running have highlighted the need to screen biopsychosocial factors.<sup>7,8,16–19,21,23,45,85</sup> Lack of sleep and a high level of fatigue have been identified as risk factors for pain in postpartum runners.<sup>7</sup> As low EA affects up to 47% of female athletes, several experts on postpartum running have stressed the importance of evaluating this.<sup>16,17,21,86,87</sup> While the difficulties of lactation have not been directly measured in runners, athletes have reported difficulties with breast feeding, supply and training schedules.<sup>13</sup> Experts have also stressed the importance of lactation consultants when working with athletes returning to sports.<sup>20,88</sup> Lastly, per a systematic review in 2019, postpartum depression is common after childbirth (up to 20%)<sup>89</sup>; however, no studies have assessed this in postpartum athletes.<sup>6,90</sup> Due to these biopsychosocial concerns, experts are recommending that the postpartum runner have access to a multidisciplinary team of providers to aid with a successful return to running.<sup>8,16</sup> (Level of evidence: III)

#### Recommendation (12/12 authors assent)

Based on consensus from experienced professionals working with postpartum runners, as well as current evidence in the general athletic population, it is recommended that runners be screened for concerns or issues with sleep, pre-existing conditions, lactation concerns, hydration, fatigue and mental health. When possible, an appropriate multidisciplinary team, consisting of a variety of healthcare professionals with expertise in the presenting concerns (eg, primary care providers, lactation consultants, pelvic health physiotherapists (PTs), mental health providers, psychiatrists, orthopaedic specialists, obstetricians/gynaecologists, urogynaecologists), should work with the runner to address these issues.

### Support items/adjuncts for return to running

#### Consensus

A unanimous consensus was reached that intravaginal support devices (eg, vaginal pessaries) can be helpful for prolapse and incontinence symptoms. Respondents agreed that runners should be educated on proper breast support (97.7%), that footwear should be assessed for fit and compatibility with running goals and current musculoskeletal profile (96.5%), and that runners who plan to run with a stroller have it assessed for appropriateness (94.1%). Respondents disagreed (92.9%) that sacroiliac joint (SIJ) belts can be helpful for some runners, and no consensus was reached on the utility of abdominal braces (71% agreed abdominal braces can be helpful for some runners) (table 4).

#### Current evidence

There is limited evidence on use of vaginal support pessaries in the postpartum period. Pessaries in addition to PFM training may improve POP symptoms<sup>91</sup> and may help with UI.<sup>92</sup> However, not all females will be candidates for pessary use, those who are may not have success with use, and intravaginal devices may not be as effective as PFM training.<sup>93</sup> The Society of Obstetricians and Gynaecologists of Canada recommends that intravaginal devices be used on an individualised basis and are considered as a first-line option for UI with high-impact exercises or when there are barriers in accessing supervised PFM training.<sup>93</sup> Such devices also promote empowerment and self-management.<sup>93</sup> No studies have been conducted on the use of absorbent items in postpartum runners. Women who exercise and experience stress UI (SUI) do report use of liners or pads to manage symptoms.<sup>94–96</sup>

The breasts can experience high magnitudes of 3D motion during running.<sup>97–99</sup> The amount, and the perceived impact, of breast motion is also influenced by individual breast size.<sup>100</sup> Motion-related breast pain has been reported in up to 40% of athletes and can negatively impact performance.<sup>101</sup> Adequate breast support is considered particularly important perinatally to accommodate breast shape and size changes, especially if lactating, as breast size can increase by 1 or more cup sizes during pregnancy.<sup>102</sup> In the general population, poor breast support is also a barrier to physical activity.<sup>103</sup> An individually fitted sports bra has been shown to reduce motion-related breast pain<sup>97</sup> while improving running economy and performance.<sup>104</sup>

Stroller running is associated with increased energy cost compared with running independently.<sup>105,106</sup> A two-handed approach to stroller running may change trunk, pelvis and hip kinematics<sup>107</sup> but spatiotemporal factors have been shown to be similar compared with independent running.<sup>105</sup>

There is no current evidence on compression garment use or the use of SIJ belts in postpartum runners. Compression garments targeting the lumbopelvic region are reported to reduce perceived symptoms of pain,<sup>108,109</sup> incontinence<sup>110,111</sup> and POI.<sup>112</sup> One study in the general postpartum population found that SIJ belts were helpful in reducing pelvic girdle pain during performance of the Active Straight Leg Raise Test.<sup>113</sup> In the general population with lumbopelvic pain, the effectiveness of SIJ belts is inconclusive and often described as having person-specific results.<sup>114–117</sup> There are also no data on footwear and postpartum runners. Experts have recommended evaluation of a postpartum runner's footwear due to potential pregnancy-related changes and incidence of running-related pain, especially in the lower extremity.<sup>16</sup>

There are currently no data examining taping (abdominal, low back, etc) for postpartum runners. The only studies examining the effect of taping in postpartum populations relate to DRA,

with no implications for running.<sup>118 119</sup> In the general running population, only lower extremity taping has been studied and there is conflicting evidence on whether it provides benefits for pain or performance.<sup>120–123</sup> (Level of evidence III)

**Recommendations (12/12 authors assent)**

Despite low-level evidence in postpartum populations, support items may be beneficial for symptom management in postpartum runners. If an intravaginal support or other continence device is desired by a postpartum runner, a collaborative pelvic healthcare team should assess the runner to determine appropriateness. Absorbent products can also be used, but runners should be encouraged to seek treatment for incontinence. Runners may benefit from a professionally guided, individualised bra fitting to select bras to suit the breast size and type of activity of the postpartum runner. Due to pregnancy-related changes, footwear should also be evaluated. Postpartum runners should be educated on considerations with stroller running for both mother and baby, and that a two-handed approach to stroller running may be favourable. Compression garments may be appropriate adjuncts to active rehabilitation in runners with lumbopelvic and/or PFD symptoms. No recommendation can be made on taping.

**Other considerations for readiness to return-to-running after childbirth**

**Consensus**

Respondents unanimously agreed that it is important to consider prior running habits—both during pregnancy and prepregnancy—as well as current training and performance goals when considering run-readiness postpartum. Respondents also agreed that it is important: (1) to prioritise the runner’s role in shared decision-making (100%); (2) to honour the runner’s wishes about when to return-to-running, even if ideal milestones have not been met (100%); (3) to consider the runner’s stress level when determining run-readiness (100%); (4) to include a multidisciplinary care team (97.7%); (5) to assess breathing technique prior to initiating running (84.7%) and (6) to consider the runner’s social support when determining run-readiness (98.8%). **Table 5** outlines additional considerations for recommending that a postpartum runner not participate in running, such as significant POP (80% agreement).

**Current evidence**

Several studies have highlighted the importance of shared decision making for patient-centred care.<sup>124 125</sup> Expert opinion encourages consideration of goals for postpartum return-to-running and highlights a multi-disciplinary approach.<sup>16 20 21</sup> Two reviews reported lack of social support as a barrier to postpartum exercise.<sup>126 127</sup>

There is no evidence in postpartum runners on the influence of returning-to-running on PFD symptoms. The American College of Obstetricians and Gynecologists (ACOG) states that symptomatic POP should be further assessed and treated.<sup>128</sup> Lochia can be present under normal circumstances for up to 8 weeks postpartum.<sup>129</sup> Persistence of vaginal bleeding (stage 1 lochia) beyond 2 weeks postpartum is likely indicative of significant pathology,<sup>129</sup> thus medical treatment should be sought and return-to-running should be delayed in this circumstance.

Pain is common in the general running population<sup>130</sup> and among postpartum runners.<sup>7 8 12 45</sup> Some causes of pain in runners (eg, bone stress injuries, medial tibial stress syndrome, etc.) will require a period of rest from running, but other causes of pain (eg, patellofemoral syndrome, etc.) do not have evidence supporting termination of running.<sup>131</sup> No evidence currently

exists on treatment of pain (ie, period of rest) in postpartum runners.

There is currently no scientific evidence that there is a relationship among DRA/abdominal wall integrity, diaphragm mechanics and breathing technique. Breathwork has been shown to not influence PFM function.<sup>132</sup> (Level of evidence: V)

**Recommendation (12/12 authors assent)**

The runner should play an active role in the plan-of-care and decision making. The runner’s medical and social history, training and goals should be considered when determining run-readiness. Runners with pain should be evaluated to determine the cause of pain, which will determine whether running is appropriate or not. Significant pelvic health symptoms should be assessed by a specialist (eg, a urogynaecologist) and may take priority over return-to-running in runners who are open to delaying running. A multidisciplinary team is encouraged to identify biopsychosocial red flags to return-to-running. It is important to identify and address barriers when designing the plan of care and return-to-exercise. No literature exists related to breathing mechanics and outcomes for perinatal runners; as such, no expert recommendation can be made.

**Education topics for postpartum runners**

**Consensus**

Respondents unanimously agreed that it is important to educate postpartum runners on (1) postpartum physiological and musculoskeletal recovery and (2) a gradual return-to-running after childbirth. Respondents also agreed that it is important to educate runners on the key milestones that indicate run-readiness (98.8%), that hydration and nutrition recommendations should be different for postpartum runners than for runners who are not postpartum (96.5%), and that runners who are lactating should be advised to express milk prior to going for a run (88.2%).

**Current evidence**

To support continued running during pregnancy, which increases the likelihood of returning to running postpartum,<sup>12</sup> education needs to be specific to running (ie, not general physical activity).<sup>11</sup> The majority of postpartum runners prefer information disseminated via websites and pelvic health PTs.<sup>12</sup> A gradual return to exercise, including running, has been recommended by several expert opinions<sup>6 16–18 20 21</sup> and is supported by RRI evidence suggesting that rapid increases in mileage or intensity increase risk.<sup>133–135</sup> Further information on gradual progression of exercise and running is presented in a companion paper.<sup>23</sup> As novice postpartum runners have higher odds of postpartum pain and up to 84% of postpartum runners have running-related pain across several body regions with the lower limbs being the most common site of pain,<sup>7 8</sup> educating runners on run readiness and how to return to running may be a priority.

No studies have investigated the relationship between breastfeeding and running. Milk secretion in the general postpartum population is associated with 700ML per day of water loss at 8 weeks postpartum,<sup>136 137</sup> which may lead to dehydration and negatively affect maternal health and exercise performance. Energy needs are also increased while lactating, with a suggested increase of ~500 kcal/day above prepregnancy caloric intake.<sup>138 139</sup> Further discussion of lactation and exercise is presented in a companion paper.<sup>23</sup> (Level of evidence: V)


**Recommendation (12/12 authors assent)**

Perinatal runners should be provided with running-specific education, during and after pregnancy, that is, individualised to their



# RETURN TO RUNNING READINESS AFTER CHILDBIRTH: KEY RECOMMENDATIONS

CHRISTOPHER SM, DONNELLY G, BROCKWELL E, BO K, DAVENPORT M, DEVIVO M, DUFOUR S, FORNER L, MILLS H, MOORE J, OLSON A, DEERING DE. CLINICAL & EXERCISE PROFESSIONAL OPINION OF RETURN-TO-RUNNING READINESS AFTER CHILDBIRTH: AN INTERNATIONAL DELPHI CONSENSUS STATEMENT. BRITISH JOURNAL OF SPORTS MEDICINE



**Figure 3** Return-to-running readiness after childbirth: infographic.

training level and goals. Educating postpartum runners on nutrition and hydration should also be a priority.

## Timeline for returning to running

### Consensus

Five themes were identified in round I for timing of return-to-running and are represented in table 6, with unanimous consensus that ‘the timeline to return to running should be person specific’. Respondents also reached consensus that one cannot start running before 3 weeks postpartum (85.5%) and that any birth injury should be completely healed before returning to running (97.6%).

### Current evidence

Pelvic health metrics—such as vaginal resting pressure, levator hiatus area, PFM strength and endurance, and bladder neck

mobility—have been shown to be altered after childbirth, particularly vaginal delivery in the general postpartum population.<sup>47 48 53</sup> Perineal trauma and surgical birth will also require adequate time for soft-tissue healing.<sup>6 140</sup> Although rare, risk for blood clots, hypertensive disorders, haemorrhage and sepsis is elevated in the first 6 weeks postpartum.<sup>129 141 142</sup> ACOG recommends all females have healthcare provider contact within 3 weeks postpartum, with a ‘comprehensive postpartum visit and transition to well-woman care’ between 4 and 12 weeks postpartum.<sup>143</sup>

Consensus (from Delphi respondents) was reached that returning to running before 12 weeks postpartum is possible. Longitudinal data investigating PFM function supports that returning to exercise within the first 12 weeks postpartum can be done successfully: PFM strength and endurance, vaginal resting pressure, POP and UI symptoms were similar at 1-year

## Consensus statement

postpartum in females who returned to exercise (including running) prior to 6 weeks postpartum and those who returned after 6 weeks postpartum.<sup>144</sup> Another longitudinal study demonstrated that early engagement in moderate-to-vigorous physical activity in the early postpartum period ( $\leq 6$  weeks) did not directly influence pelvic floor dysfunctions at 1-year postpartum, but was associated with a lower symptom burden.<sup>145</sup> Elite female athletes (including runners) often return to exercise before 6 weeks postpartum without increased incidence of incontinence.<sup>90</sup> A study of 42 elite runners (average return-to-running timeline of 6 weeks postpartum; training increased to 80% of prepregnancy levels by 14 weeks postpartum) found no association between musculoskeletal injury and timeline of return-to-running after childbirth.<sup>146</sup>

Data in postpartum recreational runners are more varied. Blyholder *et al*<sup>9</sup> reported that 49.2% of postpartum recreational runners returned within 6 weeks and 34.7% returned between 6 and 12 weeks. Moore *et al*<sup>8</sup> reported a median return-to-run time of 12 weeks (IQR 7–20 weeks), that returning-to-running increased the odds of developing SUI regardless of time frame compared with females who stopped running during pregnancy and did not return-to-running after childbirth, and that 84% of postpartum runners reported pain.<sup>8</sup> Christopher *et al*<sup>7</sup> reported a mean time of  $12.7 \pm 14.3$  weeks to first postpartum run, that 33% of postpartum runners reported running-related pain, and that timeline was not a significant risk factor for postpartum running-related pain.<sup>7</sup> However, some postpartum females have reported delaying return-to-run because they felt it was ‘too soon postpartum’.<sup>12</sup> It should be noted that the prevalence of PFD in athletes may be under-reported.<sup>147 148</sup> It should also be noted that there is no evidence on postpartum pelvic floor tissue healing timelines specifically in athletes. (Level of evidence: III)

### Recommendation (12/12 authors assent)

Given the range and complexity of factors involved (including injury, tissue healing time frames, pain and PFD symptoms), the lack of high-quality evidence, and the variability of local healthcare accessibility, a person-specific timeline of initiating postpartum running is recommended. Following a period of relative rest and recovery after childbirth, gradual progression of cardio-respiratory fitness and strength training is recommended prior to initiating running (Delphi consensus recommends a minimum of 3 weeks after childbirth prior to return-to-running). Prior training load—both before and during pregnancy—should also be considered. While many recreational runners may be able to return to running independently without significant issues, elite athletes and postpartum runners who are symptomatic (or otherwise concerned) should seek medical advice and/or evaluation by a pelvic health PT to determine run-readiness.

## DISCUSSION

To our knowledge, this is the first time an international consensus—consisting of multidisciplinary professionals—has established how postpartum run-readiness is currently determined. This Delphi survey, the corresponding literature review and expert recommendations (figure 3) start to address postpartum run-readiness and highlight knowledge gaps that need to be investigated. Due to the significant variability in postpartum runners, this study emphasises the importance of individualised, athlete centred decision-making. As not all runners will have access to health or fitness professionals, and evidence has demonstrated lack of education to perinatal runners,<sup>11 12</sup> this consensus statement also highlights the importance of

education of female runners and (where applicable) running coaches on return-to-running after childbirth.

### Research implications

Multiple gaps in research have been identified by this consensus survey and literature review. Future longitudinal studies exploring the development/progression of incontinence and prolapse during and after pregnancy in athletic populations are needed to further understand if screening and rehabilitation of postpartum runners can prevent symptoms of incontinence and prolapse when returning to running; or, if symptoms are already present, if a return-to-running progression can be performed in tandem with rehabilitation without worsening symptoms. Furthermore, the effectiveness of adjuncts to pelvic floor function (eg, compression garments or pessaries) should be explored. In addition, lactating females and females with larger breasts have historically been excluded from studies on breast support, which highlights the need for specific investigations into breast support for lactating athletes. Future studies should also evaluate the role of musculoskeletal strength, as well as gait and balance changes in postpartum RRI risk. Evaluation of pelvic floor healing timelines in athletes is also needed. Validation of all recommendations made in this consensus statement is also required. In general, more high-quality research is necessary in all areas of postpartum exercise, particularly high-impact exercise like running.

### Clinical implications

As healthcare providers and fitness professionals—particularly birth providers, primary care providers, personal trainers and PTs—are likely to be asked questions by perinatal runners, it is imperative that these providers are educated on this topic and can refer runners to the appropriate, evidence-informed information or provider to guide running during and after pregnancy.

### Limitations

Due to the lack of evidence guiding postpartum return-to-running, recommendations in this consensus statement were made based on integration of experienced professional consensus, literature review and discussion among expert researchers and clinicians in the field. As such, a narrative literature review—not a systematic review—was conducted for the literature review sections. Much of the evidence in this field is level III or below.

Respondents were predominantly white female PTs, and therefore, this study may not accurately reflect the opinions and experiences of other professionals (ie, physicians, male providers, those potentially in lower resource settings etc.) who may be the first contact and/or sole provider evaluating the runner. However, this is the first study to our knowledge that has included occupational therapists, chiropractors and running coaches. This study also included more personal trainers, exercise physiologists and physicians than the current expert opinion publications on postpartum running.<sup>16 18 19 22</sup> All the multidisciplinary participants had a voice in round 1 of the survey, thus informing the survey questions on which all participants voted. Due to the nature of Delphi methodology and multiple survey rounds, the number of respondents also decreased between rounds.

Also, several cultures may have different postpartum practices and rituals (eg, period of rest, confinement practices, avoidance of exercise, dietary requirements, breastfeeding

practices, etc.) that may conflict with the run-readiness recommendations in this Delphi study.<sup>149–152</sup> While efforts were made to recruit diverse respondents (through personal networks, social media (ie, Twitter, Instagram and Facebook) and word-of-mouth) and authors to capture sociocultural differences, the recommendations made in this consensus statement may not be applicable to every culture.

## CONCLUSION

Consensus was reached that postpartum runners were defined as anyone who self-identifies as a runner at any time after childbirth. Determining postpartum run-readiness is a multifactorial decision-making process that should be individualised and include the following components: (1) assessment of key musculoskeletal (including pelvic floor) and biomechanical milestones; (2) load and impact screening; (3) screening of biopsychosocial factors; (4) considerations of support items if needed and (5) the runner's training history, current capacity, running goals and training preferences. Due to the complexity of the postpartum experience, a multidisciplinary team approach (eg, primary care providers, lactation consultants, pelvic health PTs, mental health providers, sports medicine providers, orthopaedic specialists, physiatrists, obstetricians/gynaecologists, urogynaecologists, etc.) is recommended when feasible. Education of perinatal runners on postpartum recovery and gradual initiation of exercise is crucial. Further research is required in postpartum runners to identify specific tests and measures to determine readiness to return-to-running while mitigating injury risk and/or symptom provocation in this population.

### Author affiliations

- <sup>1</sup>Department of Rehabilitation Sciences, Doctor of Physical Therapy, Tufts University, Seattle, Washington, USA
- <sup>2</sup>Elon University, Elon, North Carolina, USA
- <sup>3</sup>Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, UK
- <sup>4</sup>Private Practice, Physiomum, Surrey, UK
- <sup>5</sup>Department of Sports Medicine, Norwegian School of Sports Sciences, Oslo, Norway
- <sup>6</sup>Department of Obstetrics and Gynecology, Akershus University Hospital, Lorenskog, Norway
- <sup>7</sup>Program for Pregnancy and Postpartum Health, Physical Activity and Diabetes Laboratory, Faculty of Kinesiology, Sport and Recreation, Women and Children's Health Research Institute, Alberta Diabetes Institute, University of Alberta, Edmonton, Alberta, Canada
- <sup>8</sup>Perinatal Physical Activity Research Group, School of Psychology and Life Sciences, Section of Sport, Exercise and Rehabilitation Science, Canterbury Christ Church University, Canterbury, UK
- <sup>9</sup>Active Pregnancy Foundation, None, UK
- <sup>10</sup>The World of my Baby (WOMB), Faculty of Health Science, McMaster University, Hamilton, Ontario, Canada
- <sup>11</sup>School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Queensland, Australia
- <sup>12</sup>LLC, Intimate Rose, Medford, Oregon, USA
- <sup>13</sup>Department of Physical Therapy, Carroll University, Waukesha, Wisconsin, USA
- <sup>14</sup>Department of Orthopedics & Rehabilitation, University of Wisconsin-Madison, Madison, Wisconsin, USA

**Twitter** Shefali Mathur Christopher @ironmompt, Gráinne Donnelly @ABSPhysio, Margie H Davenport @ExercisePreg, Marlice De Vivo @marlizedv, Hayley Mills @Drhbomb, Isabel S Moore @IzzyMoorePhD, Amanda Olson @amandaolsondpt and Rita E Deering @ritadeeringPhD

**Acknowledgements** The Delphi author group would like to extend their gratitude to all the pilot participants and the following researchers and experts who informed sections of the report relevant to their specialist fields: Dr Celeste Colman—University of Canberra, Australia; Dr Chris Mills—University of Portsmouth. Thanks, also, to research assistant Katelyn Hickey DPT.

**Contributors** SMC, RED, GD and EB convened the author group. All authors conceived the idea for this Delphi study. SMC, RED, SD and MHD performed the

thematic coding and data analysis. SMC and RED wrote the initial draft of the manuscript. All authors contributed to reviewing and giving feedback on each iteration of the survey and manuscript drafts. All authors contributed to the literature review. All authors reviewed the final manuscript.

**Funding** MHD is the Christenson Professor in Active Healthy Living.

**Competing interests** ISM is an associate editor of *BJSM*.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants but Elon University IRB 22-112 exempted this study. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

### ORCID iDs

Shefali Mathur Christopher <http://orcid.org/0000-0003-0869-8856>  
 Margie H Davenport <http://orcid.org/0000-0001-5627-5773>  
 Marlice De Vivo <http://orcid.org/0000-0002-7873-5985>  
 Lori Forner <http://orcid.org/0000-0002-5486-7350>  
 Hayley Mills <http://orcid.org/0000-0001-8895-8894>  
 Isabel S Moore <http://orcid.org/0000-0002-4746-3390>  
 Rita E Deering <http://orcid.org/0000-0002-8187-9971>

## REFERENCES

- 1 Elliott-Sale KJ, McNulty KL, Ansdell P, *et al*. The effects of oral contraceptives on exercise performance in women: A systematic review and meta-analysis. *Sports Med* 2020;50:1785–812.
- 2 McNulty KL, Elliott-Sale KJ, Dolan E, *et al*. The effects of Menstrual cycle phase on exercise performance in Eumenorrhoeic women: A systematic review and meta-analysis. *Sports Med* 2020;50:1813–27.
- 3 Bondarev D, Finni T, Kokko K, *et al*. Physical performance during the menopausal transition and the role of physical activity. *J Gerontol A Biol Sci Med Sci* 2021;76:1587–90.
- 4 Bø K, Artal R, Barakat R, *et al*. Exercise and pregnancy in recreational and elite athletes: 2016 evidence summary from the IOC expert group meeting, Lausanne. part 1—exercise in women planning pregnancy and those who are pregnant. *Br J Sports Med* 2016;50:571–89.
- 5 Bø K, Artal R, Barakat R, *et al*. Exercise and pregnancy in recreational and elite athletes: 2016 evidence summary from the IOC expert group meeting, Lausanne. part 2—The effect of exercise on the fetus, labor and birth. *Br J Sports Med* 2016;50:1297–305.
- 6 Bø K, Artal R, Barakat R, *et al*. Exercise and pregnancy in recreational and elite athletes: 2016/17 evidence summary from the IOC expert group meeting, Lausanne. part 3—Exercise in the postpartum period. *Br J Sports Med* 2017;51:1516–25.
- 7 Christopher SM, Cook CE, Snodgrass SJ. What are the Biopsychosocial risk factors associated with pain in postpartum runners? development of a clinical decision tool. *PLoS One* 2021;16:e0255383.
- 8 Moore IS, James ML, Brockwell E, *et al*. Multidisciplinary, Biopsychosocial factors contributing to return to running and running related stress urinary incontinence in postpartum women. *Br J Sports Med* 2021;55:1286–92.
- 9 Blyholder L, Chumanov E, Carr K, *et al*. Exercise behaviors and health conditions of runners after childbirth. *Sports Health* 2017;9:45–51.
- 10 Tenforde AS, Toth KES, Langen E, *et al*. Running habits of competitive runners during pregnancy and Breastfeeding. *Sports Health* 2015;7:172–6.
- 11 Donnelly GM, James ML, Coltman CE, *et al*. How does running-related advice and guidance received during pregnancy and postpartum affect women's running habits. *J Women's Health Physical Therapy* 2022;46:124–31.
- 12 James ML, Moore IS, Donnelly GM, *et al*. Running during pregnancy and postpartum, part A: Why do women stop running during pregnancy and not return to running in the postpartum period. *J Womens Health Phys Therap* 2022;46:111–23.
- 13 Davenport MH, Ray L, Nesdoly A, *et al*. We're not superhuman, we're human: A qualitative description of elite athletes' experiences of return to sport after childbirth. *Sports Med* 2023;53:269–79.
- 14 Ardern CL, Glasgow P, Schneiders A, *et al*. Consensus statement on return to sport from the first world Congress in sports physical therapy, Bern. *Br J Sports Med* 2016;50:853–64.

- 15 Brinlee AW, Dickenson SB, Hunter-Giordano A, et al. ACL reconstruction rehabilitation: clinical data, biologic healing, and criterion-based milestones to inform a return-to-sport guideline. *Sports Health* 2022;14:770–9.
- 16 Christopher SM, Gallagher S, Olson A, et al. Rehabilitation of the postpartum runner: A 4-phase approach. *J Womens Health Phys Therap* 2022;46:73–86.
- 17 Donnelly GM, Brockwell E, Rankin A, et al. Beyond the musculoskeletal system: considering whole-systems readiness for running. *J Women's Health Phys Ther* 2022;46:48–56.
- 18 Donnelly GM, Rankin A, Mills H, et al. Guidance for medical, health and fitness professionals to support women in returning to running Postnatally. *Br J Sports Med* 2020;54:1114–5.
- 19 Donnelly G, Brockwell E, Goom T. Return to running postnatal - guideline for medical, health and fitness professionals managing this population. *Physiotherapy* 2020;107:e188–9.
- 20 Deering RE, Christopher SM, Heiderscheidt BC. From childbirth to the starting blocks: are we providing the best care to our postpartum athletes? *J Orthop Sports Phys Ther* 2020;50:281–4.
- 21 Donnelly GM, Moore IS, Brockwell E, et al. Reframing return-to-sport postpartum: the 6 rs framework. *Br J Sports Med* 2022;56:244–5.
- 22 Selman R, Early K, Battles B, et al. Maximizing recovery in the postpartum period: A Timeline for rehabilitation from pregnancy through return to sport. *Int J Sports Phys Ther* 2022;17:1170–83.
- 23 Deering RE, Donnelly GM, Brockwell E, et al. Clinical and exercise professional opinion on designing a postpartum return-to-running training programme: an international Delphi study and consensus statement. *Br J Sports Med* 2024;58:183–95.
- 24 Rowe G, Wright G. The Delphi technique as a forecasting tool: issues and analysis. *Int J Forecasting* 1999;15:353–75.
- 25 Jünger S, Payne SA, Brine J, et al. Guidance on conducting and reporting Delphi studies (CREDES) in palliative care: recommendations based on a methodological systematic review. *Palliat Med* 2017;31:684–706.
- 26 Niederberger M, Spranger J. Delphi technique in health sciences: A map. *Front Public Health* 2020;8:457.
- 27 Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs* 2000;32:1008–15.
- 28 McMillan SS, King M, Tully MP. How to use the nominal group and Delphi techniques. *Int J Clin Pharm* 2016;38:655–62.
- 29 Wilkie K, Thornton JS, Vinther A, et al. Clinical management of acute low back pain in elite and Subelite rowers: a Delphi study of experienced and expert Clinicians. *Br J Sports Med* 2021;55:1324–34.
- 30 Dufour S, Bernard S, Murray-Davis B, et al. Establishing expert-based recommendations for the conservative management of pregnancy-related Diastasis rectus abdominis: A Delphi consensus study. *J Womens Health Phys Therap* 2019;43:73–81.
- 31 Trevelyan EG, Robinson PN. Delphi methodology in health research: how to do it. *Europ J Integrat Med* 2015;7:423–8.
- 32 Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends Methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67:401–9.
- 33 Sackett DL. Rules of evidence and clinical recommendations on the use of Antithrombotic agents. *Chest* 1989;95:25–45.
- 34 Foch E, Brindle RA, Pohl MB. Lower extremity Kinematics during running and hip Abductor strength in Iliotibial band syndrome: A systematic review and meta-analysis. *Gait Posture* 2023;101:73–81.
- 35 Kakouris N, Yener N, Fong DTP. A systematic review of running-related musculoskeletal injuries in runners. *J Sport Health Sci* 2021;10:513–22.
- 36 Paladine HL, Blenning CE, Strangas Y. Postpartum care: an approach to the fourth trimester. *Am Fam Physician* 2019;100:485–91.
- 37 Thein-Nissenbaum J. The postpartum Triathlete. *Phys Ther Sport* 2016;21:95–106.
- 38 Deering RE, Chumanov ES, Stiffler-Joachim MR, et al. Exercise program reduces inter-Recti distance in female runners up to two years postpartum. *J Womens Health Phys Therap* 2020;44:9–18.
- 39 Hills NF, Graham RB, McLean L. Comparison of trunk muscle function between women with and without Diastasis Recti abdominis at 1 year postpartum. *Phys Ther* 2018;98:891–901.
- 40 Deering R, Pashibin T, Cruz M, et al. Fatiguing trunk Flexor exercise decreases pain sensitivity in postpartum women. *Front Physiol* 2019;10:315.
- 41 Deering RE, Cruz M, Senefeld JW, et al. Impaired trunk Flexor strength, fatigability, and steadiness in postpartum women. *Med Sci Sports Exerc* 2018;50:1558–69.
- 42 Deering RE, Senefeld J, Pashibin T, et al. Fatigability of the Lumbopelvic stabilizing muscles in women 8 and 26 weeks postpartum. *J Womens Health Phys Therap* 2018;42:128–38.
- 43 Christopher SM, Bauer L, Maylone R, et al. Biomechanical and musculoskeletal differences between postpartum runners and nulliparous controls. *J Womens Health Phys Therap* 2022;46:11–7.
- 44 Putnick DL, Sundaram R, Bell EM, et al. Trajectories of maternal postpartum depressive symptoms. *Pediatrics* 2020;146:e20200857.
- 45 Christopher SM, Garcia AN, Snodgrass SJ, et al. Common musculoskeletal impairments in postpartum runners: an international Delphi study. *Arch Physiother* 2020;10:19.
- 46 Guzmán Rojas R, Wong V, Shek KL, et al. Impact of Levator trauma on pelvic floor muscle function. *Int Urogynecol J* 2014;25:375–80.
- 47 Hilde G, Stær-Jensen J, Siafarikas F, et al. Impact of childbirth and mode of delivery on vaginal resting pressure and on pelvic floor muscle strength and endurance. *Am J Obstet Gynecol* 2013;208:S0002-9378(12)01978-3.
- 48 Hilde G, Stær-Jensen J, Siafarikas F, et al. How well can pelvic floor muscles with major defects contract? A cross-sectional comparative study 6 weeks after delivery using Transperineal 3D/4D ultrasound and manometer. *BIOG* 2013;120:1423–9.
- 49 Martin-Martin S, Pascual-Fernandez A, Alvarez-Colomo C, et al. Urinary Incontinence during pregnancy and postpartum. associated risk factors and influence of pelvic floor exercises. *Arch Esp Urol* 2014;67:323–30.
- 50 Nygaard IE, Shaw JM. Physical activity and the pelvic floor. *Am J Obstet Gynecol* 2016;214:164–71.
- 51 Reimers C, Siafarikas F, Stær-Jensen J, et al. Risk factors for anatomic pelvic organ Prolapse at 6 weeks postpartum: a prospective observational study. *Int Urogynecol J* 2019;30:477–82.
- 52 Shin GH, Toto EL, Schey R. Pregnancy and postpartum bowel changes: constipation and fecal incontinence. *Am J Gastroenterol* 2015;110:521–9.
- 53 Stær-Jensen J, Siafarikas F, Hilde G, et al. Postpartum recovery of Levator hiatus and bladder neck mobility in relation to pregnancy. *Obstet Gynecol* 2015;125:531–9.
- 54 Vermandel A, De Wachter S, Beyltiens T, et al. Pelvic floor awareness and the positive effect of verbal instructions in 958 women early Postdelivery. *Int Urogynecol J* 2015;26:223–8.
- 55 Wallace SL, Miller LD, Mishra K. Pelvic floor physical therapy in the treatment of pelvic floor dysfunction in women. *Curr Opin Obstet Gynecol* 2019;31:485–93.
- 56 Hadizadeh-Talasz Z, Sadeghi R, Khadivzadeh T. Effect of pelvic floor muscle training on postpartum sexual function and quality of life: A systematic review and meta-analysis of clinical trials. *Taiwan J Obstet Gynecol* 2019;58:737–47.
- 57 Carvalhais A, Araújo J, Natal Jorge R, et al. Urinary Incontinence and disordered eating in female elite athletes. *J Sci Med Sport* 2019;22:140–4.
- 58 de Mattos Lourenco TR, Matsuoka PK, Baracat EC, et al. Urinary Incontinence in female athletes: a systematic review. *Int Urogynecol J* 2018;29:1757–63.
- 59 Forner LB, Beckman EM, Smith MD. Do women runners report more pelvic floor symptoms than women in Crossfit®? A cross-sectional survey. *Int Urogynecol J* 2021;32:295–302.
- 60 National Institute for Health and Care Excellence: Guidelines. *Pelvic floor dysfunction: prevention and non-surgical management*. 2021.
- 61 Borello-France DF, Handa VL, Brown MB, et al. Pelvic-floor muscle function in women with pelvic organ Prolapse. *Phys Ther* 2007;87:399–407.
- 62 Blomquist JL, Carroll M, Muñoz A, et al. Pelvic floor muscle strength and the incidence of pelvic floor disorders after vaginal and cesarean delivery. *Am J Obstet Gynecol* 2020;222:S0002-9378(19)31001-4.
- 63 Moser H, Leitner M, Baeyens JP, et al. Pelvic floor muscle activity during impact activities in continent and incontinent women: a systematic review. *Int Urogynecol J* 2018;29:179–96.
- 64 Woodley SJ, Lawrenson P, Boyle R, et al. Pelvic floor muscle training for preventing and treating urinary and Faecal Incontinence in Antenatal and postnatal women. *Cochrane Database Syst Rev* 2020;5:CD007471.
- 65 Peterson B, Hawke F, Spink M, et al. Biomechanical and musculoskeletal measurements as risk factors for running-related injury in non-elite runners: A systematic review and meta-analysis of prospective studies. *Sports Med Open* 2022;8:38.
- 66 Gluppe S, Ellström Engh M, Kari B. Women with Diastasis Recti abdominis might have weaker abdominal muscles and more abdominal pain, but no higher prevalence of pelvic floor disorders, low back and pelvic girdle pain than women without Diastasis Recti abdominis. *Physiotherapy* 2021;111:57–65.
- 67 Fuentes Aparicio L, Rejano-Campo M, Donnelly GM, et al. Self-reported symptoms in women with Diastasis rectus abdominis: A systematic review. *J Gynecol Obstet Hum Reprod* 2021;50:S2468-7847(20)30365-2.
- 68 Sperstad JB, Tennfjord MK, Hilde G, et al. Diastasis Recti abdominis during pregnancy and 12 months after childbirth: prevalence, risk factors and report of Lumbopelvic pain. *Br J Sports Med* 2016;50:1092–6.
- 69 Crommert ME, Flink I, Gustavsson C. Predictors of disability attributed to symptoms of increased Interrecti distance in women after childbirth: an observational study. *Phys Ther* 2021;101.
- 70 Eriksson Crommert M, Petrov Fieril K, Gustavsson C. Women's experiences of living with increased inter-Recti distance after childbirth: an interview study. *BMC Womens Health* 2020;20:260.
- 71 Gluppe S, Engh ME, Bø K. What is the evidence for abdominal and pelvic floor muscle training to treat Diastasis Recti abdominis postpartum? A systematic review with meta-analysis. *Braz J Phys Ther* 2021;25:664–75.

- 72 Carlstedt A, Bringman S, Egberth M, *et al.* Management of Diastasis of the rectus abdominis muscles: recommendations for Swedish national guidelines. *Scand J Surg* 2021;110:452–9.
- 73 Gluppe SB, Ellström Engh M, Bø K. Curl-up exercises improve abdominal muscle strength without worsening inter-Recti distance in women with Diastasis Recti abdominis postpartum: a randomised controlled trial. *J Physiother* 2023;69:160–7.
- 74 Goossens N, Massé-Alarie H, Aldabe D, *et al.* Changes in static balance during pregnancy and postpartum: A systematic review. *Gait Posture* 2022;96:160–72.
- 75 Bagwell JJ, Reynolds N, Katsavelis D, *et al.* Center of pressure characteristics differ during single leg stance throughout pregnancy and compared to Nulligravida individuals. *Gait Posture* 2022;97:43–7.
- 76 Ramachandra P, Kumar P, Bø K, *et al.* Comparison of static postural sway characteristics between pregnant and non-pregnant women. *J Biomech* 2023;154:111618.
- 77 Han J, Luan L, Adams R, *et al.* Can therapeutic exercises improve Proprioception in chronic ankle instability? A systematic review and network meta-analysis. *Arch Phys Med Rehabil* 2022;103:2232–44.
- 78 Plasteras CT, Rittenberg JD, Rittenberg KE, *et al.* Comprehensive functional evaluation of the injured runner. *Phys Med Rehabil Clin N Am* 2005;16:623–49.
- 79 van der Worp MP, ten Haaf DSM, van Cingel R, *et al.* Injuries in runners; a systematic review on risk factors and sex differences. *PLoS One* 2015;10:e0114937.
- 80 West TJ, Bruder AM, Crossley KM, *et al.* Unilateral tests of lower-limb function as Prognostic indicators of future knee-related outcomes following anterior Cruciate ligament injury: a systematic review and meta-analysis of 13 150 adolescents and adults. *Br J Sports Med* 2023;57:855–63.
- 81 Harrison K, Williams DSB, Darter BJ, *et al.* The running readiness scale as an assessment of Kinematics related to knee injury in novice female runners. *J Athl Train* 2023;58:120–7.
- 82 Goodson C, McLeod AR, Kearns Z, *et al.* External loading of common training drills: ranking drills to design progressive return-to-run programs. *Phys Ther Sport* 2022;58:167–72.
- 83 Casey EK, Temme K. Pelvic floor muscle function and urinary Incontinence in the female athlete. *Phys Sportsmed* 2017;45:399–407.
- 84 Pires T, Pires P, Moreira H, *et al.* Prevalence of urinary Incontinence in high-impact sport athletes: A systematic review and meta-analysis. *J Hum Kinet* 2020;73:279–88.
- 85 Christopher S, McCullough J, Snodgrass SJ, *et al.* Predictive risk factors for first-onset Lumbopelvic pain in postpartum women: A systematic review. *J Womens Health Phys Therap* 2019;43:127–35.
- 86 Whitney KE, Holtzman B, Cook D, *et al.* Low energy availability and impact sport participation as risk factors for urinary Incontinence in female athletes. *J Pediatr Urol* 2021;17:S1477-5131(21)00072-3.
- 87 Ackerman KE, Holtzman B, Cooper KM, *et al.* Low energy availability Surrogates correlate with health and performance consequences of relative energy deficiency in sport. *Br J Sports Med* 2019;53:628–33.
- 88 Deering RE, Mountjoy ML. Reds and the lactating athlete: an evidence gap. *Br J Sports Med* 2023;57:1065–6.
- 89 Kolomańska-Bogucka D, Mazur-Bialy AI. Physical activity and the occurrence of postnatal depression-A systematic review. *Medicina (Kaunas)* 2019;55:560.
- 90 Kimber ML, Meyer S, McHugh T-L, *et al.* Health outcomes following pregnancy in elite athletes: A systematic review and meta-analysis. *Med Sci Sports Exerc* 2021;53:1739–47.
- 91 Bugge C, Adams EJ, Gopinath D, *et al.* Pessaries (mechanical devices) for managing pelvic organ Prolapse in women. *Cochrane Database Syst Rev* 2020;11:CD004010.
- 92 Lipp A, Shaw C, Glavind K. Mechanical devices for urinary Incontinence in women. *Cochrane Database Syst Rev* 2014;2014:CD001756.
- 93 Dufour S, Wu M. No. 397 - conservative care of urinary Incontinence in women. *J Obstet Gynaecol Can* 2020;42:510–22.
- 94 Dakic JG, Cook J, Hay-Smith J, *et al.* Pelvic floor disorders stop women exercising: A survey of 4556 symptomatic women. *J Sci Med Sport* 2021;24:1211–7.
- 95 Brennan E, Ruiz-Mirazo E, Tang S, *et al.* Urinary leakage during exercise: problematic activities, adaptive behaviors, and interest in treatment for physically active Canadian women. *Int Urogynecol J* 2018;29:497–503.
- 96 Culleton-Quinn E, Bø K, Fleming N, *et al.* Elite female athletes' experiences of symptoms of pelvic floor dysfunction: A systematic review. *Int Urogynecol J* 2022;33:2681–711.
- 97 McGhee DE, Steele JR. Biomechanics of breast support for active women. *Exerc Sport Sci Rev* 2020;48:99–109.
- 98 Risius D, Milligan A, Berns J, *et al.* Understanding key performance indicators for breast support: an analysis of breast support effects on Biomechanical, physiological and subjective measures during running. *J Sports Sci* 2017;35:842–51.
- 99 Burbage J, Cameron L. An investigation into the prevalence and impact of breast pain, Bra issues and breast size on female horse riders. *J Sports Sci* 2017;35:1091–7.
- 100 Nolte K, Burgoyne S, Nolte H, *et al.* The effectiveness of a range of sports Bras in reducing breast displacement during treadmill running and two-step star jumping. *J Sports Med Phys Fitness* 2016;56:1311–7.
- 101 Brisbane BR, Steele JR, Phillips EJ, *et al.* Breast pain affects the performance of elite female athletes. *J Sports Sci* 2020;38:528–33.
- 102 Morris K, Park J, Sarkar A. Development of a nursing sports Bra for physically active Breastfeeding women through user-centered design. *Clothing Textiles Res J* 2017;35:290–306.
- 103 Coltman CE, Steele JR, McGhee DE. Does breast size affect how women participate in physical activity. *J Sci Med Sport* 2019;22:324–9.
- 104 Fong HB, Powell DW. Greater breast support is associated with reduced oxygen consumption and greater running economy during a treadmill running task. *Front Sports Act Living* 2022;4:902276.
- 105 Alcantara RS, Wall-Scheffler CM. Stroller running: energetic and Kinematic changes across pushing methods. *PLoS One* 2017;12:e0180575.
- 106 Sandbakk Ø, Perl R, Holmberg HC, *et al.* Energetic cost and Kinematics of pushing a Stroller on flat and uphill terrain. *Front Physiol* 2020;11:574.
- 107 O'Sullivan R, Kiernan D, Malone A. Run Kinematics with and without a Jogging Stroller. *Gait Posture* 2016;43:220–4.
- 108 Szkwara JM, Milne N, Hing W, *et al.* Effectiveness, feasibility, and acceptability of dynamic Elastomeric fabric Orthoses (DEFO) for managing pain, functional capacity, and quality of life during Prenatal and postnatal care: a systematic review. *Int J Environ Res Public Health* 2019;16:2408.
- 109 Gavrilov SG, Karalkin AV, Turischeva OO. Compression treatment of pelvic congestion syndrome. *Phlebology* 2018;33:418–24.
- 110 Szkwara JM, Milne N, Rathbone E. A prospective quasi-experimental controlled study evaluating the use of dynamic elastomeric fabric orthoses to manage common postpartum ailments during postnatal care. *Womens Health (Lond)* 2020;16.
- 111 Okayama H, Ninomiya S, Naito K, *et al.* Effects of wearing supportive underwear versus pelvic floor muscle training or no treatment in women with symptoms of stress urinary Incontinence: an assessor-blinded randomized control trial. *Int Urogynecol J* 2019;30:1093–9.
- 112 Milner M, Gamble M, Barry-Kinsella C. Covid-19, pelvic health, and women's voices: A descriptive study. *Continence* 2022;1:100012.
- 113 Fitzgerald CM, Bennis S, Marcotte ML, *et al.* The impact of a Sacroiliac joint belt on function and pain using the active straight leg raise in pregnancy-related pelvic girdle pain. *PM R* 2022;14:19–29.
- 114 Arumugam A, Milosavljevic S, Woodley S, *et al.* The effect of a pelvic compression belt on functional hamstring muscle activity in sportsmen with and without previous hamstring injury. *Int J Sports Phys Ther* 2015;10:291–302.
- 115 Hammer N, Möbius R, Schleifenbaum S, *et al.* Pelvic belt effects on health outcomes and functional parameters of patients with Sacroiliac joint pain. *PLoS ONE* 2015;10:e0136375.
- 116 Soisson O, Lube J, Germano A, *et al.* Pelvic belt effects on pelvic Morphometry, muscle activity and body balance in patients with Sacroiliac joint dysfunction. *PLoS ONE* 2015;10:e0116739.
- 117 Toyohara R, Hiramukai T, Kurosawa D, *et al.* Numerical analysis of the effects of padded pelvic belts as a treatment for Sacroiliac joint dysfunction. *BME* 2023;34:305–18.
- 118 Jobanputtra Y, Patil S. Immediate effect of Kinesio Taping on Lumbopelvic stability in postpartum women with Diastasis Recti: A review. *Cureus* 2023;15:e33347.
- 119 Ptaszowska L, Gorecka J, Paprocka-Borowicz M, *et al.* Immediate effects of Kinesio Taping on rectus abdominis Diastasis in postpartum women-preliminary report. *J Clin Med* 2021;10:5043:21..
- 120 Malhotra D, Sharma S, Chachra A, *et al.* The time-based effects of Kinesio Taping on acute-onset muscle soreness and calf muscle Extensibility among endurance athletes: A randomized cross-over trial. *J Clin Med* 2022;11:5996.
- 121 Watcharakhueakhan P, Chapman GJ, Sinsurin K, *et al.* The immediate effects of Kinesio Taping on running Biomechanics, muscle activity, and perceived changes in comfort, stability and running performance in healthy runners, and the implications to the management of Iliotibial band syndrome. *Gait & Posture* 2022;91:179–85.
- 122 Ozmen T, Aydogmus M, Dogan H, *et al.* The effect of Kinesio Taping on muscle pain, Sprint performance, and flexibility in recovery from squat exercise in young adult women. *J Sport Rehabil* 2016;25:7–12.
- 123 Rogan S, Baur H. Examination of the effects of Kinesiotape and IQ tape on neuromuscular activity during running, stair climbing and drop jumps. *Sportverletz Sportschaden* 2020;34:e1.
- 124 Tousignant-Lafamme Y, Christopher S, Clewley D, *et al.* Does shared decision making results in better health related outcomes for individuals with painful musculoskeletal disorders? A systematic review. *J Man Manip Ther* 2017;25:144–50.
- 125 Hoffmann T, Bakhit M, Michaleff Z. Shared decision making and physical therapy: what, when, how, and why? *Braz J Phys Ther* 2022;26:100382.
- 126 Makama M, Awoke MA, Skouteris H, *et al.* Barriers and Facilitators to a healthy lifestyle in postpartum women: A systematic review of qualitative and quantitative studies in postpartum women and healthcare providers. *Obes Rev* 2021;22:e13167.
- 127 Edie R, Laceywell A, Streisel C, *et al.* Barriers to exercise in postpartum women: A mixed-methods systematic review. *J Womens Health Phys Therap* 2021;45:83–92.
- 128 Pelvic organ Prolapse: ACOG practice Bulletin, number 214. *Obstet Gynecol* 2019;134:e126–42.
- 129 Symonds IM, Arulkumaran S. *Essential obstetrics and gynaecology (6th edn)*. Edinburgh: Elsevier, 2020. Available: <http://www.clinicalkey.com/dura/browse/bookChapter/3-s2.0-C20170030744>

- 130 van Gent RN, Siem D, van Middelkoop M, *et al.* Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med* 2007;41:469–80;
- 131 Arnold MJ, Moody AL. Common running injuries: evaluation and management. *Am Fam Physician* 2018;97:510–6.
- 132 Bø K, Driusso P, Jorge CH. Can you breathe yourself to a better pelvic floor? A systematic review. *NeuroUrol Urodyn* 2023;42:1261–79.
- 133 Nielsen RO, Nohr EA, Rasmussen S, *et al.* Classifying running-related injuries based upon etiology, with emphasis on volume and pace. *Int J Sports Phys Ther* 2013;8:172–9.
- 134 Ramskov D, Rasmussen S, Sørensen H, *et al.* Progression in running intensity or running volume and the development of specific injuries in recreational runners: run clever, a randomized trial using competing risks. *J Orthop Sports Phys Ther* 2018;48:740–8.
- 135 Nielsen RØ, Parner ET, Nohr EA, *et al.* Excessive progression in weekly running distance and risk of running-related injuries: an association which varies according to type of injury. *J Orthop Sports Phys Ther* 2014;44:739–47.
- 136 Ndikom CM, Fawole B, Ilesanmi RE, *et al.* Extra fluids for Breastfeeding mothers for increasing milk production. *Cochrane Database Syst Rev* 2014;2014.
- 137 Bardosono S, Morin C, Guelinckx I, *et al.* Pregnant and Breastfeeding women: drinking for two *Ann Nutr Metab* 2017;70 Suppl 1:13–7.
- 138 Lee S, Kelleher SL. Biological underpinnings of Breastfeeding challenges: the role of Genetics, diet, and environment on Lactation physiology. *Am J Physiol Endocrinol Metab* 2016;311:E405–22.
- 139 Nutrient Reference Values for Australia and New Zealand. National health and medical research Council. 2020.
- 140 Tilak M, Mann GK, Gong M, *et al.* Pelvic floor healing milestones after obstetric Anal sphincter injury: a prospective case control feasibility study. *Int Urogynecol J* 2023;34:553–61.
- 141 Li XL, Guo PL, Xue Y, *et al.* An analysis of the differences between early and late Preeclampsia with severe hypertension. *Pregnancy Hypertens* 2016;6:47–52.
- 142 Snydal S. Major changes in diagnosis and management of Preeclampsia. *J Midwifery Womens Health* 2014;59:596–605.
- 143 ACOG committee opinion No.736: optimizing postpartum care. *Obstet Gynecol* 2018;131:e140–50.
- 144 Tennfjord MK, Engh ME, Bø K. The influence of early exercise postpartum on pelvic floor muscle function and prevalence of pelvic floor dysfunction 12 months postpartum. *Phys Ther* 2020;100:1681–9.
- 145 Nygaard IE, Wolpert A, Bardsley T, *et al.* Early postpartum physical activity and pelvic floor support and symptoms 1 year postpartum. *Am J Obstet Gynecol* 2021;224:193.
- 146 Darroch F, Schneeberg A, Brodie R, *et al.* Effect of pregnancy in 42 elite to world-class runners on training and performance outcomes. *Med Sci Sports Exerc* 2023;55:93–100.
- 147 Rodríguez-López ES, Acevedo-Gómez MB, Romero-Franco N, *et al.* Urinary Incontinence among elite track and field athletes according to their event specialization: A cross-sectional study. *Sports Med Open* 2022;8:78.
- 148 Mahoney K, Heidel RE, Olewinski L. Prevalence and normalization of stress urinary Incontinence in female strength athletes. *J Strength Cond Res* 2023;37:1877–81.
- 149 Dennis CL, Fung K, Grigoriadis S, *et al.* Traditional postpartum practices and rituals: a qualitative systematic review. *Womens Health (Lond)* 2007;3:487–502.
- 150 Grigoriadis S, Erlick Robinson G, Fung K, *et al.* Traditional postpartum practices and rituals: clinical implications. *Can J Psychiatry* 2009;54:834–40.
- 151 Withers M, Kharazmi N, Lim E. Traditional beliefs and practices in pregnancy, childbirth and postpartum: A review of the evidence from Asian countries. *Midwifery* 2018;56:158–70.
- 152 Liu YQ, Petrini M, Maloni JA. Doing the month": postpartum practices in Chinese women. *Nurs Health Sci* 2015;17:5–14.