

INTRODUCTION

The need for sleep

Normal human sleep

Normal human sleep comprises two main types – non-rapid eye movement sleep (non-REM) and rapid eye movement sleep (REM).[1] Non-REM and REM sleep serve many physiologic functions and play many important roles for maintenance and restoration of the brain and body.[2, 3] Non-REM sleep is divided into three stages, representing a continuum from ‘light’ sleep in stages 1 and 2, through to ‘deep’ sleep in stage 3, or slow-wave sleep (Supplementary Figure 1).[4] Under the original classification system,[5] non-REM sleep was divided into four stages, with two stages of slow-wave sleep, i.e., stages 3 and 4. However, in the more recent, modified classification system,[4] non-REM sleep is divided into three stages, with the original stages 3 and 4 combined into a single stage of slow-wave sleep, i.e., stage 3.

Characteristics of sleep

The different sleep stages are distinguished from wake and from each other by characteristic patterns of brain activity, eye movements, and muscle tone, as assessed in the laboratory with polysomnography, electroencephalogram, electrooculogram and electromyogram measurements. However, from a behavioural perspective, sleep is defined as a reversible, naturally recurring state characterised by closed eyes, recumbent body position, reduced body movement, reduced responsiveness to external stimuli and a reduced breathing rate.[3] During non-REM sleep, the brain is relatively inactive, although it is actively regulating vital functions, and the body is movable.[1] In contrast, during REM sleep, the brain is highly

activated, while the body is essentially paralysed.[1] It was originally thought that REM sleep was the only stage in which dreaming occurs, but it is now known that dreaming also occurs in non-REM sleep, albeit about half as often compared to REM sleep.[6]

Composition of sleep

The composition of normal sleep changes across the life cycle. In early-primary children, sleep is relatively long – at least 9–10 hours – and composed of a high proportion of slow-wave sleep; whereas in the elderly, sleep is relatively short – only 5.5–6.5 hours – and comprised of very little slow-wave sleep.[7] At the ages most relevant to aspiring and established elite athletes, an 8–9-hour sleep for an adolescent (aged 15 years) contains approximately 57% light sleep, 22% deep sleep, and 21% REM sleep; and a 7–8-hour sleep for a young adult (aged 30 years) contains approximately 61% light sleep, 16% deep sleep, and 23% REM sleep.[7]

Sleep and cognitive health

Until the late 1990s it was thought that chronic sleep loss led to increased sleepiness but had relatively little effect on cognitive function due to an adaptive response.[8] However, between 1997 and 2003, three seminal laboratory-based studies provided compelling evidence to refute this position. In these studies, sleep was restricted to (a) 5 hours each night for seven nights,[9] (b) 3, 5, 7, or 9 hours each night for seven nights,[10] and (c) 4, 6, or 8 hours each night for fourteen nights.[11] Together, these studies demonstrated that chronic sleep loss does impair cognitive function, particularly in sustained attention and response speed, and that the degree of impairment increases as the sleep dose decreases.

The repetition, or practice, of a task is a critical component of learning. Nevertheless, several studies have now shown that learning can also occur in the absence of practice, and that this process of practice-independent learning is enhanced by sleep.[12-14] In one of the early studies on this topic,[15] participants learnt a basic motor task (sequential finger-tapping) in the evening, then had a night of normal sleep or total sleep deprivation. In both cases, performance on the task improved when retested in the morning (increased speed and fewer errors) but the improvement was ~30% greater after a night of sleep compared to a night of wakefulness. Most importantly, it is now suggested that “it is practice, with sleep, that ultimately leads to perfection”. [12]

With regards mental well-being, poor sleep increases discomfort, impairs mood, and reduces the capacity to cope with stressors. When sleep is deliberately restricted to 4 hours each night over 12 nights, optimism and sociability are reduced, while bodily discomfort is elevated due to increased back pain, stomach pain and general body pain.[16] During 1–2 nights of total sleep deprivation, mood is substantially impaired such that feelings of depression, anger, frustration, tension, anxiety and paranoia all increase.[17-19]

Sleep and physical health

Human growth hormone is essential for the reproduction and regeneration of cells. The secretion of growth hormone is influenced by both the circadian system and the sleep/wake system. Under normal circumstances, these two systems work in concert, such that very little growth hormone is secreted during daytime wakefulness, and the majority of growth hormone is secreted at night, during non-REM, slow-wave sleep.[20] Sleep also plays an

important role in the defence against inflammatory challenges, particularly for those at increased risk of infection.[21] At a molecular level, a week of short sleep (5.7 hours sleep each night), upregulated the expression of genes related to tumours, chronic inflammation, and stress and downregulated the expression of genes related to immune function.[22] At a broader level, when a vaccine is given before a night of total sleep deprivation, or during a week of severe sleep restriction (4 hours sleep each night), the antibody response is reduced by about 50%.[23, 24] Furthermore, when people were exposed to a ‘cold’ virus, those who averaged < 7 hours of sleep each night in the preceding two weeks were three times more likely to develop a cold than those who averaged > 8 hours of sleep each night.[25]

It was previously argued that a moderate level of sleep loss, i.e., 2 hours below the optimal dose of ~8 hours each night, does not have serious consequences for human physiology.[26] However, this view is changing as evidence of the consequences of chronic sleep loss, particularly for the cardiometabolic system, grows. For example, when sleep is restricted to ~4–5 hours each night, the capacity to process glucose is impaired,[27, 28] hunger is stimulated,[29, 30] and vascular inflammation is increased.[31, 32] Furthermore, the epidemiological data indicate that these types of effects may eventually lead to adverse health outcomes. For example, habitual sleep durations of 5–6 hours each night, independent of other lifestyle factors, are associated with higher rates of type 2 diabetes,[33] obesity,[34] cardiovascular disease[35] and respiratory infection.[25] Perhaps for these reasons, many studies have shown that habitual short sleep duration is associated with increased mortality risk.[36]

Measuring sleep

With the increasing popularity of measuring sleep within both medical and consumer fields, unsurprisingly, the number of sleep measurement tools is rapidly increasing (Table 1). Some of the more common measurements regarding sleep include: sleep architecture (sleep staging), sleep duration, measured or estimated sleep efficiency (quality), sleep onset latency (SOL: time taken to fall asleep) and wake after sleep onset (WASO). Tools to measure sleep can be divided into several primary categories including: polysomnography, activity monitoring (including research grade and consumer wearables), smartphone applications and nearable devices (designed to be placed on or near the bed, hence ‘nearable’), sleep diaries and questionnaires. A brief description of each of these categories is included below and has been comprehensively reviewed elsewhere.[37]

Polysomnography

Polysomnography is considered the gold standard of sleep monitoring and typically includes an assessment of eye movement, brain activity, heart rate, muscle activity, oxygen saturation, breathing rate and body movement.[38] Polysomnography allows for the determination of REM and non-REM. As polysomnography can be an expensive, obtrusive and complex technique, it is typically utilised for the assessment of sleep disorders or in research studies. In addition to traditional polysomnography, there are an increasing number of commercial devices that use modified versions of polysomnography to record at-home sleep.[39] These are different from true ambulatory polysomnography units in that they do not collect the full montage of signals, typically employing a small number of electrodes (1–2 in most cases, often as part of a portable headband). These devices have shown utility at collecting objective

sleep data, but in most cases they are preliminary and are not as well-validated as in-lab recordings. In the case of athletes, who may not be able to come into a laboratory, these may be a viable alternative.

Activity monitoring

Activity monitors are wearable devices that record movement,[40] with most utilising a 3-axis accelerometer to determine sleep/wake based on a specific algorithm.[41] Currently, there are generally two classifications of devices – those that are considered “research-grade” and those that are considered “commercially-available.” “Research-grade” devices have been subject to rigorous validation, and often have become scientifically accepted as a measurement of habitual sleep duration, timing, and continuity.[42] Research-grade activity monitors have been validated against polysomnography,[43] the algorithms utilised to quantify sleep and wakefulness are reported and, importantly, several of these devices have been validated in an athlete population.[44, 45] These devices are useful for longitudinal monitoring and are popular devices in both research and practice.[40]

Both research-grade and commercially-available activity monitors typically overestimate sleep duration relative to polysomnography (and underestimate relative to recall and sleep diary), underestimate awakenings and WASO relative to polysomnography (and overestimate relative to recall and sleep diary), and generally underestimate sleep latency relative to both polysomnography and sleep diary.[42] These discrepancies are larger in the case of sleep disorders, including insomnia. Further, these devices typically perform poorer in those individuals with fragmented sleep (higher amounts of wakefulness), as the deficiency in these

devices is related to their ability to detect wakefulness.[40] These discrepancies may not always reflect inaccuracy; rather it may be the case that different methodologies for measuring sleep capture different aspects of sleep-wake regulation. Despite these discrepancies, activity monitoring is considered to be a valid and reliable method for evaluating sleep-wake rhythms in free living settings.[46] Recent reviews of commercially-available devices suggest that they can be useful for monitoring sleep,[41, 47, 48] including in athlete populations,[49] but this depends on the quality of the data supporting each device.[43, 50, 51]

Nearables and smartphone applications

One subset of wearable devices is placed on the mattress and measures movement and other physiologic signals of the individual lying on the bed (typically using ballistocardiography).[52, 53] Currently, validation of available devices is sparse, little is known about the algorithms used, and these devices are not very accurate for measuring sleep.[54] Another type of wearable device sits at the bedside and detects movement and physiologic signals of the individual lying on the bed. Similarly, this technology shows promise for detecting sleep and wake,[55] but there is still a relative lack of supporting validation for available devices.

Consumer-targeted smartphone applications (apps) use motion detection, audio and/or video recording as well as questionnaires[56] for sleep assessment and self-management.[57] In general, smartphone apps are poor at determining sleep stages and sleep parameters[41] and researchers have little room to influence or access raw data.[58] Readers are directed

elsewhere for a comprehensive review of current smartphone applications for sleep assessment and self-management.[57]

Sleep diaries

Sleep diaries can be a simple and cost-effective means of assessing sleep and typically include: bed and wake time, lights-out time, daytime napping, ratings of sleepiness and alertness, caffeine and alcohol intake, exercise and light emitting device use[59] for a duration of at least one week. One example is the consensus sleep diary that resulted from the collaboration of a team of sleep experts.[60] Limitations of sleep diaries include recall bias and response expectation.

Sleep questionnaires

Questionnaires are often utilised for screening purposes or an initial assessment of sleep due to their ease of administration and low cost. Common questionnaires in insomnia research include: the Pittsburgh Sleep Quality Index (PSQI) to assess sleep quality,[61] the Sleep Hygiene Index to assess sleep hygiene,[62] the Leeds Sleep Evaluation Questionnaire to assess various aspects of sleep and early morning behaviour[63] and the Epworth Sleepiness Scale[64] to assess daytime sleepiness. However, these questionnaires have not been validated for athletes. Athlete specific questionnaires include the Athlete Sleep Screening Questionnaire[65, 66] and the Athlete Sleep Behaviour Questionnaire.[67] Similar to sleep diaries, while questionnaires may be time and cost effective, response biases may exist.

In summary, there are increasing options for scientists and athletes to monitor sleep, with some devices (primarily consumer sleep technology) having some limitations in their assessment of sleep and wakefulness. Consideration should be given to whether or not the devices are validated, and an assessment is required as to the usefulness and appropriateness of utilising unvalidated technology. Some devices may provide some basic level of sleep awareness and education; however, individuals should be cognisant of the limitations of available means of monitoring sleep. Table 1 provides current information on sleep monitoring tools with respect to their validation for detecting sleep.

REFERENCES

1. Carskadon MA, Dement WC. Monitoring and staging human sleep. In: Kryger MH, Roth T, Dement WC, eds. Principles and practice of sleep medicine. St. Louis, MO: Elsevier Saunders 2011:16-26.
2. Siegel JM. Clues to the functions of mammalian sleep. *Nature* 2005;437:1264-71. doi: 10.1038/nature04285
3. Zielinski MR, McKenna JT, McCarley RW. Functions and mechanisms of sleep. *AIMS Neurosci* 2016;3:67-104. doi: 10.3934/Neuroscience.2016.1.67
4. Berry RB, Brooks R, Gamaldo CE, *et al.* The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications. Darien, IL: American Academy of Sleep Medicine 2017.
5. Rechtschaffen A, Kales A. A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects. Washington DC: US Government Printing Office 1968.
6. Nielsen TA. A review of mentation in REM and NREM sleep: "covert" REM sleep as a possible reconciliation of two opposing models. *Behav Brain Sci* 2000;23:851-66; discussion 904-1121.
7. Ohayon MM, Carskadon MA, Guilleminault C, *et al.* Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep* 2004;27:1255-73. doi: 10.1093/sleep/27.7.1255

8. Banks S, Dinges DF. Chronic sleep deprivation. In: Kryger MH, Roth T, Dement WC, eds. Principles and practice of sleep medicine. St. Louis, MO: Elsevier Saunders 2011:67-75.
9. Dinges DF, Pack F, Williams K, *et al.* Cumulative sleepiness, mood disturbance, and psychomotor vigilance performance decrements during a week of sleep restricted to 4-5 hours per night. *Sleep* 1997;20:267-77.
10. Belenky G, Wesensten NJ, Thorne DR, *et al.* Patterns of performance degradation and restoration during sleep restriction and subsequent recovery: a sleep dose-response study. *J Sleep Res* 2003;12:1-12.
11. van Dongen H, Sha A, de Bruinier A, *et al.* Behavioural alertness and the two-process model of sleep regulation during chronic partial sleep deprivation. *J Sleep Res* 2002;11:233.
12. Walker MP, Stickgold R. It's practice, with sleep, that makes perfect: implications of sleep-dependent learning and plasticity for skill performance. *Clin Sports Med* 2005;24:301-17, ix. doi: 10.1016/j.csm.2004.11.002
13. Diekelmann S, Born J. The memory function of sleep. *Nat Rev Neurosci* 2010;11:114-26. doi: 10.1038/nrn2762
14. Walker MP, Stickgold R. Overnight alchemy: sleep-dependent memory evolution. *Nat Rev Neurosci* 2010;11:218; author reply 18. doi: 10.1038/nrn2762-c1
15. Fischer S, Hallschmid M, Elsner AL, *et al.* Sleep forms memory for finger skills. *Proc Natl Acad Sci U S A* 2002;99:11987-91. doi: 10.1073/pnas.182178199

16. Haack M, Mullington JM. Sustained sleep restriction reduces emotional and physical well-being. *Pain* 2005;119:56-64. doi: 10.1016/j.pain.2005.09.011
17. Orton DI, Gruzelier JH. Adverse changes in mood and cognitive performance of house officers after night duty. *BMJ* 1989;298:21-3. doi: 10.1136/bmj.298.6665.21
18. Caldwell JA, Caldwell JL, Brown DL, *et al.* The effects of 37 hours of continuous wakefulness on the physiological arousal, cognitive performance, self-reported mood, and simulator flight performance of f-117A pilots. *Mil Psychol* 2004;16:163-81.
19. Kahn-Greene ET, Killgore DB, Kamimori GH, *et al.* The effects of sleep deprivation on symptoms of psychopathology in healthy adults. *Sleep Med* 2007;8:215-21. doi: 10.1016/j.sleep.2006.08.007
20. Czeisler CA, Klerman EB. Circadian and sleep-dependent regulation of hormone release in humans. *Recent Prog Horm Res* 1999;54:97-130; discussion 30-2.
21. Krueger JM, Majde JA, Rector DM. Cytokines in immune function and sleep regulation. *Handb Clin Neurol* 2011;98:229-40. doi: 10.1016/B978-0-444-52006-7.00015-0
22. Moller-Levet CS, Archer SN, Bucca G, *et al.* Effects of insufficient sleep on circadian rhythmicity and expression amplitude of the human blood transcriptome. *Proc Natl Acad Sci U S A* 2013;110:E1132-41. doi: 10.1073/pnas.1217154110
23. Lange T, Perras B, Fehm HL, *et al.* Sleep enhances the human antibody response to hepatitis A vaccination. *Psychosom Med* 2003;65:831-5. doi: 10.1097/01.psy.0000091382.61178.f1

24. Spiegel K, Sheridan JF, Van Cauter E. Effect of sleep deprivation on response to immunization. *JAMA* 2002;288:1471-2.
25. Cohen S, Doyle WJ, Alper CM, *et al.* Sleep habits and susceptibility to the common cold. *Arch Intern Med* 2009;169:62-7. doi: 10.1001/archinternmed.2008.505
26. Horne JA. Why we sleep: The functions of sleep in humans and other mammals. Oxford, UK: Oxford University Press 1988.
27. Buxton OM, Pavlova M, Reid EW, *et al.* Sleep restriction for 1 week reduces insulin sensitivity in healthy men. *Diabetes* 2010;59:2126-33. doi: 10.2337/db09-0699
28. Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999;354:1435-9. doi: 10.1016/S0140-6736(99)01376-8
29. Markwald RR, Melanson EL, Smith MR, *et al.* Impact of insufficient sleep on total daily energy expenditure, food intake, and weight gain. *Proc Natl Acad Sci U S A* 2013;110:5695-700. doi: 10.1073/pnas.1216951110
30. Spiegel K, Tasali E, Penev P, *et al.* Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 2004;141:846-50. doi: 10.7326/0003-4819-141-11-200412070-00008
31. Meier-Ewert HK, Ridker PM, Rifai N, *et al.* Effect of sleep loss on C-reactive protein, an inflammatory marker of cardiovascular risk. *J Am Coll Cardiol* 2004;43:678-83. doi: 10.1016/j.jacc.2003.07.050

32. van Leeuwen WM, Hublin C, Sallinen M, *et al.* Prolonged sleep restriction affects glucose metabolism in healthy young men. *Int J Endocrinol* 2010;2010:108641. doi: 10.1155/2010/108641
33. Yaggi HK, Araujo AB, McKinlay JB. Sleep duration as a risk factor for the development of type 2 diabetes. *Diabetes Care* 2006;29:657-61. doi: 10.2337/diacare.29.03.06.dc05-0879
34. Cappuccio FP, Taggart FM, Kandala NB, *et al.* Meta-analysis of short sleep duration and obesity in children and adults. *Sleep* 2008;31:619-26. doi: 10.1093/sleep/31.5.619
35. Sabanayagam C, Shankar A. Sleep duration and cardiovascular disease: results from the National Health Interview Survey. *Sleep* 2010;33:1037-42. doi: 10.1093/sleep/33.8.1037
36. Grandner MA, Hale L, Moore M, *et al.* Mortality associated with short sleep duration: The evidence, the possible mechanisms, and the future. *Sleep Med Rev* 2010;14:191-203. doi: 10.1016/j.smr.2009.07.006
37. Halson SL. Sleep monitoring in athletes: motivation, methods, miscalculations and why it matters. *Sports Med* 2019;49:1487-97. doi: 10.1007/s40279-019-01119-4
38. Roomkham S, Lovell D, Cheung J, *et al.* Promises and challenges in the use of consumer-grade devices for sleep monitoring. *IEEE Rev Biomed Eng* 2018;11:53-67. doi: 10.1109/RBME.2018.2811735
39. Finan PH, Richards JM, Gamaldo CE, *et al.* Validation of a wireless, self-application, ambulatory electroencephalographic sleep monitoring device in healthy volunteers. *J Clin Sleep Med* 2016;12:1443-51. doi: 10.5664/jcsm.6262

40. Quante M, Kaplan ER, Rueschman M, *et al.* Practical considerations in using accelerometers to assess physical activity, sedentary behavior, and sleep. *Sleep Health* 2015;1:275-84. doi: 10.1016/j.sleh.2015.09.002
41. Kolla BP, Mansukhani S, Mansukhani MP. Consumer sleep tracking devices: a review of mechanisms, validity and utility. *Expert Rev Med Devices* 2016;13:497-506. doi: 10.1586/17434440.2016.1171708
42. Ancoli-Israel S, Martin JL, Blackwell T, *et al.* The SBSM guide to actigraphy monitoring: Clinical and research applications. *Behav Sleep Med* 2015;13 Suppl 1:S4-S38. doi: 10.1080/15402002.2015.1046356
43. Depner CM, Cheng PC, Devine JK, *et al.* Wearable technologies for developing sleep and circadian biomarkers: a summary of workshop discussions. *Sleep* 2019 doi: 10.1093/sleep/zsz254
44. Sargent C, Lastella M, Halson SL, *et al.* The validity of activity monitors for measuring sleep in elite athletes. *J Sci Med Sport* 2016;19:848-53. doi: 10.1016/j.jsams.2015.12.007
45. Fuller KL, Juliff L, Gore CJ, *et al.* Software thresholds alter the bias of actigraphy for monitoring sleep in team-sport athletes. *J Sci Med Sport* 2017;20:756-60. doi: 10.1016/j.jsams.2016.11.021
46. Ancoli-Israel S, Cole R, Alessi C, *et al.* The role of actigraphy in the study of sleep and circadian rhythms. *Sleep* 2003;26:342-92. doi: 10.1093/sleep/26.3.342
47. Hamill K, Jumabhoy R, Kahawage P, *et al.* Validity, potential clinical utility and comparison of a consumer activity tracker and a research-grade activity tracker in

- insomnia disorder II: Outside the laboratory. *J Sleep Res* 2019:e12944. doi: 10.1111/jsr.12944
48. Kahawage P, Jumabhoy R, Hamill K, *et al.* Validity, potential clinical utility, and comparison of consumer and research-grade activity trackers in Insomnia Disorder I: In-lab validation against polysomnography. *J Sleep Res* 2019:e12931. doi: 10.1111/jsr.12931
49. De Zambotti M, Cellini N, Goldstone A, *et al.* Wearable sleep technology in clinical and research settings. *Med Sci Sports Exerc* 2019;51:1538-57. doi: 10.1249/MSS.0000000000001947
50. Evenson KR, Goto MM, Furberg RD. Systematic review of the validity and reliability of consumer-wearable activity trackers. *Int J Behav Nutr Phys Act* 2015;12:159. doi: 10.1186/s12966-015-0314-1
51. Sargent C, Lastella M, Romyn G, *et al.* How well does a commercially available wearable device measure sleep in young athletes? *Chronobiol Int* 2018;35:754-58. doi: 10.1080/07420528.2018.1466800
52. Russo K, Goparaju B, Bianchi MT. Consumer sleep monitors: is there a baby in the bathwater? *Nat Sci Sleep* 2015;7:147-57. doi: 10.2147/NSS.S94182
53. Kortelainen JM, Mendez MO, Bianchi AM, *et al.* Sleep staging based on signals acquired through bed sensor. *IEEE Trans Inf Technol Biomed* 2010;14:776-85. doi: 10.1109/TITB.2010.2044797

54. Tuominen J, Peltola K, Saaresranta T, *et al.* Sleep parameter assessment accuracy of a consumer home sleep monitoring ballistocardiograph beddit sleep tracker: a validation study. *J Clin Sleep Med* 2019;15:483-87. doi: 10.5664/jcsm.7682
55. Schade MM, Bauer CE, Murray BR, *et al.* Sleep validity of a non-contact bedside movement and respiration-sensing device. *J Clin Sleep Med* 2019;15:1051-61. doi: 10.5664/jcsm.7892
56. Fino E, Mazzetti M. Monitoring healthy and disturbed sleep through smartphone applications: a review of experimental evidence. *Sleep Breath* 2018 doi: 10.1007/s11325-018-1661-3
57. Choi YK, Demiris G, Lin SY, *et al.* Smartphone applications to support sleep self-management: Review and evaluation. *J Clin Sleep Med* 2018;14:1783-90. doi: 10.5664/jcsm.7396
58. Lorenz CP, Williams AJ. Sleep apps: what role do they play in clinical medicine? *Curr Opin Pulm Med* 2017;23:512-16. doi: 10.1097/MCP.0000000000000425
59. Anderson KN. Insomnia and cognitive behavioural therapy-how to assess your patient and why it should be a standard part of care. *J Thorac Dis* 2018;10:S94-S102. doi: 10.21037/jtd.2018.01.35
60. Carney CE, Buysse DJ, Ancoli-Israel S, *et al.* The consensus sleep diary: standardizing prospective sleep self-monitoring. *Sleep* 2012;35:287-302. doi: 10.5665/sleep.1642
61. Buysse DJ, Reynolds CF, 3rd, Monk TH, *et al.* The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193-213.

62. Mastin DF, Bryson J, Corwyn R. Assessment of sleep hygiene using the Sleep Hygiene Index. *J Behav Med* 2006;29:223-7. doi: 10.1007/s10865-006-9047-6
63. Parrott AC, Hindmarch I. The Leeds Sleep Evaluation Questionnaire in psychopharmacological investigations - a review. *Psychopharmacology (Berl)* 1980;71:173-9. doi: 10.1007/bf00434408
64. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991;14:540-5.
65. Bender AM, Lawson D, Werthner P, *et al.* The clinical validation of the athlete sleep screening questionnaire: an instrument to identify athletes that need further sleep assessment. *Sports Med Open* 2018;4:23. doi: 10.1186/s40798-018-0140-5
66. Samuels C, James L, Lawson D, *et al.* The athlete sleep screening questionnaire: a new tool for assessing and managing sleep in elite athletes. *Br J Sports Med* 2016;50:418-22. doi: 10.1136/bjsports-2014-094332
67. Driller MW, Mah CD, Halson SL. Development of the athlete sleep behavior questionnaire: A tool for identifying maladaptive sleep practices in elite athletes. *Sleep Sci* 2018;11:37-44. doi: 10.5935/1984-0063.20180009

Figure Legend

Supplementary Figure 1. Examples of sleep stages (hypnograms) in healthy, young adults spending 9 hours in bed – showing wake, rapid eye movement sleep (REM), and the three stages of non-rapid eye movement sleep (non-REM), i.e., N1, N2, and N3. Panel A shows a normal sleep – it has alternating periods of non-REM and REM, most deep sleep (N3) occurs in the first third of the night, and most REM occurs in the last third of the night. Panels B, C, and D show sleep with longer than normal periods of wake at the start, middle, and end, respectively.