

## Online Supplementary Figure 1

## Frequency (%) time loss ARill by anatomical classification: Upper ARill vs. General (upper/lower) ARill

Studies	Estimate (95% C.I.)	Ev/Trt
Alonso et al. (2012)	0.217 (0.049, 0.386)	5/23
Dressendorfer et al. (2002)	0.200 (-0.151, 0.551)	1/5
Edouard et al. (2013)	0.125 (-0.104, 0.354)	1/8
Edouard et al. (2014)	0.444 (0.120, 0.769)	4/9
Edouard et al. (2015)	0.500 (0.154, 0.846)	4/8
Orhant et al. (2010)a	0.113 (0.062, 0.163)	17/151
Schwellnus et al. (2012)b	0.181 (0.092, 0.269)	13/72
<b>Subgroup Upper ARill (I<sup>2</sup>=41.95 % , P=0.111)</b>	<b>0.184 (0.107, 0.261)</b>	<b>45/276</b>
Derman et al. (2014)	0.125 (0.072, 0.178)	19/152
Nordstrom et al. (2020)	0.351 (0.275, 0.427)	53/151
Schwellnus et al. (2012)a	0.192 (0.041, 0.344)	5/26
Soligard et al. (2017)	0.185 (0.140, 0.229)	54/292
Steffen et al. (2019)	0.220 (0.158, 0.283)	37/168
<b>Subgroup General (Upper/Lower) ARill (I<sup>2</sup>=83.15 % , P=0.000)</b>	<b>0.213 (0.141, 0.285)</b>	<b>168/789</b>
<b>Overall (I<sup>2</sup>=71.32 % , P=0.000)</b>	<b>0.203 (0.151, 0.255)</b>	<b>213/1065</b>

